

Available online on 15.9.2016 at <http://ujpr.org>**Universal Journal of Pharmaceutical Research***An International Peer Reviewed Journal*

Open access to Pharmaceutical research

This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial Share Alike 4.0 License which permits unrestricted non commercial use, provided the original work is properly cited

**Volume 1, Issue 1, 2016**

Open Access

RESEARCH ARTICLE

ANTIFUNGAL, CYTOTOXIC AND PHYTOTOXICITY OF AERIAL PART OF *RANUNCULUS MURICATUS*

Muhammad Shahzad Aslam¹, Rubina Rehman², Bashir A. Choudhary², Abdul Subhan Ijaz², Muhammad Uzair², Muhammad Syarhabil Ahmad¹

¹School of Bioprocess Engineering, University Malaysia Perlis, Kompleks Pusat Pengajian, Jejawi 3, 02600 Arau, Perlis, Malaysia.

²Department of Pharmacy, Bahauddin Zakariya University, Multan, Pakistan.

ABSTRACT

Objectives: The present research is preliminary biological screening of aerial plant of *Ranunculus muricatus* (Ranunculaceae). Dichloromethane and methanol extracts of the aerial plant were investigated for their antifungal, phytotoxic and cytotoxic activities.

Methods: Anti-fungal, cytotoxicity and phytotoxicity activities were performed by agar tube dilution assay, brine shrimp lethality bioassay and lemna bioassay respectively. Dichloromethane and methanolic extracts exhibited significant phytotoxicity against Lemna minor having Paraquat as standard drug and incubation condition (28±1°C).

Results: None of extracts presented any significant cytotoxic activity having Imipenem and Etoposide as standard drug respectively. Both extract had non-significant antifungal activity but it has been noted that methanol extract showed 30% inhibition with linear growth at 70 mm, when compared with control; only against *Microsporum canis*. *Ranunculus muricatus* showed significant phytotoxicity.

Conclusion: The phytotoxicity assay is a valuable major screen for weedicide investigation. Additionally, modern studies are currently carried out to identify the allelopathic constituents by isolation, purification and structure elucidation to find out as effective herbicidal.

Keywords: Biological screening; cytotoxicity, phytotoxicity, *Ranunculus muricatus*.

Article Info: Received 9 September 2016; Revised 16 October; Accepted 30 October, Available online 15 November 2016

**Cite this article-**

Aslam MS, Rehman R, Choudhary BA, Ijaz AS, Uzair M, Ahmad MS. Antifungal, cytotoxic and phytotoxicity of aerial part of *Ranunculus muricatus*. Universal Journal of Pharmaceutical Research 2016; 1(1): 5-7.

DOI: <http://doi.org/10.22270/ujpr.v1i1.R2>

Address for Correspondence:

Muhammad Shahzad Aslam, 1School of Bioprocess Engineering, University Malaysia Perlis, Kompleks Pusat Pengajian, Jejawi 3, 02600 Arau, Perlis, Malaysia, Tel: +6019-3009674, E-mail: aslammuhammadshahzad@gmail.com

INTRODUCTION

Ranunculus muricatus also known as *Ranunculus pseudo-muricatus* Baltter and Hallb is indigenous to Atlantic, S. Europe, W. and S. W. Asia, Crimea, Caucasus, S. Siberia, Pakistan and India. Flowering period is between March and April¹. It is recognised with different names in different part of world such as spiny buttercup (English); Chambul, jaghagha, Latokari, Korgandal (Folk). This plant is slightly poisonous. Whole plant is traditionally used as decoction for periodic fever and asthma². The plant reported to contain stigmasterol-4-ene-3, 6-dione, stigmasterol, Anemonin, Aescin lactone dimethyl ether, beta-valley sterol, protocatchuic aldehyde, protocatchuic acid, and luteolin factors³. The findings of anemonin as chemical constituents in *Ranunculus muricatus* may justify the uses of these species against fever, rheumatism and rubefacient in Asian traditional

medicines⁴. Present study is conducted to document the *in vitro* biological activities of *Ranunculus muricatus* conducted to document the *in vitro* biological activities of *Ranunculus muricatus*. Antifungal, phytotoxic and cytotoxic bioassay has studied.

METHODS**Plant material**

Ranunculus muricatus was collected from Jallo pind, Lahore. Total weight of wet plant collected was 15kg while 5kg of dried plant was obtained after drying. The plant was identified by Dr. Altaf Hussain Dasti, Professor, Institute of pure and applied Biology, Bahauddin Zakariya University, Multan.

Extraction

The air-dried aerial part of plant material was grounded and extracted successively with dichloromethane and

methanol (thrice with each solvent) at room temperature occasional shaking for 24hrs. Extracts were concentrated by Rotavapor-R20 at 35°C.

Anti-fungal bioassay

Test fungi such as *Candida albicans*, *Aspergillus flavus*, *Microsporium canis*, *Fusarium solani*, *Candida glabrata* were employed for preliminary screening. Extracts were dissolved in sterile DMSO to serve as stock solution. Sabouraud dextrose agar was prepared by mixing Sabouraud 4% glucose agar and agar in distilled water. Known amount of media was dispensed into screw capped test tubes. Test tubes containing

media were autoclaved 121°C for 15 minutes. Tubes were allowed to cool to 50°C and the desired concentration of extract was added into non-solidified media. The tubes were allowed to solidify at room temperature. Each tube was inoculated with a 4mm diameter piece of in culture of fungi. All culture-containing tubes were inoculated at optimum temperature of 28-30°C for growth for 7 to 10 days. Culture was examined at least twice a weekly during the incubation. With no visible growth of microorganism is taken to represent the MIC of the test sample which is expressed in µg⁵.

Table 1: Results of *in-vitro* antifungal bioassay of *Ranunculus muricatus*

Extract	Name of Fungus	Linear Growth (mm)		% Inhibition	Standard Drug	Mic (µg/ml)
		Sample	Control			
MeOH	<i>Candida albicans</i>	100	100	0	Miconazole	110.8
	<i>Aspergillus flavus</i>	100	100	0	Amphotericin B	20.20
	<i>Microsporium canis</i>	70	100	30	Miconazole	98.4
	<i>Fusarium solani</i>	100	100	0	Miconazole	73.25
	<i>Candida glabrata</i>	100	100	0	Miconazole	110.8
DCM	<i>Candida albicans</i>	100	100	0	Miconazole	110.8
	<i>Aspergillus flavus</i>	100	100	0	Amphotericin B	20.20
	<i>Microsporium canis</i>	100	100	0	Miconazole	98.4
	<i>Fusarium solani</i>	100	100	0	Miconazole	73.25
	<i>Candida glabrata</i>	100	100	0	Miconazole	110.8

Table 2: Results of *in vitro* phytotoxic bioassay of *Ranunculus muricatus*

Extract	Plant Name	Conc. of Compound (µg/ml)	No. of Fronds		% Growth Regulation	Conc. of Standard Drug (µg/ml)
			Sample	Control		
MeOH	<i>Lemna minor</i>	1000	0		100	0.015
		100	9	20	55	
		10	17		15	
		1000	0		100	
DCM	<i>Lemna minor</i>	100	9	20	55	0.015
		10	17		15	
		1000	0		100	

Phytotoxicity bioassay

Prepared inorganic medium of 5.5-6.0 pH attained with KOH pellets. 10 vials per dose 500, 50, 5 and control were prepared. 15mg of the extract was dissolved in 15ml of the solvent. 1000, 100 and 10µl of solution to vials for testing allow the solvent to evaporate overnight. 2ml of medium was added in each vial containing a single plant a rosette of three fronds. The vials were placed in a glass dish filled with 2cm of water, sealed the container with stopcock grease and glass plate. Placed the dish along with vials in growth chamber for seven days at 25°C under fluorescent and incandescent light. Count the number of fronds per vials on day 3 and 7. Analyzed the data as percent of control with ED 50 computer program⁵.

Brine shrimp lethality bioassay

Brine shrimp cytotoxicity assay was accomplished according to the standard procedure described by McLaughlin⁶. Three concentrations (1000, 100, and 10 ppm) of the plant extracts were used in this assay. Brine shrimp larvae were hatched in a small partitioned tank in artificial seawater. Illumination was provided on one side to attract newly hatched larvae. Brine

shrimp larvae with second in star stage were used in this assay. Plant extracts of respective concentrations were added to dram vials. To each dram vial ten brine shrimp larvae were added. Negative control was prepared by evaporating 0.5ml of methanol in dram vials and then by adding sea salt solution to it. Following 24 hrs of incubation, survivors were counted by using magnifying glass. The experiment was repeated three times. Mortality data was transformed by probit analysis in finny computer program to estimate ED50 value. Percentage of mortality was also calculated at all concentrations⁶.

RESULTS AND DISCUSSION

Dichloromethane and methanol extracts of the aerial plant of *Ranunculus muricatus* were studied for their antifungal, phytotoxic and Brine Shrimp lethality bioassay. *Candida albicans*, *Aspergillus flavus*, *Microsporium canis*, *Fusarium solani* and *Candida glabrata* were employed for fugitoxic effect of the extracts. It has been noted that MeOH extract of *Ranunculus muricatus* showed 30% inhibition with

linear growth at 70mm, as compared with control; only against *Microsporium canis* at the concentration of 400µg/ml for incubation period of seven days at 27°C with reference to Miconazole as standard. While dichloromethane extract does not showed any activity as shown in Table 1. Dichloromethane and methanolic extracts of the aerial part of *Ranunculus muricatus*

showed significant phytotoxicity at concentrations of 1000 µg/ml, 100 µg/ml and 10 µg/ml against *Lemna minor*. Dichloromethane and methanolic extract of *Ranunculus muricatus* does not showed cytotoxicity even at highest level having Etoposide as standard drug containing 28±1°C as incubation condition.

Table 3: Results of *in-vitro* cytotoxic bioassay of *Ranunculus muricatus*

Extract	Dose (µg/ml)	No. of Shrimp	No. of Survivors	LD 50 (µg/ml)	STD Drug	LD 50 (µg/ml)
MeOH	1000	30	14	857.73	Etoposide	7.4625
	100	30	25			
	10	30	29			
DCM	1000	30	24	45456.4	Etoposide	7.4625
	100	30	28			
	10	30	29			

CONCLUSION

The phytotoxicity assay is a valuable major screen for weedicide investigation. Weeds are one of the main issues of poor agronomic efficiency in the developing countries. Expensive, lethal and non-specific synthetic pesticides are used now. Using natural sources pesticides will not only reduces the adverse effect of pesticides but also help us to return to natural flora. Natural herbicides destroy specific targets, while leaving the wanted crop comparatively undamaged. Additionally, modern studies are currently carried out to discover the phytotoxic constituents of the plant by isolation, purification and structure elucidation to find out as effective herbicidal.

ACKNOWLEDGEMENTS

The authors are grateful for the support provided by the department of Pharmacy Bahauddin Zakariya University, Multan and School of Bioprocess Engineering, University Malaysia Perlis, Malaysia. We also wish to acknowledge the technical support of HEJ Research institute of Chemistry, University of Karachi, Karachi, Pakistan.

AUTHOR'S CONTRIBUTION

The manuscript was carried out, written, and approved in collaboration with all authors.

CONFLICT OF INTERESTS

Declared none.

REFERENCES

- Nasir E, Ali SI. Flora of Pakistan 1986: No.172.
- Iqbal H, Sher Z, Khan ZU. Medicinal plants from Salt range Pind Dadan Khan, district Jhelum, Punjab, Pakistan. J Med Plants Research. 5(11): 2157-2168.
- Lingjie W, Xiao-Zhong G. The chemical composition of the fruit butter cup. China Modern Applied Pharmacy. 2009; 6:460-462.
- Aslam MS, Choudhary BA, Uzair M, Ijaz AS, The genus *Ranunculus*: A Phytochemical and Ethnopharmacological Review. Int J Pharm Pharmaceu Sci. 2012; 4(5): 15-22.
- Rehman A, Choudhary MI and Thomsen WJ. Bioassay technique for drug development Harwood Academic Publishers. 2001: 16-17, 23-24, 65-66.
- McLaughlin JL. Crown gall tumors on potato discs and brine shrimp lethality: two single bioassays for plant screening and fraction. In: Hostetsmann, K, ed., Methods in Plant Biochemistry. Academic Press, London. 1991. 1-31. PMID: 23675023