Available online on 15.03.2019 at <http://ujpr.org>
Universal Journal of Pharmaceutical Research
 An International Peer Reviewed Journal

Open access to Pharmaceutical research
 This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial Share Alike 4.0 License which permits unrestricted non commercial use, provided the original work is properly cited

Volume 4, Issue 1, 2019

Open Access

RESEARCH ARTICLE

STUDY ON FRESH LEAF AQUEOUS EXTRACT OF *FLACOURTIA INDICA* FOR HEPATOPROTECTIVE, ANTI-ANEMIC AND HYPOGLYCEMIC ABILITIES IN CCl₄ INDUCED HEPATOTOXICITY IN ALBINO WISTAR RATS

Idoko A*^{ID}, Ufedo-Enyo G. Emmanuel^{ID}

Department of Biochemistry, Faculty of Natural Sciences, Caritas University, Amorji– Nike, P.M.B. 01784, Enugu, Nigeria.

ABSTRACT

Objective: Hepatic injury and its associated conditions have been reportedly shown to be managed through herbal remedies. In this study, investigation of the fresh leaf aqueous extract of *Flacourtia indica* (of the family of *Salicaceae*) as hypoglycemic, anti-anemic and hepatoprotective agent in albino wistar rats induced CCl₄ hepatotoxicity was done.

Methods: Fifteen rats of either sex, weighing 175-295g, divided into five groups (I-V) of three rats each, were used. Group-I is negative control, II-positive control and III –V test groups. Groups II-V were induced 200mg/Kg/bodyweight CCl₄, for 3-days, for hepatic injury and anemia. Groups III-V were respectively treated orally with 400, 600 and 800 mg/Kg/bodyweight of fresh leaf aqueous extracts (FLAE) of *Flacourtia indica*, for 7-days.

Results: Activities of alanine aminotransferase, aspartate aminotransferase, alkaline phosphatase, concentrations of bilirubin, albumin, total protein, blood glucose and packed cell volume (PCV) and hemoglobin were assayed. Results after induction showed significant ($p < 0.05$) decrease in hemoglobin and PCV, significant ($p < 0.05$) increase in the liver function enzymes and blood glucose compared with results of liver function enzyme and blood glucose having significant ($p < 0.05$) decrease, and significant ($p < 0.05$) increase of PCV and hemoglobin after treatment with FLAE of *Flacourtia indica*. Body weight of rats induced CCl₄ was found to increase with FLAE of *Flacourtia indica* treatment.

Conclusion: It may be concluded that the potentials exhibited by FLAE of *Flacourtia indica* to manage hyperglycaemia, hepatic injury and anemia induced by CCl₄ are associated with its antioxidant activity and the presence of phytochemicals, minerals and nutrients value.

Keywords: Anti-anemic, *Flacourtia indica*, hepatoprotective, hepato-function, hypoglycemia, toxicity.

Article Info: Received 4 January 2019; Revised 5 February; Accepted 6 March, Available online 15 March 2019

Cite this article-

Idoko A, Ufedo-Enyo G. Emmanuel. Study on fresh leaf aqueous extract of *flacourtia indica* for hepatoprotective, anti-anemic and hypoglycemic abilities in CCl₄ induced hepatotoxicity in albino wistar rats. Universal Journal of Pharmaceutical Research 2019; 4(1): 17-22.

DOI: <https://doi.org/10.22270/ujpr.v4i1.234>

Address for Correspondence:

Idoko A, Department of Biochemistry, Faculty of Natural Sciences, Caritas University, Amorji – Nike, P.M.B. 01784, Enugu, Nigeria. Phone: +2348032354823, E-mail: idokoalexander1@gmail.com

INTRODUCTION

Plants have been largely used in the quest of managing health challenges as alternative of nature's providence. Traditional medicine practice is an age long practice common in developing countries¹. Several bioactive compounds in plants are known with their antioxidant and scavenging abilities². Phytochemicals are chemicals in plants with no nutritive value but highly effective in disease prevention and protection when consumed. Most of these bioactive plants' compounds have been implicated in the treatment, management and prevention of ailments. Several studies have demonstrated the use of plants' extracts as hepatoprotective and gluco-stabilizer in Albino wistar rats induced aluminium chloride hepatic toxicity³ and⁴ reported various methods of extraction, isolation,

identification and purification of bioactive compounds in plants. *Flacourtia indica*'s leaf, stem bark, fruits and root like other plants, is not exempted from these beneficial characteristics. These have been shown to possess biological, medicinal and pharmacological potentials in the prevention and treatment of hepatic disease³, cardiovascular diseases, cancer⁵, diabetes⁶, bacterial infection¹, and other conditions like anemia hyperglycemia and hypercholesterolemia^{7,8}. Various researchers have demonstrated that plants are rich sources of antioxidant vitamins such as vitamins A, C and E⁴, minerals such as Fe, Mg, Mn, N, P, Ca, Na and K⁹ and phytochemicals such as phenolics (tannins, flavonoids), carotenoids, anthocyanins, coumarin glycosides¹. The liver is an organ with multiple functions. It is involved in circulation of blood, plays

major role in metabolic reactions, seen in conversion of excess blood glucose to glycogen, carries out detoxification by secretion of bile, involved in production of blood clotting factor by production of fibrinogen, heparin and prothrombin^{10,11}. Hepatic diseases pose a universal concern to Humans and other animals, contributing a large cause of mortality and morbidity. These include fatty liver, cirrhosis; hepatitis (A, B, C, D and E), drug/chemical induced hepatic injury, hepatic cancer and alcohol induced hepatic injury¹¹. Chemical/drug induced liver toxicity is reported to be the paramount cause of hepatotoxicity. This has been linked to life style, abuse and misuse of drug, occupational, laboratory and industrial exposure to substances and chemicals like carbon tetrachloride, aluminum chloride, alcohol etc¹⁰. The mechanism through which carbon tetrachloride CCl₄ and these other chemical substances exert liver damage is understood to be linked to production of reactive oxygen species. This result in lipid peroxidation of liver tissues as a consequence of the high put of free radicals generated which subdues the liver's defense system, degenerating to inflammation, hepatic apoptosis, liver cirrhosis and fibrosis^{12,13}.

Anemia sets in due to lack of adequate and healthy red blood cells (RBC) and hemoglobin, the oxygen binding component of the blood. Anemia is a condition that is commonly affected by infants, child bearing age women/pregnant women, the young and the elderly¹⁴. Different types of anemia arise from their causes. Anemia is considered to be caused by abnormal RBC production (iron deficiency anemia, vitamin deficiency anemia, aplastic anemia thalassemia etc), destruction of red blood cell (sickle cell anemia¹⁵). Anemia has been reported to be induced by several chemical substances such as AlCl₃ and phenylhydrazine^{14,16}, reported the prevalence of anemia among the elderly with value of hemoglobin Hb < 12g/dl in women and Hb <13g/dL in men.

This study was carried out to evaluate the potential of fresh leaf aqueous extracts (FLAE) of *F. indica* as a hepatoprotective, anti-anemic and hypoglycemic agent in CCl₄ induced hepatic injury in Albino wistar rats.

MATERIALS AND METHODS

Collection and Preparation of Plant Samples

Fresh leaf materials of *F. indica* (Governor's plum) were collected from around staff quarter of Caritas University, Amorji-Nike, Enugu state, Nigeria. The required plant leaf was authenticated and a voucher number of PSB/109-12.A was given by Mr. Okafor, C.U., a botanist in plant tissue culture and biotechnology department, Faculty of Biological Science, University of Nigeria, Nsukka. The aqueous plant extracts were prepared selecting fresh leaf aerial part, weighed and squeezed in a bowl of containing water and filtered and filtrate was used for oral treatment. The volumes of the extracts to be administered were calculated according to the body weight of the rats.

Collection and Preparation of Blood Sample

Three milliliter (3ml) of blood was collected from the rats by capillary pressure insertion into the side of the

eye using capillary tubes into a plain bottle, for the collection of serum used for biochemical assay (liver function test) and about 3mls collected in an EDTA sample bottle for hematological assay (PCV and hemoglobin). The samples in bottles were stored at room temperature.

Study Animals

Albino Wistar rats of 175-294g weight, of either sex were obtained from university of Nigeria Nsukka. Animals were housed at an ambient temperature and relative humidity in the animals' house of department of Biochemistry, natural sciences, Caritas University, Amorji-Nike Enugu. The rats were allowed to acclimatize for one week prior to the experiment and had access to standardized pelletized finisher feed and clean water within the period of the acclimatization. The principle of laboratory animals' care and ethical guidelines for investigation of experimental pain in conscious animals were followed respectively^{17,18}.

Design and Animal Grouping

A total of fifteen (15) Wistar albino rats, divided into five groups (Groups I–V) of three rats each was used for this study.

Group I: Negative control consist of 3 rats, no carbon tetrachloride CCl₄ and FLAE of *F. indica* were administered.

Group II: Test control (positive control) consist of 3 rats, were administered orally with 200mg/Kg/body weight CCl₄ without FLAE of *F. indica*.

Group III: Consist of 3 rats, administered orally with 200mg/Kg/body weight CCl₄ and 400mg/kg/body weight FLAE of *F. indica*.

Group IV: Consist of 3 rats, administered orally with 200mg/Kg/body weight CCl₄ and 600mg/kg/body weight FLAE of *F. indica*.

Group V: Consist of 3 rats, administered orally with 200mg/Kg/body weight CCl₄ and 800mg/kg/body weight FLAE of *F. indica*.

At the end of induction (three days), blood sample was collected from each group for biochemical and hematological assays before treatment with FLAE of *F. indica*. After treatment with FLAE of *Flacourtia indica* for seven days, blood sample was also collected for biochemical and hematological assays.

Induction of Liver Injury and Anemia

Rats of groups II–V were induced with liver injury and anemia by single oral administration with 200mg/kg body weight of CCl₄ respectively. A confirmatory test was carried out after induction of anemia by assaying the plasma hemoglobin percentage to show that the rats were anemic.

Liver Function Assay

After collection of blood sample from rats, serum was collected by clot retraction. Serum ALT, AST, ALP, Albumin, Total protein and Bilirubin were assayed by the standard method as described by¹⁹ with the use of kits from Randox Laboratories Ltd, 55 Diamond Road, Crumlin, country Antrim, BT29 4QY, United Kingdom.

Figure 1: Liver function test after induction with 200 mg/kg body weight CCl_4 .

Letters a and b indicates significant difference ($P < 0.05$) when group I was compared with groups II, III, IV and V, respectively after CCl_4 induction of liver damage to rats in these groups. Graphs with same letters are not significantly ($P < 0.05$) different.

Hematological Assay

The Haemoglobin (Hb) and packed Cell Volume (PCV) values were determined by standard method as described in a previous study²⁰ using hematocrite and Mindray Haematology Analyser (Mindray BC-2300, Guangzhou Shihai Medical Equipment Co., Ltd, China).

Chemicals

All chemicals used were pure and of analytical grade. Liver function enzymes assay reagents for Bilirubin (BIL), Alanine Aminotransferase (ALT), Aspartate Aminotransferase (AST), Albumin (ALB), Total Protein (TP) and Alkaline phosphatase (ALP) employed kits obtained from Randox Laboratories Ltd, 55 Diamond Road, crumlin, country Antrim, BT29 4QY, United Kingdom. Aluminium trichloride AlCl_3 was purchased from BDH Laboratories/Chemicals Ltd, Poole, England.

Figure 2: Liver function Assay for CCl_4 induced rats treated with fresh leaf aqueous extract of *F. indica*.

FLF. *indica* = fresh leaf of *F. indica*. Letters a, b, c, d and e indicates significant difference ($P < 0.05$) when group II was compared with groups I, III, IV and V, respectively for 400mg/kg, 600mg/kg and 800mg/kg FLAE *F. indica* treated groups.

Statistical Analysis

Results were expressed as mean $\pm$ standard deviation and analyzed using one-way ANOVA (analysis of variance), p value (< 0.05) was considered significant. A component of graph pad instat 3 software version

3.05 and graphpad prism version 7.04 by graph pad Inc. was employed²¹.

Figure 3: Alkaline phosphatase level of CCl_4 induced rat's liver damage.

Letters a, b, c, d and e indicates significant difference ($P < 0.05$) when group I was compared with groups II, III, IV and V, respectively after CCl_4

RESULTS

Table 1 shows the liver function parameters of rats after induction with 200mg CCl_4 for liver injury. There was an observed significant ($p < 0.05$) increase in TP, ALB, BIL, ALP, ALT and AST of group I (negative control) compared to group II (positive control) and test groups (III, IV and V). Table 2 shows liver function assay of rats administered various doses (400mg/kg, 600mg/kg and 800mg/kg) of Fresh Leaf Aqueous Extract (FLAE) of *F. indica* for seven (7) days. A significant ($p < 0.05$) increase was observed in all parameters assayed in the test groups (III, IV and V) compared to group II (test control) and group I (negative control). The packed cell volume (PCV) and hemoglobin of rats after induction with CCl_4 and after treatment with Fresh Leaf Aqueous Extract (FLAE) of *F. indica* for seven (7) days is shown in table 3. The result shows a significant ($p < 0.05$) increase in PCV and hemoglobin after treatment with FLAE of *F. indica* compared to after induction with CCl_4 .

Blood glucose concentration of rats after induction with CCl_4 and after Treatment with Fresh Leaf Aqueous Extract (FLAE) of *F. indica* for seven (7) days is shown in Table 4. After FLAE *F. indica* was administered, to test groups (III, IV and V), the blood glucose concentrations of test groups were observed to decrease significantly ($p < 0.05$) compared to after induction with CCl_4 and thus acerbating induced hypoglycemia. The body weights of rats are shown in Table 5 at acclimatization, at induction with CCl_4 and at treatment with FLAE of *F. indica*. There was significant ($p < 0.05$) decrease in body weight of at induction and significant increase in body weight after treatment with FLAE of *F. indica*, which may be indicative of recovery from anemia. The results show significantly higher ($P < 0.05$) level of group II rats (positive control) compared to group I (negative control), group III (400mg/kg FLF *indica*), group IV (600mg/kg FLF. *indica*) and group V (800mg/kg FLF *indica*).

Table 1: Liver function test of rats after induction with 200mg/kg CCl₄ body weight of rats.

Group	TP(g/dl)	ALB(g/dl)	BIL(mg/dl)	ALP(U/L)	ALT(U/L)	AST(U/L)
I	10.70±7.11 ^{abcd}	4.58±3.02 ^{abcd}	12.88±0.77 ^{abcd}	1381.67±71.12 ^{abcd}	38.7±5.33 ^{abcd}	41.00±6.44 ^{abcd}
II	7.49±5.48 ^a	6.56±7.13 ^a	9.49±1.02 ^a	2612.00±36.49 ^a	77.04±2.34 ^a	76.8±4.33 ^a
III	2.58±0.95 ^b	3.72±0.23 ^b	3.38±0.59 ^b	1453.14±419.60 ^b	68.00±24.37 ^b	75.00±10.36 ^b
IV	3.26±0.49 ^c	3.29±0.62 ^c	4.09±0.70 ^c	1927.81±99.46 ^c	79.50±20.51 ^c	78.00±15.56 ^c
V	4.23±1.78 ^d	3.75±0.04 ^d	4.11±1.39 ^d	1712.58±212.73 ^d	79.50±20.51 ^d	71.50±6.36 ^d

Results are mean ± standard deviation, Values in the same column bearing similar superscripts are significantly different at $P<0.05$. (n=3). Key: I: Negative Control Group, II: positive control and III, IV and V: Test groups. TP: Total Protein, ALB: Albumin, BIL: Bilirubin, ALP: Alkaline Phosphatase, ALT: Alanine Transaminase, AST: Aspartate Transaminase.

Table 2: Liver function assay of Rats after Treatment with 400, 600 and 800mg/kg/body weight of Fresh Leaf Aqueous Extract (FLAE) of *F. indica*.

Group	TP(g/dl)	ALB(g/dl)	BIL(mg/dl)	ALP(U/L)	ALT(U/L)	AST(U/L)
I	10.07±10.82 ^{abc}	4.73±1.22 ^a	8.98±1.01 ^{ab}	1701.01±173.23 ^{abc}	34.01±7.33 ^{abc}	35.02±7.24 ^{abc}
II	6.98±7.63 ^a	4.52±3.11	9.19±0.12 ^a	2034.01±84.12 ^a	67.00±2.08 ^a	72.19±7.12 ^a
III	2.25±1.88 ^b	2.34±0.97 ^a	1.54±0.90	1003.85±134.52 ^b	14.33±9.01 ^b	59.50±0.71 ^b
IV	2.75±0.45 ^c	1.38±0.58	2.15±2.14 ^b	1565.34±550.95 ^c	8.33±4.04 ^c	40.33±2.52 ^c
V	2.26±0.89	0.46±0.11	0.51±0.33	992.22±68.31	15.67±4.04	48.33±9.87

Results are mean ± standard deviation, Values in the same column bearing similar superscripts are significantly different at $P<0.05$. (n=3). Key: I: Negative Control Group, II: positive control and III, IV and V: Test groups, FLAE: Fresh Leaf Aqueous Extract, TP: Total Protein, ALB: Albumin, BIL: Bilirubin, ALP: Alkaline Phosphatase, ALT: Alanine Transaminase, AST: Aspartate Transaminase.

Table 3: Packed Cell Volume and Hemoglobin of rats after induction with CCl₄ and after treatment with FLAE of *F. indica*.

Group	After Induction		After Treatment	
	PCV	Haemoglobin	PCV	Haemoglobin
I	27.36±0.69 ^a	10.00±4.56 ^u	39.89±2.09 ^a	16.03±2.05 ^u
II	34.54±7.15 ^b	10.78±2.84 ^v	49.87±15.09 ^b	20.67±8.15 ^v
III	31.67±2.89 ^c	10.56±0.96 ^w	50.50±9.19 ^c	16.84±3.06 ^w
IV	43.00±4.58 ^d	14.00±2.00 ^x	41.00±7.55 ^d	13.67±2.52 ^x
V	36.50±4.95 ^e	12.89±1.17 ^y	45.33±4.16 ^c	15.11±1.39 ^y

Results are mean ± standard deviation, Values in the same row bearing similar superscripts are significantly different at $P<0.05$. (n=3). Key: I: Negative Control Group, II: positive control and III, IV and V: Test groups, FLAE: Fresh Leaf Aqueous Extract.

There were significant differences ($P<0.05$) observed in comparing group I with groups III, IV and V. Also, significant differences ($P<0.05$) were observed when the treated groups (III, IV and V) are compared with one another, not necessary in a dose dependent manner. As shown in Figure 1, the results of induction with 200mg/kg CCl₄ reveal significantly higher ($P<0.05$) levels of groups II (positive control), III, IV and V compared to group I (negative control).

Results in Figure 4 also reveal significantly higher ($P<0.05$) differences in the concentration of Alkaline Phosphatase (ALP) when group I (negative control) is compared with group III (400mg/kg FLF. *indica*), group IV (600mg/kg FLF. *indica*) and group V (800mg/kg FLF. *indica*). When the concentrations of ALP in groups III, IV and V were compared with one another, significant differences ($P<0.05$) were observed in a dose dependent pattern.

Figure 4: Alkaline phosphatase for CCl₄ induced rats treated with fresh leaf aqueous extract of *F. indica*.

FLF. *indica* = fresh leaf of *F. indica*. Letters a, b, c, d and e indicates significant difference ($P<0.05$) when group II was compared with groups I, III, IV and V, respectively for 400mg/kg, 600mg/kg and 800mg/kg *F. indica* treated groups.

DISCUSSION

Carbon tetrachloride (CCl₄) induced hepatic injury is shown in Table 1 and (Figure 1 and Figure 3). The rise in the level of ALT, AST, ALB, ALP, BIL and TP of groups II, III, IV and V when compared to group I indicates a CCl₄ induced liver damage. This is consistent with²², who reported the use of 1.5ml/kg body weight of CCl₄ orally administered to rats to induce liver damage. These liver function enzymes are found to be located in the cytosol of the liver cell and thus, are easily released into the serum after cellular liver damage²³. The mechanism of action involved in CCl₄ hepatic injury is understood to be linked to the liver phase II detoxification action. The liver, in the process of detoxification transforms Carbon tetrachloride (CCl₄) in the presence and action of cytochrome P₄₅₀ enzyme component to produce peroxy trichloromethyl and trichloromethyl free radicals¹³. These free radicals results in lipid peroxidation by

reacting covalently with biomolecules (proteins, nucleic acids, lipids etc) in the presence of oxygen. Thus the liver becomes damaged and obviously its cell membrane becomes degenerate, permeable and leaks out its cellular contents of AST, ALT, TP, ALP, BIL and ALB²⁴.

Table 4: Blood Glucose Concentration (mg/dl) of Rats after Induction with CCl₄ and after Treatment with FLAE of *F. indica*.

Group	After Induction	After Treatment
I	100.39±1.66 ^a	89.88±25.71 ^a
II	88.26±1.19 ^b	99.93±2.76 ^b
III	102.33±1.52 ^c	95.00±5.29 ^c
IV	93.33±4.16 ^d	80.00±12.29 ^d
V	82.67±15.37 ^e	81.00±11.14 ^e

Results are mean ± standard deviation, Values in the same row bearing similar superscripts are significantly different at $P < 0.05$. (n=3). Key: I: Negative Control Group, II: positive control and III, IV and V: Test groups, FLAE: Fresh Leaf Aqueous Extract.

After induction with CCl₄, the level of blood glucose was raised (Table 4), hemoglobin and PCV (Table 3) levels were decreased. This could suggest that CCl₄ induced anemia was possible owing from the destruction of red blood cells and shortage of circulating mineral iron and vitamins¹⁵. Hyperglycemia induced by CCl₄ could be due pancreatic injury caused by generation of free radical, cell membrane lipid peroxidation and subsequent destruction of pancreatic β - islet cells³.

Table 5: Body Weight of rats before induction with CCl₄, after induction with CCl₄ and after treatment with FLAE of *F. indica*.

Group	At acclimatization	At induction with CCl ₄	Treatment with FLAE of <i>F. indica</i>
I	184.49±11.55 ^a	218.21±7.70 ^a	206.45±16.05 ^a
II	179.10±13.13 ^b	221.15±14.42	207.55±6.16 ^b
III	222.48±13.93 ^c	187.76±28.90 ^c	235.10±34.22 ^c
IV	246.63±48.53 ^d	227.95±41.98 ^d	241.27±63.52 ^d
V	227.20±45.07 ^e	219.63±39.17 ^e	274.43±38.20 ^e

Results are mean ± standard deviation, Values in the same row bearing similar superscripts are significantly different at $P < 0.05$. (n=3). Key: I: Negative Control Group, II: positive control and III, IV and V: Test groups, FLAE: Fresh Leaf Aqueous Extract.

Administration of CCl₄ induced rats with FLAE of *F. indica* shows reduction in the concentrations of liver function enzymes (TP, AST, ALT, ALP, and BIL) in the serum and blood glucose (Table 2 and Table 4 and Figure 2 and Figure 4). Similarly, after treatment with FLAE of *F. indica* the levels of ALB, Hb and PCV increased as shown in Table 2 and Table 4. The reduced serum levels of the liver function enzymes indicate the recuperative, regenerative and healing effect of FLAE of *F. indica* on the hepatic cells. This is in support of a previous study³, who reported that treatment with *F. indica*'s ethanol extract stem bark with 500 mg/kg and 700 mg/kg in rats liver revealed regeneration of hepatocytes and absence of inflammation. It appears FLAE of *F. indica* exert its effects by antioxidant and free radical scavenging strength by furnishing the body with antioxidants

phytochemicals (tannins, flavonoids, carotenoids, anthocyanins), minerals (Fe, Mg, Mn, Na, K) and vitamins (A, C and E)^{4,22}.

Blood glucose was found to decrease with treatment with FLAE of *F. indica*. This could be associated with its antioxidant ability of increasing insulin production and regeneration of the β - islet cells of pancreas that was ones destroyed by CCl₄ induction^{22,7} buttresses that the presence of minerals in plants enhance effective function of the glycolytic pathway enzymes for the breakdown of glucose. That minerals enhances the phosphorylation conversion reaction of glucose to glucose 6-phosphate by the action of the enzyme hexokinase or glucokinase and phosphorylation of fructose 6-phosphate by the action of phosphofructokinase (PFK) to fructose 1, 6-bisphosphate⁷. The hematological indices of PCV and Hb after treating the CCl₄ anemia induced rats with FLAE of *F. indica* revealed increased levels of PCV and Hb. The anti-anemic and haem regenerating effects of FLAE of *F. indica* as depicted by the results of this study, could be associated to some extent on the antioxidants phytochemical and mineral elements it contains^{14,9}. Antioxidant phytochemicals such as saponins, flavonoids and alkaloids have been reported for their anti-anemic abilities, prevent thrombosis and aggregation of platelet and promote enhanced blood circulation^{25,26}. Thus, FLAE of *F. indica* was able to increase the levels of PCV and Hb because of these phytochemicals and mineral elements contents. This action could be made possible by its enhanced removal of the toxic effects caused by CCl₄ and creating flourishing iron utilization for the production of heme and subsequent release of new red blood cells¹⁴. FLAE of *F. indica* could have exert its effects in the improved production of Hb and PCV by enhancing the production of erythropoietin in the bone marrow stem cells and subsequent synthesis of new blood cells²⁷. In addition, there was an observed reduced body weight of anemic rats (CCl₄ induced groups) when compared to the groups treated with FLAE of *F. indica* (Table 5). This reduction in body weight of anemic rats and increase or weight gain in the treated group with FLAE of *F. indica* is consistent with the report of previous study¹⁴. The association of weight loss with anemia is not very clear. However, it appears to be related to defect in carbohydrates digestion in the small intestine of anemic rats due to insufficient amount of the enzyme, disaccharidases, thus leading to undigested carbohydrates²⁸.

CONCLUSION

The use of fresh leaf aqueous extract of *F. indica* in this study reveals that the plant possesses anti-anemia, hypoglycemic and hepato-healing potentials. This is obviously seen in the reduced levels of blood glucose, liver function assay, and in the raised levels of the hematological parameters, coupled with weight gain after treating CCl₄ induced groups with FLAE of *F. indica*.

AUTHOR'S CONTRIBUTION

All authors have worked equally for this work.

ACKNOWLEDGEMENTS

The authors extend their thanks and appreciation to the Caritas University, Nigeria to provide necessary facilities for this work.

CONFLICT OF INTEREST

Authors have declared that no conflict of interest is linked with this work.

REFERENCES

1. Eramma N, Devaraja G. Antibacterial potential and phytochemical analysis of *Flacourtia indica* (Burm.f.) Merr. root extract against human pathogens. *Indo American J Pharm Res* 2013; 3(5).
2. Ihcène B, Asma B, Amel B, Abdelfattah EF, Mahfoud M. Antioxidant effect of alpha lipoic acid on hepatotoxicity induced by aluminium chloride in rats. *Int J Pharm Sci Rev Res* 2014; 29(2): Pages: 19-25.
<https://doi.org/10.1155/2018/2817036>
3. Idokoa A, Emmanuel UG, Yakubu O, Ugwu-dike P, Patricia N. Hepatocurative and gluco-stabilizing potentials of ethanol extract of stem bark of *flacourtia indica* in aluminium chloride induced toxicity in albino wistar rats. *Curr Trends Biomedical Eng Biosci* 2018; 17(3): 1-7. DOI: 10.19080/CTBEB.2018.17.555965.
<https://doi.org/10.1155/2018/2817036>
4. Ammar A, Naoufal L, Azam B, Dennis G. Watson and David A. Lightfoot. Phytochemicals extraction, isolation, and identification of bioactive compounds from plant extracts. *Plants* 2017; 6(42): 1-23.
<https://doi.org/10.3390/plants6040042>
5. Vivekananthan DP, Penn MS, Sapp SK, Hsu A, Topol EJ. Use of antioxidant Vitamins for the prevention of Cardiovascular diseases. Meta-analysis of Randomized trials. *Lancet* 2003; 361: 2017-2023.
[https://doi.org/10.1016/S0140-6736\(03\)13637-9](https://doi.org/10.1016/S0140-6736(03)13637-9)
6. Abubakar SM, Umar SA, Alexander I, Abubakar N, Abdulazeez MA, Sule MS. Evaluation of hypoglycaemic, hypolipidaemic and non toxic effect of hydro-methanolic extracts of *Ziziphus mauritiana*, *Ziziphus spina christi* fruit and glibenclamide on alloxan induced diabetic rats. *J Drug Deliv Therap* 2018; 8(3): 82-92.
<https://doi.org/10.22270/jddt.v8i3.1711>
7. Idoko A, Ikpe VPO, Rita ON, Nelson NO, Effiong JU, Alhassan AJ, Muhammed IU, Abubakar N, Abubakar SM, Ugwu-dike PO. Hypoglycemic and lipid profile lowering effect of *chromolaena odorata* (linn) in albino wistar rats fed different concentrations of cholesterol enriched diet. *Univ J Pharm Res* 2018; 3(1): 37-42.
<https://doi.org/10.22270/ujpr.v3i1.17>
8. Idoko A, Ikpe VPO, Nelson NO, Effiong JU, Alhassan AJ, Muhammad IU, Abubakar, N, Abubakar SM. Effects of Lime Juice and Honey on Lipid Profile of Cholesterol Enriched Diet Fed Rat Model. *Annual Res Rev Biol* 2017; 20(3): 1-10. <https://doi.org/10.9734/ARRB/2017/37213>
9. Abhishek M, Thangadurai D, Shivanand B, Sangeetha J. Proximate analysis and mineral composition of potential minor fruits of Western Ghats of India. *Agronomy* 2017; Series A. Vol. LX: 340-346.
10. Alan F. An overview of the liver: A series of fact sheets written in the field of liver disease, Hepatitis C Support Project (HCSP publication); 2015 version 1; 1-5. Reviewed January 2018.
11. Tarasankar M, Surajit M, Nilanjan P, Durgesh RK, Subarna G. A review on hepatic diseases and development of herbal drugs for the treatment of liver complications. *World J Pharm Res* 2015; 4(6): 677-691.
12. Gupta AK, Ganguly P, Majumder UK, Ghosal S. Hepatoprotective and antioxidant effect and steroidal saponins of *Solanum xanthocarpum* and *Solanum nigrum* in paracetamol induce hepatotoxicity in rats. *Pharmacologyonline* 2009; 1: 757-768.
13. Weber LW, Boll M, Stampfl A. Hepatotoxicity and mechanism of action of haloalkanes: carbon tetrachloride as a toxicological model. *Crit Rev Toxicol* 2003; 33: 105-36. <https://doi.org/10.1080/713611034>
14. Luka CD, Abdulkarim M, Adoga GI, Tijjani H, Olatunde A. Anti-anaemic potential of aqueous extract of *spinacia oleracea* leaf in phenylhydrazine-treated rats. *New York Sci J* 2014; 7(6): 14-18.
15. Hoffman R, Edward B, Leslie S, Helen H, Jeffrey W, John Anastasi. Hematologic aspects of parasitic diseases. In: *Hematology: Basic Principles and Practice*. 6th ed. Philadelphia, Pa.: Saunders Elsevier; 2012. <http://www.elsevier.com>. Accessed January 17, 2019.
16. Khoo SB. Haematinics for Anaemia in the Elderly. *Malaysian Fam Phys* 2011; 6(1):1-8.
17. Zimmerman M. Ethical guidelines for investigations of experimental pain in conscious animals. *Pain* 1995; 16(2): 109-11. [https://doi.org/10.1016/0304-3959\(83\)90201-4](https://doi.org/10.1016/0304-3959(83)90201-4)
18. National Institute of Health. Guidelines for the care and use of laboratory animals, National Academic Press, 1937; 85-23.
19. Mauro P, Renze B, Wouter W. In: Tietz text book of clinical chemistry and molecular diagnostics. 4th edition. Carl AB, Edward R, David EB, editors. Elsevier; Enzymes. 2006; 604-616.
<https://doi.org/10.1007/s12291-012-0287-7>
20. Barbara JB, Imelda B. Basic haematologic techniques. In: Lewis SM, Bain BJ, Bates I, editors. *Dacie and Lewis Practical Haematology*. 11th ed. London: Churchill Livingstone; 2001; 19-46.
<https://doi.org/10.1016/B0-443-06660-4/X5001-6>
21. Graphpad Instat3 and graphpad prism Softwares. 2000. Available: www.graphpad.com. Retrieved on 24th November, 2018.
22. Anindita B, Gangarao B. *Flacourtia indica* (burm.f.) prevents ccl4 induced rat liver damage by augmenting antioxidant enzyme activity. *Eur J Pharm Med Res* 2016; 3(4): 263-270.
23. Bagban IM, Roy SP, Chaudhary A, Das SK, Gohil KJ, Bhandari KK. Hepatoprotective activity of the methanolic extract of *Fagonia indica* Burm in carbon tetra chloride induced hepatotoxicity in albino rats. *Asian Pacific J Trop Biomed* 2012; 1457-1460.
[https://doi.org/10.1016/S2221-1691\(12\)60437-7](https://doi.org/10.1016/S2221-1691(12)60437-7)
24. Recknagel RO, Glende EA, Dolak JA, Waller RL. Mechanisms of carbon tetrachloride toxicity. *Pharmacol Ther* 1989; 43: 139-154. [https://doi.org/10.1016/0163-7258\(89\)90050-8](https://doi.org/10.1016/0163-7258(89)90050-8)
25. Shi J, Arunasalam K, Yeung D, Kakuda Y, Mittal G, Jiang Y. Saponins from edible legumes: chemistry, processing, and health benefits. *J Med Food* 2004; 7: 67-78.
<https://doi.org/10.1089/109662004322984734>
26. Falcone A, Musto P, Rasario P, Rosella M. Compounds and Methods for treatment of chemotherapy-induced Anemia. *Europ J Haemat* 1997; 58:34-319.
27. Ganong WF. A review of medical physiology appleton and Lange Pubiisher. 1997:96.
28. Vieira MR, Galzvaio LC, Fernandes MIM. Relation of the disaccharidases in the small intestine of the rat to the degree of experimentally induced iron-deficiency anaemia. *Braz J Med Biol Res* 2000; 33: 539-544.