Available online on 15.1.2017 at <http://ujpr.org>
Universal Journal of Pharmaceutical Research
 An International Peer Reviewed Journal

Open access to Pharmaceutical research
 This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial Share Alike 4.0 License which permits unrestricted non commercial use, provided the original work is properly cited

Volume 1, Issue 2, 2016

Open Access

REVIEW ARTICLE

MURRAYA KOENIGI-A BOON IN DIFFERENT PATHOLOGICAL CONDITIONS

Igwe J Chibueze^{1b}, Emenike IV^{1b}

Department of Pharmaceutical Microbiology and biotechnology Gombe state University, Nigeria.

ABSTRACT

Since very long period of time medicinal plants or their bioactive compounds have been utilized by majority of world population particularly in developing countries for primary and traditional healthcare system. At present scenario, people are more interested to use herbal drugs because they are considered as safe and inexpensive having no adverse effects. Different parts of the plants like roots, leaves, stem, bark, fruits and seeds have been used in treatment of different diseases and strengthening the immune system. *Murraya koenigii*, is a herb from mainly Asian origin, it has therapeutic applications such as in bronchial disorders, piles, vomiting, skin diseases, night blindness, dysentery, diarrhoea, bites of poisonous animals, bruises and eruption etc. The present review is an attempt for description of *M. koenigii*, its phytochemical constituents and various pharmacological activities.

Keywords: *Murraya koenigii*, phytochemistry, pharmacological activities.

Article Info: Received 6 November 2016; Revised 3 December; Accepted 30 December, Available online 15 January 2017

Cite this article-

Chibueze IJ, Emenike IV. *Murraya koenigii*- a boon in different pathological complications. Universal Journal of Pharmaceutical Research 2016; 1(2): 36-40.

DOI: <http://doi.org/10.22270/ujpr.v1i2.RW5>

Address for Correspondence:

Igwe J Chibueze, Department of Pharmaceutical Microbiology and Biotechnology, Gombe state University, Nigeria, E-mail: Igwejames42@yahoo.com

INTRODUCTION

Murraya koenigii (family Rutaceae) is a herb having characteristic aroma and different potential medicinal values. It is deciduous shrub or tree up to 6 m in height and 15-40 cm in diameter with short trunk, thin smooth grey or brown bark and dense shady crown¹. The most important chemical constituents responsible for its intense characteristic aroma are P-gurjunene, P-caryophyllene, P-elemene and O-phellandrene. The plant is rich source of carbazole alkaloids. Phytocompounds like koenimbine, koenine, mahanimbine, murrayazolidine, murrayazoline, murrayacine, girinimbine, mukoeic acid, etc. have also been isolated and characterized². The stem of *M. koenigii* is an aromatic and more or less deciduous shrub or small tree upto 6 meters in height and 15 to 40 cm in diameter. The main stem is dark green to brownish. The bark of the stem can be peeled off longitudinally which exposes the white wood underneath. Flowers are small, white fragrant ebracteate, calyx deeply five cleft, pubescent³. Petals five, free, whitish, glabrous and with dotted glands. Fruits occur in close clusters, small ovoid or sub-globose, glandular, thin pericarp enclosing one or two seeds having spinach green color⁴. Fresh leaves, dried leaf powder, and essential oil are widely used for

flavouring soups, curries, fish and meat dishes, eggs dishes, traditional curry powder blends, seasoning and ready to use other food preparations. Bark and roots are used as stimulant and externally to cure eruptions and bites of poisonous animals⁵. It is traditionally used as a whole or in parts as anti-emetics, anti-diarrheal, febrifuge, blood purifier, antifungal, depressant, anti-inflammatory, body aches, for kidney pain and vomiting. Green leaves are eaten raw for cure of dysentery, diarrhoea and for checking vomiting⁶. Leaves and roots are also used traditionally as bitter, anthelmintic, analgesic, curing piles, inflammation, itching and are useful in leucoderma and blood disorders⁷.

Table 1: Botany of plant.

Kingdom	Plantae
Sub-kingdom	Tracheobionta
Super division	Spermatophyta
Division	Magnoliophyta
Class	Magnoliopsida
Subclass	Rosidae
Order	Sapindales
Family	Rutaceae
Species	<i>Murraya koenigii</i>
	L. Spreng

Flowers are white, ebracteate, scented and small in size. Calyx deeply five cleft, pubescent. Petals five, free, whitish, glabrous and with dotted glands. It bears fruits in close clusters/ bunches, small, ovoid or sub-glucose, glandular, thin pericarp enclosing one or two seeds having spinach green color.

(a). Whole plant

(b). Leaves

Figure 1: *Murraya koenigii*.

Origin and Distribution

Murraya koenigii is basically found in tropical Asia like the foothills of Himalayas of India, Sri Lanka, Myanmar, Indonesia, Southern China and Hainan. It reproduces the means of seeds which germinate freely under partial shade. It is also available in other part of Asian region like in moist forests of 500-1600m height in Guangdong, S Hainan, S Yunnan (Xishuang banna), Bhutan, Laos, Nepal, Pakistan, Sri Lanka, Thailand, Vietnam. Together with South Indian immigrants, curry leaves reached Malaysia, South Africa and Réunion Island. They are hardly found outside the Indian sphere of influence⁸.

PHYTOCHEMISTRY

Murraya koenigii is a rich source of different organic compounds with diverse chemical composition.

Leaves

Fresh young leaves contains yellow colored volatile oil rich in vitamin A, calcium, girinimbine, isomahanimbine, koenine, koenigine, koenidine and koenimbine. Leaves are aromatic and contain proteins, carbohydrates, fiber, minerals, carotene, nicotinic acid and vitamin C. Mature leaves contains 63.2% moisture, 1.15% total nitrogen, 6.15% fat, 18.92% total sugars, 14.6% starch, 6.8% crude fiber, ash 13.06%, acid insoluble ash 1.35%, alcohol soluble extractive 1.82%. The leaves contain high amount of oxalic acid, leaves also contains crystalline glycosides, carbazole alkaloids, koenigin, resin, fresh leaves contain yellow color 2.5% volatile oil⁹.

Bark

Bark mainly contains the carbazole alkaloids as murrayacine, murrayazolidine, murrayazoline, mahanimbine, girinimbine, koenioline, xynthyletin. The alcoholic extract of stem bark shows the presence of koenigine-quinone A and koenigine quinone B⁹.

Fruits

The pulp of fruits contains 64.9% moisture, 13.35% of vitamin C, 9.76% total sugar, 0.17% non-reducing sugar, trace amount of minerals (1.97% phosphorus, 0.082% potassium, 0.811% calcium, 0.166% magnesium and 0.007% iron)¹⁰.

Roots

Roots contains murrayanol, mukoenine- A, -B and C and murrastifoline -F. bis - 2- hydroxy- 3- methyl carbazole, bismahanine, bi koeniquinone- A and bismurrayaquinone A, Koenoline (1- methoxy-3- hydroxymethyl carbazole). Roots were also found to contain girinimbine¹¹.

Seeds

Mahanimbine, girinimbine, koenimbine, isomahanine and mahanine were isolated from seeds of *M. koenigii*¹².

pharmacological activity profile of *M. koenigii*

1. Antioxidant and free radical-scavenging activity:

Antioxidant activity has been reported by a number of workers. In a study antioxidative properties of the extract of *M. koenigii* leaves were done using different solvents. They were evaluated on the basis of oil stability index together with their radical scavenging ability against 1-1-diphenyl-2-picrylhydrazyl. The methylene chloride extract and the ethyl acetate soluble fraction of the 70 % acetone extract was prolonged^{13,14}.

2. Cytotoxic Activity- In a study the alkaloid koenoline isolated from the root bark of *M. koenigii* is found to exhibit cytotoxic activity against KB cell culture system¹⁵. Carbazole alkaloids isolated from the stems are found to have significant effects in the growth of the human leukaemia cell line HL-60. Mahanine, pyrafoline-D and murrayoline-I (Carbazole alkaloids) showed significant cytotoxicity against HL-60 cells and cause a significant loss in mitochondrial membrane potential. The results obtained suggested it, s cytotoxic activity potential¹⁶.

3. Hypoglycemic activity

Leaves feeding produce hypoglycemia by increasing the hepatic glycogenesis as evident by increased activity of glycogen synthetase¹⁷. In a study a decrease in glycogenolysis and gluconeogenesis is reported and was evident from decreased activity of glycogen phosphorylase and gluconeogenic enzymes. A significant reduction in fasting blood sugar and postprandial blood sugar was observed by feeding (12 gm) leaves powder to non insulin dependent diabetes mellitus patients. The results obtained suggested it, s hypoglycemic activity potential¹⁸.

4. Antimicrobial and anti-fungal activity

In a study Murrayanine, girinimbine and mahanimbine isolated from stem bark showed anti fungal activity against human pathogenic fungi. 1- formyl-3 methoxy-6- methyl carbazole and 6,7-dimethoxy-1- hydroxy-3- methyl carbazole were reported to possess antibacterial and anti fungal property. Extract containing murrayanol and or isomahanine is used as microbicide in variety of industries due to high safety, strong activity, little odor and without coloring effect¹⁹.

5. Anti-inflammatory activity

In a study stem bark of alcoholic extract in a dose of 1 gm/kg body weight showed protective effect against carrageenan-induced inflammation. Crude root extract also showed anti-inflammatory activity in rat model. The mast cell stabilization and antihistaminic effects of EEMK were suggested to be the probable mechanisms for its anti- inflammatory action and thus attains its

therapeutic value. Study concludes anti-inflammatory potential of *M. koenigii*²⁰.

6. Immunomodulatory activity

In a study methanolic extract of *M. koenigii* showed significant increase in phagocytic index by rapid removal of carbon particles from blood stream. The extract also increased the antibody titre against ovalbumin and protection towards cyclophosphamide-induced myelosuppression in albino mice. Oral administration of the aqueous extract of leaves at doses of 250 and 500 mg/kg significantly enhanced the delayed-type hypersensitivity reaction induced by ovalbumin. The extract also potentiated the production of circulating antibody titre significantly in response to ovalbumin²¹.

7. Alzheimer disease therapy

In a study administration ethanolic extract of *M. koenigii* Leaves for 15 days produces significant dose-dependent improvement of memory. The results also indicated to reduce the brain cholinesterase activity and total cholesterol level. Diet rich in *M. koenigii* leaves produced significant dose dependent improvement in the memory scores of young and aged mice and significantly reduced the amnesia induced by scopolamine (0.4 mg/kg, intraperitoneally) and diazepam (1 mg/kg, intraperitoneally)²².

8. Anti-obesity and anti-hyperlipidemic activities

In a study the dichloromethane and ethyl acetate extracts of *Murraya koenigii* leaves significantly reduced the body weight gain, plasma total cholesterol and triglyceride levels significantly. The observed anti-obesity and antihyperlipidemic activities of these extract are correlated with the carbazole alkaloids, Mahanimbine. When it was given orally (30 mg/kg/day) significantly lowered the body weight gain. These findings demonstrate the excellent pharmacological potential of mahanimbine to prevent obesity²³.

9. Anti-amnesic and wound-healing activity-

In a study aqueous extract of *M. koenigii* accelerates the wound-healing process by decreasing the surface area of the wound. Aqueous extract of leaves showed marked reduction in wound area in comparison with the control group from 4th day onwards in albino rats by excision wound model²⁴.

10. Kidney protective activity

In a study aqueous extract of leaves produced a significant dose- dependent decrease in serum urea and creatinine levels ($P < 0.001$), and a marked increase in the levels of plasma antioxidant capacity ($P < 0.01$) in diabetic rats, compared with the control (non-diabetic) subjects. Histological studies of the kidneys of these animals showed comparable tissue regeneration by the aqueous extract²⁵.

11. Antipyretic activity

In a study ethanolic extract of leaves of *M. koenigii* was investigated for antipyretic activity in rats using yeast-induced pyrexia model. Ethanolic extract at a single dose of 300 mg/kg produced significant antipyretic activity ($P < 0.01$) in albino rats as compared with the standard drug paracetamol²⁶.

12. Anti-ulcer activity

Antiulcer activity of aqueous and solvent ether extracts of *Murraya koenigii* was studied in reserpine induced gastric ulcer model in albino rats. Aqueous and solvent ether extracts of *Murraya koenigii* effective in gastric ulceration and suggested as protective as ranitidine.

The extract dose of *Murraya koenigii* 200-400 mg/kg produced significant inhibition of gastric secretion. The results obtained suggested that the extract possesses significant antiulcer activity²⁷.

13. Anti-trichomonal activity

In an earlier study carbazole alkaloids and their derivatives from *M. koenigii* leaves showed antitrichomonal activity against *Trichomonas gallinae*. Girinimbine and girinimbilol with IC₅₀ values of 1.08 and 1.20 mg/mL were the most active. Acetylation of girinimbilol and mahanimbilol improved their activities to 0.60 and 1.08 mg/MI²⁸.

14. Anthelmintic activity

Ethanolic and aqueous extracts from *M. koenigii* leaves were investigated for their anthelmintic activity against *Pheretima posthuma*. Both the extracts exhibited significant anthelmintic activity at concentration of 100 mg/mL. The alcoholic extract produced more significant anthelmintic activity than petroleum ether extract²⁹.

15. Cosmetic use

Hyaluronidase inhibitors are extracted from *M. koenigii* and are formulated in a cream base. *M. koenigii* extract is included in a skin-lightening cosmetic for its moisturizing, antioxidant and hyaluronidase inhibitory activity. Herbal composition containing *M. koenigii* stem extract as one of the ingredient showed skin lightening and rough skin improving effect. *M. koenigii* was studied for sun protection^{30,31}.

16. Anti-diarrhoeal activity

In a study bioactive alkaloids like, kurryam and koenimbine obtained from fractionated n-hexane extract of the seeds of *M. koenigii* were showed inhibitory action in reference to castor oil-induced diarrhoea and prostaglandin E₂-induced enter pooling in strain of Wistar rats in charcoal meal test in Wister rats, these compounds were found to exhibit significant reduction in gastrointestinal motility and play mandate role in studying the modulatory role in disease progression³².

17. Anti-cancer Activity

Intraperitoneal inoculation of Dalton's Ascitic Lymphoma cells in the mice produced an enormous increase in the cancer cell count which indicated that there is progression of cancer in the animals³³. The decrease in the cancer cell number observed in the ether extract of *Murraya koenigii* the treated mice indicates that the test drug is having significant inhibitory effect on the tumour cell proliferation. The increase in tumour weight may be due to accumulation of peritoneal fluid as an abnormal enlargement of peritoneal cavity was observed in tumour-induced mice. Treatment with extract of *Murraya koenigii* reduced the tumour weight and hence increased the life span³⁴.

Table 2: Pharmacological activity shown by extract of different parts of *Murraya koenigii*.

S. N.	Plant Part	Pharmacological Activity	Extract
1.	Leaf	Anti-inflammatory Anti-amnesic, memory enhancer, anti-tumor Hypocholesterolemic, wound healing activity, anti-helminthic Anti-fungal Analgesic and antinociceptive, radioprotective and chemoprotective, anti-oxidant, phagocytic activity, anti-lipid peroxidative Anti-ulcer, cardiovascular	Ethanol, petroleum ether, chloroform, methanol Petroleum ether Ethanol Petroleum ether, alcohol and acetone Methanol Aqueous
2.	Bark, leaf	Anti-bacterial	Petroleum ether, alcohol
3.	Stem bark	Anti-cancer	Petroleum ether
4.	Seeds	Antidiarrhoeal	n-hexane
5.	Roots, stem	Cytotoxicity	Aqueous
6.	Leaf, fruit	Anti-diabetic	Aqueous, methanol

CONCLUSION

At present scenario, people are moving towards the use of herbal medicine for any kind of treatment as it seems to be economical and more beneficial without any adverse effects. According to WHO (World Health Organization), 80% of the population of developing countries still rely on plant-based medicines. The presence of various beneficial constituents in plants has always motivated scientists to carry out research for investigations for finding new therapeutic agents for treatment of different diseases. Based on tremendous pharmacological activities and plenty of literature available, *M. koenigii* may be utilized to alleviate the symptoms of variety of diseases. Almost each and every part of the plant has numerous therapeutic values. Various parts of it have numerous medical applications; modern drugs can be developed after extensive investigation of its bioactivity, mechanism of action, pharmaco-therapeutics, toxicity and after proper standardization and clinical trials. Wide spread availability of *M. koenigii* makes it suitable candidate for further pre-clinical and clinical research. From the available literature it can be stated that *Murraya koenigii* is a versatile medicinal plant having rich source of biologically active compounds. Thus, it can be consider being a most suitable candidate for new drug discovery evaluated by means of scientific experimental animal models and clinical trials.

AUTHOR'S CONTRIBUTION

All authors have worked equally for this work.

ACKNOWLEDGEMENTS

The authors extend their thanks and appreciation to the Gombe state University, Nigeria to provide necessary facilities for this work.

CONFLICT OF INTEREST

No conflict of interest associated with this work.

REFERENCES

- Ito C. Studies on medicinal resources of rutaceous plants and development to pharmaceutical chemistry. *Natural Med* 2000; 54: 117-122.
- Rahman MM, Gray AI. A benzoisofuran one derivative and carbazole alkaloids from *Murraya koenigii* and their antimicrobial activity. *Phytochemistry* 2005; 66:1601-1606. <https://doi.org/10.1016/j.phytochem.2005.05.001>
- Iyer UM, Mani UV. Studies on the effect of curry leaves supplementation (*Murraya koenigii*) on lipid profile, glycated proteins and amino acids in non-insulin-dependent diabetic patients. *Plant foods hum nutr.* 1990; 40(4): 275-282. <https://doi.org/10.1007/bf02193851>
- Adebajo AC, Olayiwola G, Verspohl EJ, Iwalewa EO, Omisore NOA, Bergenthal D, et al. Evaluation of the ethnomedical claims of *Murraya koenigii*. *Pharm Biol.* 2004; 42(8): 610-620. <https://doi.org/10.1080/13880200490902518>
- Palaniswamy UR, Caporuscio C, Stuart J. A chemical analysis of antioxidant vitamins in fresh curry leaves by reverse phase high performance liquid chromatography with UV detection. *Acta Horti* 2003; 620: 475-478. <https://doi.org/10.1021/jf0481711>
- Tachibana Y, Kikuzaki H, Lajis NH, Nakatani N. Anti oxidative activity of carbazoles from *Murraya koenigii* leaves. *J Agric Food Chem* 2001; 49: 5589-5594. <https://doi.org/10.1021/jf010621r>
- Tachibana Y, Kikuzaki H, Lajis NH, Nakatani N. Anti oxidative activity of carbazoles from *Murraya koenigii* leaves. *J Agric Food Chem* 2001; 49: 5589-5594. <https://doi.org/10.1021/jf010621r>
- Tachibana Y, Kikuzaki H, Lajis NH, Nakatani N. Comparison of Anti oxidative properties of Carbazole Alkaloids from *Murraya koenigii* Leaves. *J Agri Food Chem* 2003; 51:6461-6467. <https://doi.org/10.1021/jf034700+>
- Fiebig Manfred, Pezzuto John M., Soejarto Djaja D., Plant Anticancer Agents. Part 40, Koenoline A further cytotoxic carbazole alkaloid from *Murraya koenigii*. *Phytochemistry* 1985; 24(12):3041-3043. [https://doi.org/10.1016/0031-9422\(85\)80052-2](https://doi.org/10.1016/0031-9422(85)80052-2)
- Nutan MTH, Hasnat A, Rashid MA, Antibacterial and cytotoxic activities of *Murraya koenigii*. *Fitoterapia* 1998; 69(2):173-175. <https://doi.org/10.1186%2F1472-6882-14-87>

11. William F, Lakshminarayan S, Chegu H. Effect Of Some Indian vegetables on the glucose and insulin response in diabetic subjects. *Int J Food Sci Nutr* 1993; 44(3):191-196. <https://doi.org/10.3109/09637489309017439>
12. Tembhurne SV, Sakarkar DM. Hypoglycemic effects of fruit juice of *Murraya koenigii* (L.) in alloxan induced diabetic mice. *Int J Pharm Tech Res* 2009;1(4):1589-1593. PMID: 21589767
13. Khuntia TK, Panda DS. Evaluation of antibacterial, antifungal and anthelmintic activity of *Murraya koenigii* Spreng. *Pharma Sci Monit* 2011; 2(2):105-110.
14. Darvekar VM, Patil VR, Choudhari AB. Anti-inflammatory activity of *Murraya koenigii* Spreng on experimental animals. *J Nat Prod Plant Resour* 2011; 1(1):65-69.
15. Rakesh K Sindhu, Sandeep Arora. Phytochemical and pharmacognostical studies on *Murraya koenigii* (L.) Spreng roots. *Drug Inven Tod* 2012; 4(1):325-333.
16. Adesina SK, Olatunji OA, Brgenthal D, Reisch J, New biogenetically significant constituents of *Clausena anisata* and *Murraya koenigii*. *Pharmazie* 1988; 43(3): 221-222.
17. Kureel SP, Kapil RS, Popli SP. Terpenoid Alkaloids Form *Murraya koenigii* Spreng- II: Constitution of Cyclomahanimbine, Bicyclomahanimbine and Mahanimbidine. *Tetrahedron Lett.* 1969; 44:3857-3862. [https://doi.org/10.1016/S0040-4039\(01\)88531-2](https://doi.org/10.1016/S0040-4039(01)88531-2)
18. Reisch J, Adebazo AC, Kumar V, Aladesanmi AJ. Two carbazole alkaloids from *Murraya koenigii*. *Phytochemistry* 1994; 36(4):1073-1076. [https://doi.org/10.1016/S0031-9422\(97\)00345-2](https://doi.org/10.1016/S0031-9422(97)00345-2)
19. Adebajo Adeleke C, Olugbade Tiwalade A, Elujoba Anthony A, Aladesanmi Adetunji J, J Reisch Johannes, 2', 3' Epoxyindicolactone from *Murraya koenigii*. *Niger J Nat Prod Med* 1997; 1(1): 21-24. <https://doi.org/10.4314/njnp.v1i1.11794>
20. Li Q, Zhu Liangfeng, et al. Monoterpene and sesquiterpene rich oils from the leaves of *Murraya* species: chemotaxonomical significance. *Biochem Syst Ecol.* 1988; 16(5):491-494. <https://doi.org/10.20959/wjpps20172-8740>
21. Mac Leod, Alexander J, Pieris Nirmala M. Analysis of essential oil of *Murraya koenigii* and *Pandanus latifolius*. *Phytochem* 1982; 21(7):1653-1657. [https://doi.org/10.1016/S0031-9422\(82\)85034-6](https://doi.org/10.1016/S0031-9422(82)85034-6)
22. Ningappaa MB, Dineshaa R, Srinivasa L. Antioxidant and free radical scavenging activities of polyphenol-enriched curry leaf (*Murraya koenigii* L.) extracts. *Food Chem* 2008; 106(2):720-728. <https://doi.org/10.1016/j.foodchem.2007.06.057>
23. Birari R, Javia V, Bhutani KK, Antiobesity and lipid lowering effects of *Murraya koenigii* (L.) Spreng leaves extracts and mahanimbine on high fat diet induced obese rats. *Fitoterapia* 2010; 81(8):1129-33. <https://doi.org/10.1016/j.fitote.2010.07.013>
24. Mathur A, Verma SK, Singh SK, Prasad GBKS, Dua VK. Investigation of the antimicrobial, antioxidant and anti-inflammatory activity of compound isolated from *Murraya koenigii*. *Int J Appl Biol Pharm Technol* 2011; 2(1):470-477.
25. Parmar S, Gangwal A, Sheth N, Mast cell membrane stabilization and anti-histaminic actions possible mechanism of action of anti-inflammatory action of *Murraya koenigii*. *J Curr Pharm Res* 2010, 2 (1), 21-25.
26. Ningappaa MB, Dhananjayaa BL, Dineshaa R, Harsha R, Srinivas L. Potent antibacterial property of APC protein from curry leaves (*Murraya koenigii* L.). *Food Chem* 2010; 118, 747-750. <https://doi.org/10.1016/j.foodchem.2009.05.059>
27. Patidar DK. Anti-ulcer activity of aqueous extract of *Murraya koenigii* in albino rats. *Int J Pharma Bio Sci* 2011; 2(1):524-529. <https://doi.org/10.4103%2F0257-7941.147434>
28. Adebajo AC, Olayiwola G, Verspohl EJ, Iwalewa EO, Omisore NOA. Bergenthal Evaluation of the ethnomedical claims of *Murraya koenigii*. *Pharm Biol* 2004; 42: 610-620. <https://doi.org/10.1080/13880200490902518>
29. Khuntia TK, Panda DS. Evaluation of antibacterial, antifungal and anthelmintic activity of *Murraya koenigii* Spreng. *Pharma Sci Monit* 2011; 2(2):105-110. [https://doi.org/10.1016/S2222-1808\(12\)60175-3](https://doi.org/10.1016/S2222-1808(12)60175-3)
30. Inahata Keishiro, Shin Kunio, Microbicides useful in pharmaceutical cosmetic and food preparation Japan. Kokai Tokkyo koho JP. 1996; 225.
31. Nanba Tsuneo, Hatsutori Yukio, Shimomura Kenji, Nakamura Masami. Extraction of hyaluronidase inhibitors from *Azadirachta indica* or other plants for manufacturing cosmetics or for therapeutic use. Jpn. Kokai Tokkyo Koho JP 1995; 95. <https://doi.org/10.1016/j.indcrop.2017.11.019>
32. Mandal S, Nayak A, Kar M, Banerjee SK, Das A, Upadhyay SN et al. Antidiarrhoeal activity of carbazole alkaloids from *Murraya koenigii* Spreng (Rutaceae) seeds. *Fitoterapia* 2010; 8(1):72-74. <https://doi.org/10.1016/j.fitote.2009.08.016>
33. Ghasemzadeh A, Jaafar HZE. Antioxidant potential and anticancer activity of Malaysian young ginger (*Zingiber officinale* Roscoe) varieties grown under different CO2 concentration. *J Med Pla Res* 2011; 5 (14), 3247-3255.
34. Itharat PJ, Houghton E, Eno-Amooquaye PJ, Burke JH. In vitro cytotoxic activity of Thai medicinal plants used traditionally to treat cancer. *J Ethnopharmacol* 2004; 90 (1), 33-38. <https://doi.org/10.1016/j.jep.2003.09.014>