Available online on 15.11.2019 at <http://ujpr.org>
Universal Journal of Pharmaceutical Research
An International Peer Reviewed Journal

Open access to Pharmaceutical research
 This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial Share Alike 4.0 License which permits unrestricted non commercial use, provided the original work is properly cited

Volume 4, Issue 5, 2019

Open Access

REVIEW ARTICLE

AN OVERVIEW ON CURRENT REGULATION AND EVALUATION OF BIOCIDAL PRODUCTS

Ash ŞAHİNER¹, Evren ALGIN YAPAR^{2*}

¹Department of Biology, Faculty of Science, Ege University, 35100 Bornova, Izmir, Turkey.

²Department of Analysis and Control Laboratories, Turkish Medicines and Medical Devices Agency, 06100 Sıhhiye, Ankara, Turkey.

ABSTRACT

A biocidal product is a substance or mixture prepared to limit, destroy, neutralize or control the effects of a harmful microorganism, plants and animals. The active substance in a biocidal product can be a natural oil or extract, a chemical substance or a microorganism, virus or fungus. Biocides consist of four main groups: disinfectants, preservatives (wood, paint, etc.), pest control and other type of biocidal products. A biocidal substance can also be added to a product to make the product itself into a biocidal product. These products range from disinfectants, hand sanitizers, preservatives, insect repellents, to rodenticides and insecticides and are used to protect humans, animals, materials and articles by controlling the intended target organism by a chemical or biological action. To make sure the use of biocidal products do not have unacceptable risks for people, animals and the environment, they are regulated to control their marketing, sale and use. In the current study biocidal products have been overviewed in the scope of current European Union regulations, product types and conformity tests.

Keywords: Biocidal products, conformity tests for biocidal products, disinfectants.

Article Info: Received 1 September 2019; Revised 9 October; Accepted 7 November, Available online 15 November 2019

Cite this article-

ŞAHİNER A, ALGIN YAPAR E. An overview on current regulation and evaluation of biocidal products. Universal Journal of Pharmaceutical Research 2019; 4(5): 34-38.

DOI: <https://doi.org/10.22270/ujpr.v4i5.315>

Address for Correspondence:

Assos. Prof. Evren ALGIN YAPAR, Department of Analysis and Control Laboratories, Turkish Medicines and Medical Devices Agency, 06100 Sıhhiye, Ankara, Turkey. Tel: +90 312 565 53 70, E-mail: evrenalginy@yahoo.com

INTRODUCTION

According to the Biocidal Products Regulation [BPR, Reg. (EU) 528/2012] of European Commission (EC), a biocidal product is defined as a product or substance intended to eliminate, control or prevent the effects of harmful organisms for human and animal health and to control organisms harmful to natural or manufactured materials¹. The use of chemical biocides is a fundamental protection in the prevention and control of microbial growth in medical, veterinary, domestic and industrial environments². Biocides are used a great extent in the healthcare environment for the disinfection of equipments, surfaces, water, and for antiseptics of skin and wound. Besides biocides are used for the preservation of pharmaceuticals and sterilization of medical devices³. In the 20th century, the development of cationic biocides such as quaternary ammonium compounds, biguanides, aldehydes, peroxides and phenolics had been an enormous increase in the number of active compounds used for disinfection, sterilization and preservation⁴. Biocides are the main armory in the disinfection programme of food industry to control pathogenic and

spoilage micro-organisms⁵. Biocidal products also constitute the antimicrobial component of nanomaterials used in food packaging in recent years⁶. Preservatives are an important part of biocides and used to protect industrial products, cosmetics, metal and wood materials, textile products from microbial spoilage. Isothiazolinone biocides are broadly used in various industrial applications for the control of microbial growth. Between the biocides, 5-chloro-2-methyl-4-isothiazolin-3-one (CMIT), 2-methyl-4-isothiazolin-3-one (MIT), 1, 2-Benzisothiazolin-3-one (BIT), 4,5-dichloro-2-methyl-4-isothiazolin-3-one (DCO IT) are the most commonly used as preservative active substance⁷. Effective and proper use of insecticide and rodenticide group biocides is also important for the protection of human and environmental health. Permethrin, tetramethrin, cypermethrin are commonly used insecticide active substance for product type⁸. Misuse of biocidal products brings about some problems such as resistance development and toxicity. The spreading usage of products containing low concentrations of commonly used biocides have raised some concerns about the possible development of

microbial resistance. Laboratory studies have demonstrated that bacteria can become resistant to a biocide, and that resistant bacteria can develop cross-resistance to other biocides and antibiotics⁹. High concentrations of biocides generally have toxic effects not only for humans but also for the environment. Therefore, a legal guideline for environmental risk evaluation for biocidal products is being prepared¹⁰. The toxicity of some biocides associated with dermatitis, occupational asthma and irritation has been reported^{11,12,13}.

The use of commercially available biocidal products is regulated according to the EU Biocidal Products Regulation. In this study biocidal products have been overviewed in the scope of current EU regulations, product types and conformity tests.

RULES FOR BIOCIDAL PRODUCTS

The aim of the Biocidal Products Regulation (Reg. (EU) 528/2012) is to determine the rules for the production and the use of biocidal products, as well as to provide a protection from health and environmental risks. The main rule is that biocidal products must be authorized before they are placed on the market. This regulation applies to biocidal products containing or forming one or more active substances.

A biocidal product is a substance or mixture designed to limit, destroy, neutralize or control the effects of a harmful organism. All biocidal products contain one or more active substances. The active ingredients give the product the desired biocidal properties. Active substances used in biocidal products must be authorised within the EU before they can be put on the market. Before the approval of an active substance, its effect on the environment and human health is assessed. To ensure legal certainty, Active substances from the Union list approved for use in biocidal products has been established. In order to market an active substance that has not been previously authorized or under examination, It must be applied for authorization to The European Chemical Agency (ECHA)¹⁴.

The following conditions must be met in order to license a biocidal product:

- The active substances must be approved for the type of product concerned and the conditions specified for these active substances must be met.
- It must be sufficiently effective.
- The biocidal product must not produce unacceptable resistance or cross-resistance on target organisms. There should be no unacceptable effects such as unnecessary pain and suffering for vertebrates.
- The biocidal product or its residues must not have an unacceptable effect on food and feed, directly or indirectly.
- The biocidal product or its residues must not have an unacceptable effect on the environment, groundwater, surface water, drinking water, soil and air.
- It should not have undesirable effects on non-target organisms.

- It should not have undesirable effects on the ecosystem and biodiversity.
- The chemical identity, quantities and technical equivalence of the active and inactive substances in the biocidal product should be determined. Toxicological and ecotoxicological information should be provided where appropriate.

PRODUCT TYPES

In order to make a suitable assessment of exposure and risk to health and the environment as a result of a biocidal product's use, the EU has determined the different product types of biocidal products. According to the Biocidal Products Regulation (EU) 528/2012, biocides are classified in four main categories and twenty-two specific product types¹.

Disinfectants

PT 1- Human hygiene: Products are used for human hygiene. The primary purpose of these products is to disinfect human skin and scalp.

PT 2- Disinfectants and algacides: The aim of these products is to disinfect surfaces and equipment that are not contact with food or feeding stuff. Areas of use include air conditioning systems, aquariums, swimming pools, and private, public and industrial areas. It is also used for disinfection of air, water other than human or animal consumption, waste water, hospital waste and soil. They are incorporated into textiles, textures, paints and other articles to produce treated products.

PT 3- Veterinary hygiene: Antiseptics and disinfectants that are used for veterinary hygiene. Used for disinfecting animal shelters and transport equipment and surfaces.

PT 4- Food and feed area: It is used disinfecting equipment, surfaces or piping related to the production, transportation, storage of food or feed.

PT 5- Drinking water: Products in this group are used for disinfection of drinking water for human and animal consumption.

Preservatives

PT 6- In-can preservatives: It is used for the protection of cosmetic products, medical products or medical devices and manufactured products with microbial deterioration and control to ensure shelf life.

PT 7- Film preservatives: These products are used for the protection of films and coatings by controlling microbial deterioration or algae growth for the purpose of protecting the objects such as paint, plastic, wall adhesives, paper.

PT 8- Wood preservatives: These products are used for the protection of wood and wood products by controlling organisms that damage wood, including insects.

PT 9- Fibre, leather, rubber and polymerised materials preservatives: These preservatives are used for the preservation of polymers and fibres, such as textile, rubber, leather or paper by the control of microbiological degradation.

PT 10- Construction material preservatives: The purpose of these products are to preserve masonry, construction materials and composite materials except

for wood by the control of bacterial, fungal and algal attack.

PT 11- Preservatives for liquid-cooling and processing systems: Products in this group are used to protect liquids used in cooling water and other water systems from microorganisms.

PT 12- Slimicides: This group is used to control or prevent the formation of slime in equipment and materials used in industrial areas.

PT 13- Working or cutting fluid preservatives: These products aim to control microbial deterioration in fluids used for metal, glass or other materials.

Pest control

PT 14- Rodenticides: These products aim to control of rodents, except for attraction or repulsion.

PT 15- Avicides: Purpose of use is to control birds, except for attraction or repulsion.

PT 16- Molluscicides, vermicides and products to control other invertebrates: Used for the control of invertebrates such as molluscs, worms and other, by means other than attraction or repulsion.

PT 17- Piscicides: These products purpose is to control of fish.

PT 18- Insecticides, acaricides and products to control other arthropods: These products are used for the control of arthropods such as insects, crustaceans and spiders.

PT 19- Repellents and attractants: These are used to control harmful organisms by attracting or repelling, are used for directly on the skin or indirectly in environments of human or animal.

PT 20- Control of other vertebrates: These products are used for controlling vertebrates other than those included by the other product types.

Other biocidal products

PT 21- Antifouling products: These products aim to control the growth of fouling microorganisms, plant or animal species on vessels, aquaculture equipment or other construction used in water.

PT 22- Embalming and taxidermist fluids: These products are used for the disinfection and preservation of human or animal corpses and parts.

Conformity tests of biocidal products

Physical and Chemical Properties of the Active Substance

When evaluating the physicochemical properties of the active substances in the biocidal product, the appropriate method should be selected and experimental results given priority, provided that it operates within the validity range. Data on physicochemical properties should be reliable¹⁵.

Physical parameters

Appearance: The physical parameters of the product should be measured and reported at 20 °C ambient temperature, and 101.3 kPa atmospheric pressure. The physical state of the product may be solid, liquid, or gaseous. The colour and odour must be reported at parameters mentioned above, Odour associated with the active substance described in laboratories or production plants, must be reported. This can be e.g. odourless, characteristic of aromatic compounds, ammonia-like, biting, faint, pungent, slight, sweetish or

other. Substances that are hazardous by inhalation should not be investigated for their odour properties¹⁵.

Melting and freezing points: The melting point must be measured up to 360 °C. Generally, it should be determined if the freezing point of liquid substances is above -20°C. EC method A.1 (Melting / Freezing Temperature) should be used as the test method. The use of Differential Scanning Calorimetry (DSC) or Differential Thermo Analysis (DTA) is recommended¹⁶.

Acidity and alkalinity: For water-containing active substances, the pH of the active substance itself must be tested according to the CIPAC method MT 75.3¹⁷. If solid and nonaqueous liquid active substances are to be used in biocidal products, the pH of an aqueous dilution of 1% of the active substance should be determined. Acidity (pH <4) or alkalinity (pH >10) of the products should be determined according to the CIPAC method MT 191¹⁸.

Boiling point: The boiling point must be measured up to 360 °C according to EC method A.2 (Boiling temperature)¹⁹. The boiling point should be measured at of 101.3 kPa (Standart atmospheric pressure) unless decomposition occurs.

Density: Density for liquids and solids is tested according to OECD Test Guideline 109 (Density of Liquids and Solids)²⁰. The relative density of gases can be calculated from their molecular weight and the Ideal Gas Law. Polymer density should be determined by buoyancy methods.

Additional physical tests may be performed depending on the product type.

Chemical Properties:

Approved analytical methods should be used for the determination of active substances and, where necessary, residues. Methods should be able to fully characterize analytes and quantitatively identify them. Therefore, the methods should be validated by laboratories.

Table 1: Tolerance limits of the active substance content.

Declared nominal content of active [g/kg or g/L]	Tolerance limit
≤25	±15% (homogenous formulations) ±25% (non-homogenous preparations)
25 - 100	±10%
100 - 250	±6%
250 - 500	±5%
> 500	±2.5%

The method validation parameters should include recovery, repeatability, selectivity/specificity, limit of detection and limit of quantification. Methods should avoid hazardous substances and where possible, use commonly available techniques/equipment. The active substance analysis is carried out using titrimetric, spectroscopic or chromatographic methods according to the chemical properties of the substance. HPLC, GC, GC-MS, AAS, ICP systems are the most commonly used analytical methods. In the analyzes, it is evaluated

whether the content of the active substance is in accordance with the manufacturer's declaration. It is accepted that the content of the active substance will vary with each batch and as a result of sampling and analytical errors. These changes must be within a certain limit. The tolerance limits applied to the active substances are given in the Table 1¹⁵.

Storage stability tests:

Tests to prove that the biocidal product is stable during storage and shelf life.

Accelerated storage test:

Stability tests are performed with CIPAC method MT 46.3 accelerated stability results are used to indicate that the biocidal product will remain stable over a two-year shelf life. Accelerated stability data are also indicative of the stability of the biocidal product when exposed to a higher temperature than optimum temperatures. If the biocidal product should not be stored at high temperatures, the label should contain a warning statement. If the active substance is sensitive to high temperature, the accelerated stability test can be carried out at lower temperatures for longer periods²¹.

Long term storage test:

Long term stability test is carried out at 25°C for 2 years to determine whether the product will remain stable in the commercial packaging during the shelf life. During this time the condition of the product is supported by chemical analysis data.

Low temperature stability test (liquids):

The relevant test method for low temperature stability is the CIPAC method MT 39.3. The product is stored for seven days at 0 °C, and its stability is examined. Some types of formulations may need to be investigated for stability to freeze/thaw²².

Efficacy Tests:

To determine the claimed efficacy of the biocidal product or active substance, its biological/microbiological activity needs to be evaluated using appropriate methods. Efficacy is described as the ability of a biocidal product to perform the claimed activity when used according to the instructions recommended on the product label. Studies should be conducted using appropriate methods to demonstrate that the product is sufficiently effective against organisms under conditions of use (concentration, duration, application method, etc.).

There are different types of studies to determine the efficacy of the product²³.

Screening tests:

Screening tests are generally not related to practical/field conditions and are only performed with the active substance. Therefore, such tests are used during product development.

Laboratory studies: These are the studies performed according to the standard criteria for determining the efficiency in the laboratory.

Simulation tests in the laboratory:

Simulation tests are more appropriate to demonstrate effectiveness. It simulates the field conditions where the product will be used under laboratory conditions. For example, for a product intended for disinfection of hard surfaces, a suspension test and surface test with the relevant EN standards is sufficient.

Field tests: Field tests are a good indicator to see how the efficacy of the biocidal product is affected by field conditions. The results of biocidal treated samples or areas are compared with those of untreated control samples or areas.

Efficacy tests should be carried out in accordance with CEN, ISO, OECD, ASTM standard protocols. If standard methods are not available, validated in-house methods can be used. For the disinfectant group, the product must have at least "cidal" effect on the target organisms of the relevant standard when tested according to the phase 1 and phase 2 step 1 suspension methods of the EN standards. An appropriate method should be selected considering the area where the disinfectant is used²⁴⁻²⁷. For preservatives group, biocidal activity is mostly a static activity showed on challenge tests on some target organisms, in the related product matrix. When a curative effect is claimed, it is sufficient to show that the microbial reduction in the treated samples is significantly greater than that of the untreated control samples. For pest control products, only biological activity can be demonstrated for a target organism (eg, control of mice or mosquito control).

Testing for Skin Irritation:

Disinfectants used for human hygiene and insecticides that contact with the skin should be tested for irritation. *In vitro* and *in vivo* assays are used to study the skin irritation potential of a biocidal product²⁸. The EC method B.46 and OECD Test Guideline 439, known as Reconstructed Human Epidermis Model Test, are the methods used for *in vitro* skin irritation²⁹⁻³⁰. *In vivo* tests can be used if there are certain limitations to conducting the *in vitro* test to study the irritation potential of the biocidal product. EC method B.4 Acute Toxicity: Dermal Irritation/Corrosion, OECD Test Guideline 404: Acute Dermal Irritation/Corrosion test methods may be used³¹⁻³².

CONCLUSION

As a conclusion, the use of chemical biocides is a fundamental protection in the prevention and control of microbial growth in medical, veterinary, domestic and industrial environments. Biocides are used a great extent in the healthcare environment for the disinfection of equipments, surfaces, water, and for antisepsis of skin and wound. Misuse of biocidal products brings about some problems such as resistance development and toxicity. The spreading usage of products containing low concentrations of commonly used biocides has raised some concerns about the possible development of microbial resistance. Laboratory studies have demonstrated that bacteria can become resistant to a biocide, and that resistant bacteria can develop cross-resistance to other biocides and antibiotics. High concentrations of biocides generally have toxic effects not only for humans but also for the environment. In this context biocidal products need to be enlarged in regulations and control in terms of traceability due to their extensive and increase in use.

AUTHOR'S CONTRIBUTION

The manuscript was carried out, written, and approved in collaboration with all authors.

ACKNOWLEDGEMENTS

The authors extend their thanks and appreciation to the Turkish Medicines and Medical Devices Agency to provide necessary facilities for this work.

CONFLICT OF INTEREST

No conflict of interest associated with this work.

REFERENCES

- Commission Regulation, EC 2012. Regulation (EU) No 528/2012 of the European Parliament and of the Council of 22 May 2012 concerning the making available on the market and use of biocidal products.
- Ribeiro M, Simões LC, Simões M. Biocides. Reference Module in Life Sciences, Encyclopedia of Microbiology 4th ed. Elsevier Inc 2018; 478-490.
- Maillard JY. Antimicrobial biocides in the healthcare environment: efficacy, usage, policies, and perceived problems. *Ther Clin Risk Manag* 2005; 1(4):307-320. PMID: 18360573
- Russell AD. Introduction of biocides into clinical practice and the impact on antibiotic-resistant bacteria. *J Appl Micro* 2002; 92:121-35. <https://doi.org/10.1046/j.1365-2672.92.5s1.12.x>
- Holah J, Taylor J, Dawson D and Hall K. Biocide use in the food industry and the disinfectant resistance of persistent strains of *Listeria monocytogenes* and *Escherichia coli*. *J Appl Micro* 2002; 92:111-120. <https://doi.org/10.1046/j.1365-2672.92.5s1.18.x>
- Bradley EL, Castle L, Chaudhry Q. Applications of nanomaterials in food packaging with a consideration of opportunities for developing countries. *Trends Food Sci Tech* 2011; 22(11):604-610. <https://doi.org/10.1016/j.tifs.2011.01.002>
- Williams, TM. The Mechanism of Action of Isothiazolone Biocide. NACE International Conference Paper, 2006.
- Fang F, Bernigaud C, Candy K, Melloul E, Izri A, Durand R, Botterel F, Chosidow O, Huang W, Guillot J. Efficacy assessment of biocides or repellents for the control of *Sarcoptes scabiei* in the environment. *Parasit Vectors* 2015; 8:416.
- Russell AD. Biocide use and antibiotic resistance: the relevance of laboratory findings to clinical and environmental situations. *The Lanc Infect Dis* 2003; 3(12): 794-803.
- Coors A, Vollmar P, Heim J, Sacher F, Kehrer A. Environmental risk assessment of biocidal products: identification of relevant components and reliability of a component-based mixture assessment. *Environ Sci Eur* 2018; 30(1):3. <https://doi.org/10.1186/s12302-017-0130-0>
- Vyas A, Pickering CAC, Oldham LA, Francis HC, Fletcher AM, Merrett T, Niven RMCL. Survey of symptoms, respiratory function, and immunology and their relation to glutaraldehyde and other occupational exposure among endoscopy nursing employee. *Occupational Environmental Medicine* 2000; 57:752-759. <https://doi.org/10.1136/oem.57.11.752>
- Rutala WA, Weber DJ. The benefits of surface disinfection. *AJIC* 2004; 32:22631. <https://doi.org/10.1016/j.ajic.2004.04.197>
- Rideout K, Teschke K, Dimich-Ward H, Kennedy SM. Considering risks to healthcare workers from glutaraldehyde alternatives in high-level disinfection. *J Hosp Infect* 2005; 59:4-11. <https://doi.org/10.1016/j.jhin.2004.07.003>
- ECHA 2014, Transitional Guidance on Regulation (EU) No 528/2012 of the European Parliament and of the Council of 22 May 2012 concerning the making available on the market and use of biocidal products (Biocidal Products Regulation, the BPR). European Chemicals Agency, Helsinki, Finland, May 2014.
- ECHA 2018, Guidance on the Biocidal Products Regulation Volume I: Identity of the active substance/physico-chemical properties/analytical methodology–Information Requirements, Evaluation and Assessment. Parts A+B+C
- Commission Regulation EC No 440/2008 laying down test methods pursuant to Regulation EC No 1907/2006 of the European Parliament and of the Council on the Registration, Evaluation, Authorisation and Restriction of Chemicals (REACH) (originally published as Council Regulation EC No 440/2008, corrigendum according to Official Journal of the European Union dated June 3, 2008, L 143/55), Annex, Part A, A.1. Melting/Freezing Temperature, Official Journal of the European Union, L 142, 51.
- Collaborative International Pesticide Analytical Council, Ltd. CIPAC Handbook, MT 75.3 Determination of pH Values, CIPAC, Hatching Green, Harpenden, Hertfordshire, England, 1999.
- Collaborative International Pesticide Analytical Council, Ltd. CIPAC Handbook, MT 191 acidity or alkalinity of formulations CIPAC, Hatching Green, Harpenden, Hertfordshire, England, 2005.
- Official Journal of the European Communities L 383 A, 15-21, 1992.
- OECD Guidelines for the Testing of Chemicals, Section 1, Test No. 109: Density of liquids and solids, 2012.
- Collaborative International Pesticides Analytical Council Ltd. (CIPAC) Handbook, “MT 46.3 Accelerated Storage Procedure”, (Dobrat & Martin, Eds.) Published by Black Bear Press Ltd., 2000.
- Collaborative International Pesticides Analytical Council Ltd. (CIPAC) Handbook, “MT 39.3 Low temperature stability of liquid formulations”, (Dobrat & Martin, Eds.) Published by Black Bear Press Ltd., 2000.
- ECHA (2018) Guidance on the Biocidal Products Regulation Volume II: Efficacy Part A: Information Requirements.
- European Committee for Standardization. Chemical disinfectants and antiseptics – Quantitative suspension test for the evaluation of bactericidal activity of chemical disinfectants and antiseptics used in food, industrial, domestic, and institutional areas–test method and requirements, (phase 2/step 1) European Standard EN 1276, 2010.
- European Committee for Standardization. Chemical disinfectants and antiseptics– Quantitative suspension test for the evaluation of fungicidal activity of chemical disinfectants and antiseptics used in food, industrial, domestic, and institutional areas – Test method and requirements, (phase 2/step 1) European Standard EN 1650+A1, 2013.
- European Committee for Standardization. Chemical disinfectants and antiseptics– Quantitative suspension test for the evaluation of bactericidal activity for instruments used in the medical area. Test method and requirements (phase 2/step 1), European Standard EN 13727+A2, 2015.
- European Committee for Standardization. Chemical disinfectants and antiseptics– Quantitative suspension test for the evaluation of fungicidal activity of chemical disinfectants for instruments used in the medical area - Test method and requirements (phase 2, step 1), European Standard EN 13624, 2013.
- ECHA 2018 Guidance on the Biocidal Products Regulation Volume III: Human health Part A: Information Requirements Version 1.2 May 2018.
- EC Testing Method B.46. *In vitro* skin irritation: reconstructed human epidermis model test. Council Regulation EC No 761/2009
- OECD Guidelines for the Testing of Chemicals. Test No. 439: *In vitro* skin irritation– Reconstructed human epidermis test method, 2013.
- OECD Guidelines for the Testing of Chemicals. Test No. 404: Acute dermal irritation/corrosion, 2002.
- EC Testing Method B.4. Acute toxicity: dermal irritation/corrosion. Council Regulation (EC) No 440/2008– Part B.