Available online on 15.1.2020 at <http://ujpr.org>
Universal Journal of Pharmaceutical Research
 An International Peer Reviewed Journal
 Open access to Pharmaceutical research

This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial Share Alike 4.0 License which permits unrestricted non commercial use, provided the original work is properly cited

Volume 4, Issue 6, 2019

Open Access

REVIEW ARTICLE

ETHNOBOTANY, PHYTOCHEMISTRY AND PHARMACOLOGY OF *OCHNA SCHWEINFURTHIANA*: A REVIEW

Yusuf AJ^{1*}, Abdullahi MI¹, Muhammad AA²

¹Department of Pharmaceutical and Medicinal Chemistry, Usmanu Danfodiyo University, Sokoto, Nigeria.

²Department of Pharmaceutical and Medicinal Chemistry, Ahmadu Bello University, Zaria, Nigeria.

ABSTRACT

Ochna schweinfurthiana (Os, Family: Ochnaceae) is a small evergreen tree used in ethnomedicine to treat different ailments; it is also used in agri-horticulture and as ornaments, dyes among others. It is a rich source of complex dimers of flavonoids and used for treatment of pain, inflammation, and arthritis. Chemical investigations carried out on the different parts of the plant have been confined to phenolic compounds majorly, bioflavonoids, glycosides, steroids and terpenes. The plant, *O. schweinfurthiana* have shown a wide spectrum of biological and pharmacological properties which include antimicrobial, cytotoxic/antiproliferative, genotoxicity, antinociceptive, anti-inflammatory, antioxidant and antiplasmodial. This review comprehensively summarizes the potential effects of the plant *O. schweinfurthiana*, chemically and pharmacologically. However, more researches in the aspect of phytochemical and biological studies are needed to exhaustively isolate bioactive compounds and evaluate their effects on other ailments as claimed by the traditional healers.

Keywords: Anti-microbial, antimalarial, flavonoids, *Ochna schweinfurthiana*, Ochnaceae, phenolics, pharmacological.

Article Info: Received 2 November 2019; Revised 19 December; Accepted 28 December, Available online 15 January 2020

Cite this article-

Yusuf AJ, Abdullahi MI, Muhammad AA. Ethnobotany, phytochemistry and pharmacology of *Ochna schweinfurthiana*: a review. Universal Journal of Pharmaceutical Research 2019; 4(6):75-79.

DOI:<https://doi.org/10.22270/ujpr.v4i6.341>

Address for Correspondence:

Yusuf AJ, Department of Pharmaceutical and Medicinal Chemistry, Usmanu Danfodiyo University, Sokoto, Nigeria. Tel: 08036386793, E-mail: amynajega@gmail.com

INTRODUCTION

Ochna schweinfurthiana (Os) belonging to the Ochnaceae family is a small tree that was named after a German botanical collector and taxonomist Dr. Georg August Schweinfurth; it is a small tree that measures up to 4 m tall; the plant is commonly known as the brick-red *Ochna* in English, Jan-taru in Hausa language, Hiéké in Yoruba and Sa'aboule in Foulfouldé^{1,2}. The plant has found use as medicine, for agricultural, social and religious purposes¹. This review centered on the ethnobotanical, phytochemical and pharmacological properties of *O. schweinfurthiana*.

Botanical Description

Ochna originated from a Greek word "*Ochne*" which means wild pear and it was named by Linnaeus in 1951 as *Ochna* because of the resemblance of their leaves with those of wild pear³. It is an old world genus of mainly trees, shrubs and shrublets which comprises of about 85 species⁴ and it is widely distributed in tropical Asia, Africa and America⁵ of which eleven (11) species are found in India⁶. *Ochna*'s are usually called Mickey Mouse plants, because of the appearance of the black drupelets fruits. The Ochnaceae family is mainly composed of trees and shrubs with an estimated 33

genera and 550 species⁷ well distributed around the world especially in tropical Africa, Australia, Madagascar Asia, the Mascarene Island and America⁸. They are notably known for their unusual shiny leaves, with parallel veins that are closely spaced, alongside toothed margins with conspicuous stipules^{1,7}. The largest genera are *Ouratea*, *Ochna*, *Campylospermum*, *Sauvagesia* and *Quiina* with (200, 85, 65, 39 and 34 species) respectively (Table 1).

Table 1: Ochnaceae subfamilies and their estimated number of species.

Subfamily	Estimated number of species
Ouratea	200
Ochna	85
Campylospermum	65
Sauvagesia	39
Quiina	34
Total	423

Morphology

O. schweinfurthiana is a small evergreen tree or shrub that grows up to 4 m tall and it has a dark grey bark that is fissured and cracked, separating into square

segments¹⁰. The leaves are olive-green (1-13.5x1.7-5.5cm) that oblanceolate to oblong or elliptic, apex somewhat rounded, base tapers into the petiole, margins rather bluntly toothed (serrulate), sometimes appearing almost scalloped, net-veining conspicuous on the upper surface and young leaves are coppery^{1,11}. It bears bright yellow flowers (1.5 cm diameter) which are sweetly-scented from September to November, very short-lived, normally appearing before or with the young leaves¹⁰. In addition, it appears in a condensed raceme with 4–10 flowers on a short central stem and the petals fall very early¹¹.

Figure 1: Leaf and fruits of *O. Schweinfurthiana*.

The fruits of *O. schweinfurthiana* are 1-5 oval appear between August and January, attached at the base are 2-4 black berries when ripe; they are enlarged, borne on brick-red persistent sepals turning cherry to brick-red. The bark is dark grey, thick, and deeply fissured into a grid-like pattern^{10,11}.

Figure 2: Whole plant of *O. Schweinfurthiana*.

Taxonomy

Kingdom: Plantae Order: Malpighiales
Family: Ochnaceae Genus: Ochna
Subfamily: Ochnoideae
Species: *O. schweinfurthiana*⁹

Common names

English: Brick-red Ochna, Hausa:Jan-taru
Yoruba: Hiéké, Fulfulde:Sa'aboule

Habitat, Distribution and Ecology

The plant grows in open deciduous woodland in tropical regions in Africa from Guinea to southern and northern Nigeria and across central Africa to Sudan, Uganda, Zimbabwe, Mozambique, Tanzania and Angola. It has medium water requirement when young and grows fast, flowers from September to November. It required low maintenance and attracts insects and birds^{4,10}.

Ethnomedicinal uses

Several preparations (powdered and decoctions) of the leaves and/or root of the *O. schweinfurthiana* have found general use as antimicrobial (wound dressing, eye infection), analgesic, anti-inflammatory and anthelmintic agents¹. The leaf is also used as an antiseptic, stimulant, febrifuge, laxative, enemaetc¹. In Northern Cameroon, *O. schweinfurthiana* is used to treat different diseases such as rubella, burns, stomachache and multiple sclerosis¹²; the root of the plant is also used in the treatment of stomach and eye aches as well as headache, while the leaves are used for toothaches treatment².

The pulverized bark is used to treat malaria, febrifuges and as anthelmintic, while the decocted root leaves and/or bark is used in wound dressing¹². In Northern Nigeria, the *O. schweinfurthiana* is used to treat typhoid fever, measles and fungal skin infections¹². The macerated roots of *O. schweinfurthiana* has been reportedly used to induce/speed the delivery process and for miscarriage¹³.

Other uses

The plant is used in agri-horticulture; the bark and flowers *O. schweinfurthiana* are cultivated for ornaments, dyes, stains, inks, tattoos and mordant among others. The wood is used for farming, forestry, hunting and fishing apparatus. The leaf has social, religious, superstitious and magic values among others¹.

Phytochemical screening

Abdullahi et al.,¹² reported the presence of flavonoids, steroids/terpenes and saponins in the acetone leaf extract of *O. schweinfurthiana* and the methanol leaf extract indicated the presence of flavonoids and saponins. However, flavonoids, saponins, glycosides, tannins and steroids/terpenes were reported on the methanol stem extract of *O. schweinfurthiana*¹⁴. A study conducted by Ibrahim et al., reported the presence of carbohydrates, steroids/triterpenes, glycosides, saponins, tannins and flavonoids in the methanol leaf extract of *O. schweinfurthiana*¹⁵.

Bioactive constituents

Chemical constituents isolated from *O. schweinfurthiana* fall under the following class of secondary metabolites phenolics-flavonoids, and glycosides. *O. schweinfurthiana* have been reported to contain phenolic derivatives (such as flavonoids, bioflavonoids) which appear as free or in polymerized forms. Isolation and characterization of quercetin-3-O-β-D-glucopyranosyl-(1→6)-α-rhamnoside (quercetin rutinoid) from the *n*-butanol soluble fraction of methanolic leaf extract of *O. schweinfurthiana* was reported¹⁶. A novel biflavonoid compound, tri-methoxy lophirone was isolated from the chloroform soluble fraction of the methanol root extract of *O. schweinfurthiana*¹⁷.

Ndongo et al.,¹⁸ reported the isolation of seven flavonoids, hemerocallone, 6,7-dimethoxy-3',4'-dimethoxyisoflavone, amentoflavone, agathisflavone, cupressuflavone, robustaflavone, and epicatechin, and three other chemical constituents, lithospermoxide, β-D-fructofuranosyl-α-D-glucopyranoside and 3β-O-D-glucopyranosyl-β-stigmasterol from the ethyl acetate the stem bark extract of *O. schweinfurthiana*.

Figure 3: Chemical constituents isolated from *O. Schweinfurthiana*.

The roots of *O. schweinfurthiana* were reported to contain three new compounds viz; 4'''-methoxylophirone A, 4,4',4'''-trimethoxylophirone A. In addition, six known compounds were also isolated and characterized, including calodenone, calodenine B, lophirone A, gerontoisoflavone A, and 3β-O-D-glucopyranosyl-β-sitosterol². Six known compounds were isolated from the powdered bark of *O.*

schweinfurthiana and they include, hemerocallone, 6,7-dimethoxy-3'-4'-dimethoxyisoflavone, lithosperm - oside, amentoflavone, agathisflavone and β-D-fructofuranosyl-α-D-glucopyranoside¹⁹.

BIOLOGICAL AND PHARMACOLOGICAL ACTIVITIES

Antimicrobial activity

The acetone and methanol leaf extracts of *O. schweinfurthiana* had a remarkable antibacterial effect against *S. aureus*, *S. typhi*, *K. pneumonia* and *P. aeruginosa* with a zone of inhibition ranging from 15 – 21 mm; the extracts had an MIC and MBC values of 10–20 mg/ml and 20–40 mg/ml, respectively¹². Quercetin-3-*O*- β -D-glucopyranosyl(1 $\rightarrow$ 6)- α -rhamnoside from the *n*-butanol soluble fraction of the methanolic leaf extract of *O. schweinfurthiana* showed an *in vitro* inhibitory effect against some bacterial isolates such as *S. aureus*, MRSA, *S. pyogenes*, *E. coli*, *K. pneumoniae*, *S. typhi* and *P. aeruginosa* with an MIC and MBC range between 2.5–5.0 and 5–20 μ g/ml, but there was no effect against *B. subtilis*, *C. ulcerans* and *C. albicans*¹⁶. Earlier studies showed that trimethoxy lophirone A from the chloroform soluble fraction of the methanol root extract of *O. schweinfurthiana* inhibited the growth of some selected human pathogens including *S. aureus*, *S. pyogenes*, *P. aeruginosa*, *K. pneumoniae* and *S. typhi* with an MIC and MFC values of 5 μ g/ml and 20 μ g/ml, respectively¹⁷. Crude methanol stem extract of *O. schweinfurthiana* and its chloroform (CF), ethylacetate (EF) and *n*-butanol (BF) fractions inhibited the growth of MRSA, *S. aureus*, *S. pyogenes*, *S. typhi*, *S. dysenteriae*, *K. pneumoniae*, *N. gonorrhoea*, *P. aeruginosa*, *C. albicans*, *C. tropicalis*; the mean zone of inhibition of extract and fractions ranges from 20–29 mm; moreover, chloroform fraction showed greater antimicrobial activity with an MIC value of 1.25 mg/ml against all the test organisms except *P. aeruginosa*¹⁴.

Cytotoxic and Antiproliferative effect

The methanol and ethylacetate stem bark extracts of *O. schweinfurthiana* demonstrated cytotoxicity against HeLa cells; amentoflavone and agathisflavone were also active¹⁸. Antiproliferative effect of *O. schweinfurthiana* extract was evaluated against *Glioblastoma multiforme* (GBM U-1242 MG) cell line and the extract reduced cell count by 20 % with an IC₅₀ value 823.51 μ g/ml²⁰. The aqueous stem bark of *O. schweinfurthiana* did not show any cytotoxic effect on Vero monkey kidney cell line after 48 h incubation with an LC₅₀ value 50 $\pm$ 1 μ g/ml¹⁹.

Genotoxicity

Djova et al., reported that the extracts of *O. schweinfurthiana* were non genotoxic in a study they carried out; as none of the plant extracts demonstrated a dose dependent increase or revertent colonies $\geq$ the number of negative control revertent colonies; thus, the plant *O. schweinfurthiana* may not contain any genotoxic substances that may lead to mutations either by substitution or by reversion in the genetic material¹⁹.

Antinoceptive and anti-inflammatory effect: The methanol leaf extract of *O. schweinfurthiana* significantly inhibited the writhing response induced by acetic acid in a dose dependent manner; the highest dose exhibited maximum inhibition of pain (84.3 %). In addition, the extract was also able to attenuate pain response in a similar manner though with a slower onset of action in the tail flick model¹⁵. The aqueous bark extract of *O. schweinfurthiana* exhibited good anti-inflammatory effect in both ferrous oxidation-Xylenol Orange (Fox) and BSA denaturation assays;

the extract demonstrated good 15-lipoxygenase inhibitory effect with an IC₅₀ value of 32.2 $\pm$ 0.36 μ g/ml, however, an IC₅₀ of 130 $\pm$ 5.78 μ g/ml was recorded by the extract in the inhibition of heat induced albumin denaturation¹⁹.

Antioxidant effect

Messiet et al., evaluated the antioxidant activity of some compounds including 4"-methoxylophirone A, calodenone, calodenine B, lophirone A, gerontoisoflavone A from the roots of *O. schweinfurthiana* using DPPH radical scavenging and ferric reducing-antioxidant power (FRAP) assays². In the DPPH radical scavenging assay, calodenine B showed prominent effect with SC₅₀=0.17 $\pm$ 0.04 μ M and EC₅₀=4.25 μ M, gerontoisoflavone A exhibited weak activity in all the models applied with SC₅₀=19.00 μ M and SC₅₀=78.67 μ g EAA/mg/dw in DPPH and FRAP respectively. The antioxidant property of the leaf, stem bark and fraction of *O. schweinfurthiana* was evaluated²¹.

Antiplasmodial effect

An *in vivo* study showed that the methanol leaf extract of *O. schweinfurthiana* exerted a suppressive effect against *Plasmodium berghei* at a lower dose (50 mg/kg); Ibrahim et al., concluded that, the extract possess blood schizonticidal activity as it was able to suppress malaria at the early stage²². Moreso, the extract reduced the level of parasitaemia with 100 % cure at the lowest dose (50 mg/kg); the percentage inhibition of parasitaemia was higher than the chemo suppression which might be due to non-selectivity of the extract to the proliferative process of the parasite^{22,23}.

Antiplasmodial effect of the ethylacetate roots extract of *O. schweinfurthiana* and some compounds including 4"-methoxylophirone A, (4E,7Z)-3,8-dicarboxy-1-(*O*- β -D-glucopyranosyl-(1 $\rightarrow$ 6))-*O*- β -D-glucopyranosyl-2,9 dihydroxyhexeicos-4,7-diene, calodenine B, lophirone A and gerontoisoflavone A were investigated *in vitro*; 4"-methoxylophirone A showed good antiplasmodial effect against *P. falciparum* strain 3D7; this effect as explained by Messi et al.,² might be related to the presence of a methoxy group on position C-4" which has been known to enhance lipophilicity thereby enhancing its movement into the cells²⁴; other compounds were found to be inactive². Cold and hot aqueous leaf extracts of *O. schweinfurthiana* possess inhibitory effect against *P. falciparum in vitro*; thus there was significant reduction of parasitaemia.

The high dose (80 μ g/ml) exhibited 86.42 % (cold extract) and 85.06 % (hot extract) reduction of parasitaemia. On the other hand, no significant difference was observed on the plasmodium lactate dehydrogenase (PLDH) activity of the treated extract when compared with the standard drug²⁵.

Toxicity

Toxicity level of *O. schweinfurthiana* was assessed in mice both orally and intraperitoneally. The methanol leaf extract of the plant produced an LD₅₀ 774.6 mg/kg, i.p. while the oral LD₅₀ value for the extract was about 5000 mg/kg; according to this study, the leaf of *O. schweinfurthiana* is intraperitoneal toxic and orally safe²².

CONCLUSION

O. schweinfurthiana exhibit a variety of biological effects; the plant is considered to be effective against cancer, malaria, oxidative stress, pain, inflammation and a wide range of activity against microbes; thus, the pharmacological actions have been attributed to the presence of different classes of secondary metabolites such as biflavonoids, glycosides, steroids and terpenes among others. In addition, the mechanisms of action of the observed effects and evaluation of other pharmacological properties of *O. schweinfurthiana* need closer attention and it should be the objective of new researches on *O. schweinfurthiana*.

AUTHOR'S CONTRIBUTION

All authors have worked equally for this work.

ACKNOWLEDGEMENTS

The authors extend their thanks and appreciation to the Usmanu Danfodiyo University, Sokoto, Nigeria to provide necessary facilities for this work.

CONFLICT OF INTEREST

No conflict of interest associated with this work.

REFERENCES

- Burkill MH. The useful plants of west Tropical Africa, Families J-L. Royal Botanic Gardens Kew 1985; 4: 275.
- Messi AN, Mbing JN, Ndongo JT, Nyegue MA, Tchinda AT, Yemeda MF, Pegnyemb DE. Phenolic compounds from the roots of *Ochna schweinfurthiana* and their antioxidant and antiparasitic activities. *Phytochemistry Lett* 2016; 17: 119-125. <https://doi.org/10.1016/j.phytol.2016.07.011>
- Muema MJ. Phytochemical and antimicrobial investigation of *Ochnathomasiana* Engl. and Gilg. MSc Thesis, Department of Chemistry, Kenyatta University 2015; 30– 35. http://dx.doi.org/10.1093/jac/48.suppl_1.5
- Verdcourt B. Ochnaceae. Flora of tropical east Africa. Royal Botanic Gardens, Kew, Richmond, United Kingdom 2005;60.
- Rendle AB. The classification of flowering plants. Cambridge University Press, London. 1952; 2:12-18.
- Kirtikar KR, Bussu BD. Indian medicinal plants. Periodical expert book agency, Delhi. 2012; 2:1539.
- Christenhust MJM, Byng JW. The number of known plants species in the world and its annual increase. *Phytotexa* 2016; 261(3): 201-217.
- Mabberley DJ. Mabberley's plant-book. A portable dictionary of plants, their classification and uses. Cambridge university press, Cambridge 2008; 342.
- Simpson DP. Cassell's Latin Dictionary (5 ed.). London: Cassell Ltd. 1979; 883. ISBN 0-304-52257-0.
- Hyde MA, Wursten BT, Ballings P, Coates PM. Flora of Zimbabwe: Species information: *Ochna schweinfurthiana* F. Hoffm. www.zimbabweflora.co.zw/speciesdata/species.php?speciesid.2019; retrieved on 28 September, 2019.
- Anonymous. *Ochna schweinfurthiana*: Plant Book. Available online at www.plantbook.co.za/ochna-schweinfurthiana 2019; retrieved 28 September, 2019
- Abdullahi MI, Iliya I, Haruna AK, Sule MI, Musa AM, Abdullahi SM. Preliminary phytochemical and antimicrobial investigations of leaf extracts of *Ochna schweinfurthiana*. *African J Pharm Pharmacol* 2010; 4(2): 083-086.
- Bruschi P, Morganti M, Mancini M, Signorini MA. Traditional healers and laypeople: A qualitative and quantitative approach to local knowledge on medicinal plants in Muda (Mozambique). *J Ethnopharm* 2011; 138: 543-563. <https://doi.org/10.1016/j.jep.2011.09.055>
- Danmusa UM, Nasir IA, Abdullahi MI, Ahmad AA, Abdulkadir IS. Phytochemical analysis and antimicrobial activities of methanolic stem extracts of *Ochna schweinfurthiana* F. Hoffm. *J Pharmacy Pharmacog* 2015; 3(6): 171-182.
- Ibrahim ZYY, Musa AM, Abdullahi MI, Uba A, Yusuf AJ, Aliyu IM, Ya'u J. Phytochemical and antinociceptive studies on the *Ochna schweinfurthiana* (Ochnaceae). *Int J Advances Pharm Biol Chem* 2015a; 4(4): 838-843.
- Abdullahi MI, Musa AM, Haruna AK, Sule MI, Abdullahi SM, Abdulmalik M, Akinwande Y, Abimiku AG, Iliya I. Anti-microbial flavonoid diglycoside from the leaf of *Ochna schweinfurthiana* Hoffm. (Ochnaceae). *Nigerian J Pharm Sci* 2011; 10(2): 1-7. <https://doi.org/10.1016/j.phytol.2016.07.011>
- Abdullahi MI, Musa AM, et al. Isolation and characterization of anti-microbial biflavonoid from the chloroform-soluble fraction of methanol root extract of *Ochna schweinfurthiana* (Ochnaceae). *African J Pharm Pharmacol* 2014; 8(4): 93-99. <https://doi.org/10.1076/phbi.41.2.92.14245>
- Ndongo JT, Issa ME, Messi AN, Ngo Mbing J, Cuendet M, Pegnyemb DE, Bochet CG. Cytotoxic flavonoids and other constituents from the stem bark of *Ochna schweinfurthiana*. *Natural Prod Res* 2015; 17:1684-1687. <https://doi.org/10.1080/14786419.2014.991321>
- Djova SV, Nyegue MA, Messi AN, Afagnigni AD, Etoa FX. Phytochemical study of aqueous extract of *Ochna schweinfurthiana* F. Hoffm powder bark and evaluation of their anti-inflammatory, cytotoxic and genotoxic properties. *Hindawi Evidence-Based Complementary and Alternative Medicine*, 2019; <https://doi.org/10.1155/2019/8908343>
- Abdullahi MI, Musa AM, Tajuddeen N, Mohammed GM, Yusuf AJ, Iliya I. Anti-proliferative study and isolation of Ochnaflavone from the ethylacetate-soluble fraction of *Ochnakibbiensis* Hutch and Dalziel. *Nat Prod Res* 2016; <https://doi.org/10.1080/14786419.2016.1274892>
- Nyegue MA, Ngo Mbing J, Djova SV, et al. Evaluation of the antioxidant activity of the leaves, stem-barks extracts and fractions of *Ochna schweinfurthiana* F. Hoffm (Ochnaceae). *African J Pharm Pharmacol* 2016; 10(17): 370–378. <https://doi.org/10.1155/2019/8908343>
- Ibrahim ZYY, Abdullahi MI, Musa AM, Maje IM, Yusuf AJ, Aliyu IM. Acute toxicity profile and *Plasmodium berghei* inhibitory activity of *Ochna schweinfurthiana* (F. Hoffm) Ochnaceae leaf extract in Laboratory animals. *Int J Pharma Sci Res* 2015b; 6(10):1302–1306.
- Salawu OA, Tijani AY, et al. Anti-malarial activity of ethanolic stem bark extract of *F. bialalbida* (Del) a. Chev (Mimosoidae) in mice. *Arc App Sci Res* 2010; 2: 261-268. <https://doi.org/10.1155/2011/618209>
- Monks MR, Ferraz A, Bordignon SA, Machado KR, Richter MF, Da Rocha AB. *In vitro* cytotoxicity of extracts from Brazilian Asteraceae. *Pharm Biol* 2002; 40: 494-500. <https://doi.org/10.1076/phbi.40.7.494.14681>
- Omoniwa BP. *In-vitro* antiparasitic activity of cold and hot aqueous extracts of *Ochna schweinfurthiana* leaf. *J Bacteriol Parasitol* 2017; 8(5 Suppl): 30.