Available online on 15.09.2020 at <http://ujpr.org>
Universal Journal of Pharmaceutical Research
 An International Peer Reviewed Journal

Open access to Pharmaceutical research
 This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial Share Alike 4.0 License which permits unrestricted non commercial use, provided the original work is properly cited

Volume 5, Issue 4, 2020

Open Access

RESEARCH ARTICLE

PREVALENCE OF INTESTINAL HELMINTHIASIS AND THEIR ASSOCIATION WITH EOSINOPHILIA AMONG SCHOOLCHILDREN IN WADIDHAHAR DISTRICT AT SANA'A GOVERNORATE, YEMEN

Arwa Mohammed Othman¹, Abdulsalam Mohammed Al-Mekhalafi²

¹Department of Microbiology and Immunology, Faculty of Medicine and Health Sciences, Sana'a University, Sana'a, Yemen.

²Parasitology Department, Faculty of Medicine and Health Sciences, Sana'a University, Sana'a, Yemen.

ABSTRACT

Objectives: Allergy and parasitic infections are common causes of blood eosinophilia. Intestinal helminthiasis remains a major health problem in many developing countries. Eosinophils are an effector immune cell against parasites. The objective of this study was to determine the prevalence of intestinal helminthiasis and eosinophilia among primary schoolchildren in Wadi Dhahr district, Sana'a governorate, Yemen.

Subjects and methods: Four hundred and fourteen pupils were enrolled in this study. Blood and stool samples were collected from each pupil. Stool samples were examined using Hoffman's sedimentation method for the presence of worms' ova. Blood smears were stained with Giemsa stain and used for counting the eosinophils' percentages.

Results: Out of 414 specimens, the prevalence of intestinal helminths infections was 79(19%) among schoolchildren; (52 (18%) males and 27 (21%) females). Eosinophilia was 134 (32%) observed in the participated pupils; 86 (30%) males and 48 (37%) females. The age group 5-10 years had the highest frequent rate of eosinophilia with a statistically significant difference ($\chi^2=10.4$, $P=0.002$) and 60% of infected students with intestinal worms had eosinophilia ($\chi^2=32.8$, $P<0.001$). Females had a higher rate of eosinophilia than males. The most common helminthiasis infections were hymenolepiasis (10.6%) followed by ascariasis (4.1%), enterobiasis (2.7%), and taeniasis (2.4%). The high rate of eosinophilia was recorded among students infected with *Ascaris lumbricoides* (82%) followed by *Enterobius vermicularis* (73%), *Trichuris trichiura* (67%), *Hymenolepis nana* (59%), and *Taeniasaginata* (30%).

Conclusions: Intestinal helminthiasis still a common health problem among schoolchildren in WadiDhahar district at Sana'a governorate, Yemen, and was found to be associated with high blood eosinophils. Eosinophilia was statistically significant among students infected with ascariasis, hymenolepiasis, and enterobiasis but not with taeniasis and trichuriasis.

Keywords: Intestinal helminthiasis, eosinophilia, schoolchildren, WadiDhar, Yemen.

Article Info: Received 1 July 2020; Revised 3 August; Accepted 5 September, Available online 15 September 2020

Cite this article-

Othman AM, Al-Mekhalafi AM. Prevalence of intestinal helminthiasis and their association with eosinophilia among schoolchildren in WadiDhahar district at Sana'a Governorate, Yemen. Universal Journal of Pharmaceutical Research 2020; 5(4):11-15.

DOI: <https://doi.org/10.22270/ujpr.v5i4.433>

Address for Correspondence:

Arwa Mohammed Othman, Department of Microbiology and Immunology, Faculty of Medicine and Health Sciences, Sana'a University, Sana'a, Yemen, Tel: +967-774486417, E-mail: arwaothman@hotmail.com

INTRODUCTION

Typically, eosinophils' count in the blood is often less than 6% of the total blood leukocyte count. An increase in the eosinophil number, more than 6%, is called eosinophilia. Varied disorders and etiologies may increase blood eosinophils which mainly include worm infections and allergic diseases. Parasitic infections are the most common cause of persistent eosinophilia in developing countries while atopic diseases such as eczema and asthma are the most common causes in developed countries^{1,2}. In some parasitic infections, increase in the eosinophil number may be the only symptom³.

Eosinophilia may vary depending on the parasite development stage, parasite location in the body, parasite load, and co-infections with more than one parasite. Helminths which migrate into the tissues as part of their growth life cycle such as ascariasis induce persistent eosinophilia in both blood and tissues^{4,5}. Globally 1.5 billion individuals were infected with intestinal helminths; of which more than 267 million pre-school children and above 568 million schoolchildren are considered at risk of morbidity, particularly in developing countries⁶. In 2010, an estimated 819 million individuals were infected with *Ascaris lumbricoides*, 464 million with *Trichuris*

trichura, and 438 million with *Ancylostoma* worms in the world⁷. In children, worm infections can cause significant malnutrition, delayed physical development, cognitive impairment while in adults; worms can reduce their productivity^{8,9,10}. Helminth infections have been associated with poor personal hygiene, environmental sanitation, and limited access to potable water¹¹ particularly in Yemen where most of people live under poverty line that lacks the effective programs for parasitic infection control and prevention^{12,13,14}. It was reported that the prevalence rate of intestinal parasitic infection was 62.7% recorded among schoolchildren in Ibb city¹³. Also, a study performed by Al-Mekhlafi *et al.*,¹⁴ reported that the overall prevalence of intestinal parasitic infections was 17.2% among schoolchildren. In Yemen, there are several investigations that focused on the prevalence of intestinal parasitic infection among children school and associated with environmental factors. To date, data on the prevalence of intestinal helminthiasis associated with eosinophilia levels among schoolchildren are not available for Sana'a governorate, Yemen. Therefore, this study aimed to determine the prevalence of intestinal helminthiasis and its association with eosinophilia among schoolchildren at WadiDhahr district in Sana'a governorate, Yemen.

MATERIALS AND METHODS

Study design and area

A cross-sectional survey was conducted between January 2016 to June 2018 at two primary schools (AL-Wahda and AL-Mutanabi) situated in WadiDhahr locating about 15Km north of Sana'a, the capital of Yemen.

Study population

A total of 414 pupils, aged between 5-15 years, were enrolled in this study. Samples were chosen by using a simple random method from each school. One hundred and ninety-six samples were collected from students presented at AL-Wahda school while two hundred and eighteen samples were collected from AL-Mutanabi school.

Exclusion criteria

Students who were taking medications that may affect eosinophilia (e.g. penicillins and cephalosporins) or who had received anti-helminthic drugs within three

months from the beginning of the study. Also, students known to have a food allergy or asthma were excluded from the study.

Ethical considerations

This study was approved by the Faculty of Medicine and Health Sciences, Sana'a University. Also, permission to conduct this study was approved by the office of educational authorities and school heads. Prior samples and data collection, the study objective was clarified to the parents, school teachers, and students. Oral consent was obtained from pupils' parents to participate in this study.

Sample collection

Two samples were obtained from each participant: blood and stool specimens. One gram of stool sample was emulsified in 7ml of 10% formalin for fixation¹⁵. Blood samples were collected by finger prick using disposable lancets. Blood smears were made from capillary blood on a glass slide, left to air dry, and then fixed with absolute methanol¹⁶.

Examination of stool specimens

Stool samples were examined using Hoffman's sedimentation method¹⁵. Helminthic ova are concentrated by passing the fecal suspension through a gauze followed by centrifugation for two minutes at 1000 rpm. The upper liquid phase was discarded using a pipette. Two slides per fecal sample were prepared and read by two investigators¹⁵.

Examination of blood films

Differential blood count:

Fixed blood films were stained by Giemsa method. Blood films were washed and left to dry in air. Dried films were then examined microscopically using 40X and 100X objectives. One hundred white blood cells (WBC) were counted to determine the eosinophil percentages in the peripheral blood of each student. Eosinophil count greater than 6% was considered to be eosinophilia¹⁶.

Statistical analysis

The obtained results were analyzed using SPSS (Version 20). The statistical analysis was performed by using an odds ratio (OR), 95% confidence interval (CI), Chi-square test (χ^2), and probability (*P*) value <0.05 (significant) were used to determine the association between eosinophilia and helminths infection.

Table 1: Characteristics of schoolchildren participate in the study.

Characteristics	Males (n =285)		Females (n =129)		Total	
	No.	%	No.	%	No.	%
Age groups						
5-10 years	45	16	22	17	67	16
11-15 years	240	84	107	82	347	84
Mean age±SD*	12±1.7		12±1.7			
Intestinal helminths						
Infected	52	18	27	21	79	19
Non-infected	233	82	102	79	335	81
Eosinophilia						
Yes	86	30	48	37	134	32
No	199	70	81	63	280	68
Total					414	100

*SD: standard deviation

Table 2: Eosinophilia among students according to their gender, age groups and Intestinal helminthiasis

	Eosinophilia		Normal		OR	C.I	χ^2	P value
	No.	%	No.	%				
Age groups								
5-10	33	49	34	51	2.4	1.39-4.02	10.4	0.002*
11-15	101	29	246	71				
Intestinal helminths								
Infected	47	60	32	40	4.2	2.5-6.98	32.8	<0.001*
Non-infected	87	26	248	74				
Gender								
Males	86	30	199	70	1.4	0.89-2.12	2	0.17
Females	48	37	81	63				

*P value is significant (≤ 0.05)

RESULTS

The present results showed that out of 414 participated schoolchildren; 285(69%) were males and 129(31%) were females. Also, 67(16%) and 347(84%) of student participants were aged, respectively, between-group of 5-15 years and 11-15 years. The frequency rate of infection with intestinal helminths was 79(19%) recorded among both males and females, while 335(81%) of participated were negative for an intestinal infection. The eosinophilia was 134(32%) reported among the schoolchildren (Table 1). The current findings found that the age group 5-10 years had the highest frequent rate of eosinophilia compared to the age group 11-15 years with a statistically significant difference ($\chi^2=10.4$, $P=0.002$). In addition, it was observed that the higher rate of eosinophilia (60%) was recorded among students infected with intestinal parasites whereas 26% of non-infected students had eosinophilia ($\chi^2=32.8$, $P<0.001$). The frequency of eosinophilia was higher among the females (37%) compared to the males (30%) with no statistical difference as listed in Table 2.

Table 3: Types of worm infections among schoolchildren.

Infection type	No.	%
Hymenolepiasis	44	10.6
Ascariasis	17	4.1
Enterobiasis	11	2.7
Taeniasis	10	2.4
Trichuriasis	6	1.4
Schistosomiasis	2	0.5
Mixed infection	10	2.4

The present results revealed that the hymenolepiasis was the most common intestinal worms among schoolchildren (44; 10.6%), followed by ascariasis (17; 4.1%), enterobiasis (11; 2.7%) and taeniasis (10; 2.4%). Also, 10(2.4%) of students were infected with more than one intestinal helminths (Table 3). Table 4 shows that the most frequent of eosinophilia rate was recorded among infected students with *A. lumbricoides* (82%) followed by *E.vermicularis* (73%), *T. trichiura* (67%), *H. nana* (59%), and *T. saginata*(30%). Moreover, the eosinophilia was found in all pupils who were infected with more than one intestinal helminth. There was a statistical significance among students infected with *A. lumbricoides*, *H. nana* ($P<0.001$), and *E. vermicularis* ($P<0.007$). Nevertheless, no significant

difference ($P>0.05$) between infected participants with *T. saginata* and *T. trichiura*.

DISCUSSION

Intestinal parasitic infections are still major public health problems in developing countries and affect the poorest and most deprived communities. Soil-transmitted helminths impair the nutritional status and affect the general intelligence of the people they infect. Malnutrition has a significant impact on the growth and physical development of the infected children^{6,17}. Current study confirms intestinal helminthiasis to remain a medical health problem in children from the two elementary schools at WadiDhahr district with a prevalence rate of 19%. Intestinal helminths in current study are almost similar to those reported from Nepal (20.4%) and Egypt (21%)^{18,19}. The prevalence rate of intestinal helminthiasis among schoolchildren in the present study was lower than that reported from Nigeria (59%)²⁰, India (50%), and Ethiopia (48%)^{20,21,22}. The high prevalence of intestinal helminthiasis may reflect poor adherence to preventive measures which helps re-infection to occur after dewormed programs performed by WHO. Hymenolepiasis was the most frequent intestinal helminthiasis among schoolchildren followed by ascariasis and enterobiasis. Current finding is consistent with studies conducted in Saudi Arabia and Egypt which reported *H. nana* to be the most common intestinal worm among schoolchildren^{23,24}. However, current study disagrees with studies performed in Nigeria which found *A. lumbricoides* to be the most frequent intestinal helminth among Nigerian schoolchildren, and also from a study conducted in Nepal which reported *T. trichiura* to be the most common intestinal worm among schoolchildren^{20,25}. The pattern of intestinal helminthiasis is found to vary geographically within the country (rural versus urban) and from country to country. This variation might be due to changes in the climate and other environmental factors that may support the survival of the infective stage in the soil for a long time^{26,27}. Eosinophilia was common among school children infected with intestinal worms. Current finding is in agreement with those reported by Jiero et al., 2015 and Darlan et al., 2017, who found a significant association between soil-transmitted worms and high blood eosinophil count^{28,29}. This could be explained by fact that worm

infections induce immune responses via T helper cell type 2 subset (Th2 cells). Th2 cells produce interleukin-4 (IL-4), IL-5, IL-10, and IL-13 which stimulate more production of eosinophils from bone marrow resulting in peripheral blood eosinophilia^{17,30}

³³. The presence of blood eosinophilia without intestinal helminthiasis could be attributed to that students may have allergies, other worm infections such as urinary schistosomiasis, or a light number of parasites in the students' gastrointestinal tracts.

Table 4: Association of eosinophilia with different types of intestinal worms

Species of worms	Eosinophilia		Normal		χ^2	C.I	P value
	No.	%	No.	%			
<i>Ascaris lumbricoides</i>	14	82	3	18	20.2	0.026-0.329	<0.001*
<i>Hymenolepis nana</i>	26	59	18	41	16.1	0.150-0.542	<0.001*
<i>Enterobius vermicularis</i>	8	73	3	27	8.4	0.045-0.654	0.007*
<i>Trichuris trichiura</i>	4	67	2	33	3.3	0.042-1.293	0.089
<i>Taeniasaginata</i>	3	30	7	70	0.03	0.285-4.399	0.59

* P value is significant (≤ 0.05)

Eosinophilia was significantly more frequent among children infected with *Ascaris lumbricoides*, *Enterobius vermicularis*, and *H. nana*. Similar observations were reported by other studies^{28, 34-37}. This could be explained by presence of tissue-invading larvae that migrate from the small intestine into blood circulation to reach lungs and elicit pulmonary inflammation^{4, 38-40}. Nevertheless, eosinophilia was non-significant among schoolchildren infected with *T. trichiura* and *Taeniasaginata*. This may be interpreted by worms that do not invade host tissues and thus do not come in contact with host immune system⁴¹. Limitations for this study was neither availability of nearby laboratory to perform complete blood count in order to calculate absolute eosinophil counts nor refrigerator for sample storage until they were being investigated.

CONCLUSION

Current study indicates that eosinophil percentages for schoolchildren who were infected with intestinal helminthiasis were significantly higher than in schoolchildren who were not infected.

ACKNOWLEDGEMENTS

The authors are grateful to all students and teachers who participated in this study.

CONFLICT OF INTEREST

There is no conflict of interest related to this work.

AUTHORS' CONTRIBUTIONS

Authors contributed equally to the design, implementation, statistical analysis and manuscript drafting. All authors read and approved the final manuscript.

REFERENCES

1. Rothenberg M. Eosinophilia. N Engl J Med 1998; 338 (22):1592-1600.
<https://doi.org/10.1056/NEJM199805283382206>
2. Kovalszki A, Weller PF. Eosinophilia. Prim Care 2016; 43(4):607-617.
<https://doi.org/10.1016/j.pop.2016.07.010>
3. Ardiç N. An overview of Strongyloides stercoralis and its infections. Mikrobiyol Bul 2009; 43:169-77.
PMID: 19334396
4. Nutman TB. Evaluation and differential diagnosis of marked, persistent eosinophilia. Immunol Allergy Clin North Am 2007; 27:529-549.

5. Chinchilla RH. Eosinophilia y parasitosis. Rev Med Costa Rica y Centroamerica 2010; 593:241-244.
6. WHO. Soil-transmitted helminth infections; 2020.
7. Pullan RL, Smith JL, Jasrasaria R, Brooker SJ. Global numbers of infection and disease burden of Soil transmitted helminth infections in 2010. Parasites and Vectors 2014; 21(7):37.
<https://doi.org/10.1186/1756-3305-7-37>
8. Hotez PJ, Bundy DP, Beegle K, et al. Helminth infections: soil-transmitted helminth infections and schistosomiasis. Oxford University Press, Washington, 2006.
9. Papier K, Williams GM, Luceres-Catubig R, et al. Childhood malnutrition and parasitic helminth interactions. Clin Infect Dis 2014; 59(2):234-43.
<https://doi.org/10.1093/cid/ciu211>
10. Assoum M. The effects of helminth infections on child physical and cognitive development: an integrated pathophysiological and socioeconomic approach. Adv Life Sci Med 2015; 1(1):1-23.
11. Gelaw A, Anagaw B, Nigussie B, Silesh B, Yirga A, Alem M, Endris M, Gelaw B. Prevalence of intestinal parasitic infections and risk factors among schoolchildren at the University of Gondar Community School, Northwest Ethiopia: a cross-sectional study. BMC Public Health 2013; 3:304.
<https://doi.org/10.1186/1471-2458-13-304>
12. Alshahethi MA, Edrees WH, Mogalli NM, et al. Distribution and risk factors for Giardia lamblia among children at Amran Governorate, Yemen. Universal J Pharm Res 2020; 5(3):34-37.
<https://doi.org/10.22270/ujpr.v5i3.413>
13. Qasem EA, Edrees WH, Al-Shehri WA, Alshahethi MA. Frequency of intestinal parasitic infections among schoolchildren in Ibb city-Yemen. Universal J Pharm Res 2020; 5(2):42-46.
<https://doi.org/10.22270/ujpr.v5i2.388>
14. Al-Mekhlafi AM, Abdul-Ghani R, Al-Eryani SM, Saif-Ali R, Mahdy MA. School-based prevalence of intestinal parasitic infections and associated risk factors in rural communities of Sana'a, Yemen. Acta Trop 2016; 163:135-141.
<https://doi.org/10.1016/j.actatropica.2016.08.009>
15. Hoffman WA, Pons JA, Janer JL. The sedimentation-concentration method in Schistosomiasis mansoni. Puerto Rican J Public Health Trop Med 1934; 2: 283-298.
16. Bain B, Lewis M. Preparation and staining methods for blood and bone marrow films In: Dacie and Lewis (ed) Practical Haematology. 10th edn, Philadelphia, Churchill Livingstone 2006, pp59-77.
17. Jardim-Botelho A, Raff S, Rodrigues Rde A, et al. Hookworm, *Ascaris umbricoides* infection and polyparasitism associated with poor cognitive

- performance in Brazilian schoolchildren. *Trop Med Int Health* 2008; 13(8):994-1004.
<https://doi.org/10.1111/j.1365-3156.2008.02103.x>
18. Kunwar R, Acharya L, Karki S. Decreasing prevalence of intestinal parasitic infections among school-aged children in Nepal: a systematic review and meta-analysis. *Trans R Soc Trop Med Hyg* 2016; 110(6):324-332. <https://doi.org/10.1093/trstmh/trw033>
 19. Farghly AM, Mohamed SM, Abdel-Rahman SA, et al. The relation between the prevalence of soil-transmitted parasites in the soil and among school children in Zagazig district, Sharkya Governorate, Egypt. *J Parasit Dis* 2016; 40(3):1021-1029.
<https://doi.org/10.1007/s12639-014-0627-z>
 20. Karshima SN. Prevalence and distribution of soil-transmitted helminth infections in Nigerian children: a systematic review and meta-analysis. *Infect Dis Poverty* 2018;7(1):69. <https://doi.org/10.1186/s40249-018-0451-2>
 21. Salam N, Azam S. Prevalence and distribution of soil-transmitted helminth infections in India. *BMC Public Health* 2017; 17(1):201.
<https://doi.org/10.1186/s12889-017-4113-2>
 22. Chelkeba L, Mekonnen Z, Alemu Y, Emanu D. Epidemiology of intestinal parasitic infections in preschool and school-aged Ethiopian children: a systematic review and meta-analysis. *BMC Public Health* 2020; 20(1):117.
<https://doi.org/10.1186/s12889-020-8222-y>
 23. Omar MS, Abu-Zeid HA, Mahfouz AA. Intestinal parasitic infections in schoolchildren of Abha (Asir), Saudi Arabia. *Acta Trop* 1991;48(3):195-202.
[https://doi.org/10.1016/0001-706x\(91\)90047-n](https://doi.org/10.1016/0001-706x(91)90047-n)
 24. Ibrahim FA. Prevalence and predisposing factors regarding intestinal parasitic infections among rural primary school pupils at Minia Governorate, Egypt. *J Public Health Afr* 2011; 2(2):e29.
<https://doi.org/10.4081/jphia.2011.e29>
 25. Shrestha A, Schindler C, Odermatt P, et al. Intestinal parasite infections and associated risk factors among schoolchildren in Dolakha and Ramechhap districts, Nepal: a cross-sectional study. *Parasit Vectors* 2018; 11(1):532.
<https://doi.org/10.1186/s13071-018-3105-0>
 26. Silver ZA, Kaliappan SP, Samuel P, et al. Geographical distribution of soil transmitted helminths and the effects of community type in South Asia and South East Asia - A systematic review. *PLoS Negl Trop Dis* 2018;12(1):e0006153.
<https://doi.org/10.1371/journal.pntd.0006153>
 27. Ruberanziza E, Owada K, Clark NJ, et al. Mapping soil-transmitted helminth parasite infection in Rwanda: estimating endemicity and identifying at-risk populations. *Trop Med Infect Dis* 2019; 4(2):93.
<https://doi.org/10.3390/tropicalmed4020093>
 28. Jiero S, Ali M, Pasaribu S, Pasaribu A. Correlation between eosinophil count and soil-transmitted helminth infection in children. *Asian Pacific J Trop Dis* 2015; 5(10):813-816.
[https://doi.org/10.1016/S2222-1808\(15\)60936-7](https://doi.org/10.1016/S2222-1808(15)60936-7)
 29. Darlan DM, Tala ZZ, Amanta C, Warli SM, Arrasyid NK. Correlation between soil-transmitted helminth infection and eosinophil levels among primary school children in Medan. *Open Access Maced J Med Sci* 2017;5(2):142-146.
<https://doi.org/10.3889/oamjms.2017.014>
 30. Moreau E, Chauvin A. Immunity against helminths: interactions with the host and the intercurrent infections. *J Biomed Biotech* 2010; 2010:428593.
<https://doi.org/10.1155/2010/428593>
 31. Shamri R, Xenakis JJ, Spencer LA. Eosinophils in innate immunity: an evolving story. *Cell Tissue Res* 2011; 343(1):57-83. <https://doi.org/10.1007/s00441-010-1049-6>
 32. Alcántara-Neves M, Brito IG, Veiga R, et al. Effects of helminth co-infections on atopy, asthma and cytokine production in children living in a poor urban area in Latin America. *BMC Res Notes*. 2014; 7:817.
<https://doi.org/10.1186/1756-0500-7-817>
 33. Weatherhead JE, Hotez JP. Worm infections in children. *Ped Rev* 2015; 36(8):341-354.
<https://doi.org/10.1542/pir.36-8-341>
 34. Kumar S, Jeong Y, Ashraf MU, Bae YS. Dendritic cell-mediated Th2 immunity and immune disorders. *Int J Mol Sci* 2019; 20:2159-2186.
<https://doi.org/10.3390/ijms20092159>
 35. Sumagaysay JB, Emverda FM. Eosinophilia and incidence of soil-transmitted helminthic infections of secondary students of an indigenous school. *Asian J Health* 2011; 1 (1):172-184.
 36. De-Farias SV, Silva E, Medeiros S, et al. Serum total IgE, Ascaris lumbricoides specific IgE and eosinophils in parasites-infected children in a tropical area. *World Allergy Org J* 2015; 8 Suppl 1:A223.
<https://doi.org/10.1186/1939-4551-8-S1-A223>
 37. Schroeder JC, Jones D, Maranich A. Peripheral eosinophilia found in pediatric Enterobius vermicularis infections. *Clin Pediatr (Phila)* 2019; 58(1):13-16.
<https://doi.org/10.1177/0009922818805193>
 38. Shin MH, Lee YA, Min DY. Eosinophil-mediated tissue inflammatory responses in helminth infection. *Korean J Parasitol* 2009; 47 Suppl: S125-S131.
<https://doi.org/10.3347/kjp.2009.47.S.S125>
 39. Mejia R, Nutman T. Evaluation and differential diagnosis of marked, persistent eosinophilia. *Semin Hematol* 2012; 49(2):149-159.
<https://doi.org/10.1016/j.iac.2007.07.008>
 40. Curtis C, Ogbogu PU. Evaluation and differential diagnosis of persistent marked eosinophilia. *Immunol Allergy Clin North Am*. 2015; 35(3):387-402.
<https://doi.org/10.1016/j.iac.2015.04.001>
 41. Heyneman D. Cestodes In: Baron S. Medical Microbiology. 4th edition, The University of Texas, Galveston, 1996, chapter 89.