Available online on 15.09.2020 at <http://ujpr.org>
Universal Journal of Pharmaceutical Research
 An International Peer Reviewed Journal
 Open access to Pharmaceutical research

This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial Share Alike 4.0 License which permits unrestricted non commercial use, provided the original work is properly cited

Volume 5, Issue 4, 2020

Open Access

RESEARCH ARTICLE

PHYTOCHEMICAL PURIFICATION OF ACTIVE CONSTITUENTS ISOLATED FROM ROOT OF THE MEDICINAL HERB, *CARALLUMA QUADRANGULA*

Jamal AN. Al-Mahweety¹ , Ammer Al-Fadaly² , Waled Abdo Ahmed³

¹School of Chemical Sciences, Faculty of Applied Science, Sana'a University, Yemen.

²School of Chemical Sciences, Faculty of Science, Sana'a University, Yemen.

³School of Chemical Sciences, Faculty of Education, Tamar University, Yemen.

ABSTRACT

Objective: Present study aim for the purification of quantitative phytochemical compounds from roots of *Caralluma quadrangula* belongs to the family Asclepiadaceae. This type of plants can be use as folk medicine to take care of wide diversity of health and diseases situation.

Methods: Preliminary phytochemical analysis for different type of chemical compounds by using various chromatographic techniques. The phytochemical characterizations were evaluated by nuclear magnetic resonance and mass spectrometry.

Results: The quantitative phytochemical analysis of this species exhibited the presence four pure compounds, hydroxyoplopan-4-one (4.5 mg), dihydroxyeudesm-4(15)-ene (5.0 mg), and quercetin- rhamnopyranosyl-D-glucopyranose (Rutin) (7.0 mg).

Conclusion: From this study, it can be concluded that the species found four pure compounds from *C. quadrangula*.

Keywords: *Caralluma quadrangula*, hydroxyoplopan-4-one, dihydroxyeudesm-4(15)-ene, quercetin- rhamnopyranosyl- D-glucopyranose (Rutin).

Article Info: Received 5 July 2020; Revised 1 August; Accepted 29 August, Available online 15 September 2020

Cite this article-

Al-Mahweety JAN, Al-Fadaly A, Ahmed WA. Phytochemical purification of active constituents isolated from root of the medicinal herb, *Caralluma quadrangula*. Universal Journal of Pharmaceutical Research 2020; 5(4):33-36. DOI: <https://doi.org/10.22270/ujpr.v5i4.437>

Address for Correspondence:

Jamal AN. Al- Mahweety, School of Chemical Sciences, Faculty of Applied Science, Sana'a University, Yemen, Tel: +967-771161212, E-mail: jamal.nasser2009@gmail.com

INTRODUCTION

Medically significant genus *Caralluma* is widely studied for its stem and fruits. It is belong to the family Asclepiadaceae, which comprises 200 genera and 2500 species¹. About 200 species belong to genus *Caralluma* distributed throughout Africa and Asia. The greater part of species are native in Indian sub-continent and Arabian Peninsula². A number of *Caralluma* species use as anti-hyperglycemic goings-on of their crude extracts or their corresponding fractions^{3,4}. The investigation of the chemical and biological members of genus *Caralluma*^{3,5} the anti-hyperglycemic activity of the extracts, fractions of the major pregnane glycoside of the aerial parts of *C. quadrangula* was investigated in Kingdom Saudi Arabia as novel. We use the extract of *C. quadrangula* as herbal medicine in Saudi, for the treatment of freckles, diabetes, vitiligo and melasma and for thirst, hunger^{5,6}. Several countries the species of *Caralluma* are fit to be eaten and variety division for the traditional medicine organization⁷. These plants can be use as folk medicine as remedies to health situation and treat large multiplicity of diseases⁸. The species of *C. arabica* use as traditionally for an

emollient and diuretic In United Arab of Emirates. Also used to care for diabetes, hypertension and liver diseases. The *C. Arabica flower* used for wounds and cuts, while the juice of the stem is given to sick people to speed convalescence of burns, itchy skin and sunburns^{9,10}. The *C. attenuate* species in Indian (Andhra Pradesh) use for eaten raw as an anti-diabetic agent, although the juice of the plant beside the black pepper is suggested in the treatment of migraine¹¹. The different applications of *Caralluma* plants in folk medicine have prompted the phytochemical and biological investigations of their constituents¹². The pregnane glycosides, flavone glycosides, megastigmane glycosides, bitter principles, triterpenes and saponins isolated from *Caralluma*^{13,14,15,16}.

MATERIALS AND METHODS

Purified every one of chemical constituent by subsequent standard procedures^{17,18} and all chemicals used systematic Reagent evaluation.

Plant material:-

Roots of *Caralluma quadrangula* (Asclepiadaceae) were collected from Sana'a 2014. The plant identified

by Dr. Hessen Ibrahim and was deposited voucher sampling of plant in Herbarium, Department of Phytochemistry (Sana'a University).

Extraction and Isolation:-

Shade dried roots were crushed and sieved. Next powder was stored in air closing container. Than weighed and extracted with soxhlet extractor by using solvents Chloroform with consecutive solvent extraction. To concentrate the extracts and removal of final traces of solvent than vapor^{19,20}. After that, recrystallization was done to purify the crude extracts. Melting point was taken by using Fisher-John apparatus. The ¹H NMR and ¹³C NMR spectra were taken on Bruker 100 MHz and 400 MHz, spectrometer, using an internal standard like TMS.

Extraction and isolation

Extracted by using Soxhlet (2 Kg) of *C. quadrangula* roots powder with solvents (3X, 8 hours each) and then evaporated collective extracts to give a brown gummy residue (8 g) after than separation and purified by silica gel flash column chromatography (FCC) with CHCl₃ containing increasing percentages of MeOH as eluent and collected 20 ml for each fraction. Fractions 3-10 were combined and rechromatographed by C.C. with CHCl₃-MeOH (8:2) to afford JA1 (4.5 mg) identified as 10 α -hydroxyoplopan-4-one (1), CHCl₃-MeOH (7:3) to afford JA3 (5.0 mg) identified as 1 β , 6 α -dihydroxyeudesm-4(15)-ene (2), CHCl₃-MeOH (6:4) to afford JA4 (5.0 mg) identified as and CHCl₃-MeOH (3:7) to afford JA4 (7.0 mg) identified as quercetin- rhamnopyranosyl- D-glucopyranose (Rutin) (4). NMR data used to identified for each pure compounds.

Figure 1: 10 α -Hydroxyoplopan-4-one (1).

¹H-NMR (100 MHz, CDCl₃) δ : 2.75 (1H, m, H-3), 2.30 (3H, s, H-15), 1.50 (3H, s, H-13), 1.10 (3H, d, H-11), 0.87 (3H, d, H-12); ¹³CNMR (MHz, CDCl₃) δ : 209.4 (C-14), 73.0 (C-8), 56.0 (C-3), 54.6 (C-9), 48.2 (C-5), 45.8 (C-4), 41.0 (C-7), 28.6 (C-10), 27.5 (C-1), 24.4 (C-2), 21.9 (C-6), 21.0 (C-11), 19.2 (C-13), 18.5 (C-15), 15.0 (C-12).

Dihydroxyeudesm-4(15)-ene (2). ¹H-NMR (100 MHz, CDCl₃) δ : 5.10 (1H, brs, H-15), 5.01 (1H, brs, H-15),

3.79 (1H, t, H-6 β), 3.42 (1H, dd, H-1 α), 2.33 (1H, ddd, H-3 α), 2.24 (1H, sept, H-11), 2.07 (1H, ddd, H-3 β), 1.91 (1H, s, H-8), 1.85 (1H, ddd, H-2 α), 1.75 (1H, brd, H-5 α), 1.53 (1H, m, H-2 β), 1.53 (1H, m, H-8), 1.43 (1H, brs, 1-OH), 1.28 (1H, m, H-7 α), 1.20 (1H, m, H-9 α), 1.18 (1H, m, H-9 β), 1.02 (3H, d, H-13), 0.87 (3H, d, H-12), 0.72 (3H, s, H-14); ¹³C-NMR (MHz, CDCl₃) δ : 147.4 (C-4), 108.2 (C-15), 79.1 (C-1), 67.8 (C-6), 56.4 (C-5), 50.1 (C-7), 42.1 (C-10), 36.9 (C-9), 36.1 (C-3), 32.2 (C-2), 26.5 (C-11), 21.8 (C-13), 19.1 (C-8), 16.6 (C-12), 12.0 (C-14).

Figure 2: dihydroxyeudesm-4(15)-ene (2).

Quercetin-L-rhamnopyranosyl-(1 $\rightarrow$ 6)-Dglucopyranose (3). ¹H NMR (100 MHz, CDCl₃) δ : 6.22 (1H, d, H-6), 6.40 (1H, d, H-8), 7.68 (1H, s, H-2'), 6.90 (1H, d, H-5), 7.59 (1H, d, H-6'), 5.10 (1H, d, H_{glc}-1), 3.49 (1H, m, H_{glc}-2), 3.43 (1H, m, H_{glc}-3), 3.50 (1H, m, H_{glc}-4), 3.58 (1H, m, H_{glc}-5), 3.30 (2H, m, H_{glc}-6), 4.51 (1H, br, H_{Rha}-1), 3.10 (1H, m, H_{Rha}-2), 3.43 (1H, m, H_{Rha}-3), 3.54 (1H, m, H_{Rha}-4), 3.31 (1H, m, H_{Rha}-5), 1.17 (3H, d, H_{Rha}-6). ¹³C NMR (CDCl₃) δ : 158.1 (C-2), 135.1 (C-3), 180.0 (C-4), 160.0 (C-5), 100.0 (C-6), 166.1 (C-7), 95.1 (C-8), 163.0 (C-9), 104.7 (C-10), 123.0 (C-1'), 118.2 (C-2'), 146.1 (C-3'), 150.1 (C-4'), 115.8 (C-5'), 123.4 (C-6'), 105.0 (C_{Glc}-1), 75.6 (C_{Glc}-2), 77.8 (C_{Glc}-3), 74.9 (C_{Glc}-4), 77.2 (C_{Glc}-5), 69.0 (C_{Glc}-6), 103.1 (C_{Rha}-1), 71.2 (C_{Rha}-2), 72.1 (C_{Rha}-3), 74.1 (C_{Rha}-4), 70.4 (C_{Rha}-5), 19.2 (C_{Rha}-6).

Figure 3: Quercetin -rhamnopyranosyl-D-glucopyranose.

Figure 4: ¹H NMR of 10 α -hydroxyoplopan-4-one (1).

Figure 5: ^{13}C NMR of 10 α -hydroxyoplopan-4-one (1). Figure 6: ^1H NMR Of dihydroxyeudesm-4(15)-ene (2). Figure 7: ^{13}C NMR of dihydroxyeudesm-4(15)-ene (2).

RESULTS AND DISCUSSION

Compound 1: The ^1H NMR showed one multiplet proton at δH 2.75 (1H, m), 5.46, High intensity Peaks at δ 2.30, 1.50, 1.10 and 0.88 are corresponding to methyl groups (Me- (15, 13, 14 and 12). 4 methyl, 4 methylene, 5 methine and 2 quaternary carbons presence in ^{13}C NMR spectrum. Carboxylic group signals become visible at δ 209.5. In addition of β -hydroxyl group to C8 is visible from a peak at δ 73.1. hydroxyoplopan-4-one, it has never been isolated before from *Caralluma quadrangula*, it reported from *Cassia buds*²¹.

Compound 2: The ^1H NMR showed signals for three angular methyl singlet's at δH 0.95, 0.85 and 0.71. proton of H-6 and H-1 appeared at δ 3.79 and 3.42. Olefinic protons present at δ 5.10 and 4.95 for H-15.

^{13}C NMR showed fifty carbon signal including three CH_3 , five CH_2 , five CH and two quaternary carbons. The double bond carbons appeared at δ 147.4 and 108.1. The significant signal for the 1β , 6α -dihydroxyeudesm-4(15)-ene would be the signals for two carbon attached to hydroxyl group, which is C-1 and C-6 that appeared at δ 79.2 and 67.8^{22,23}.

Compound 3: The ^1H NMR spectrum exhibited signals which were typical of a flavone compound. In addition to the presence of five aromatic protons; one was represented by two *meta*-coupled protons at δH 6.23 (*d*, H-6) and 6.42 (*d*, H-8). ^{13}C NMR experiments showed one methyl, 15 methines, one methylene and 10 quaternary carbon atoms, one being the flavone carbonyl (C 180.0)²⁴. NMR spectral data confirmed the sugar part assigned as glucose and rhamnose.

Figure 8: ^1H NMR of quercetin -rhamnopyranosyl-D-glucopyranose (3). Figure 9: ^{13}C NMR of quercetin -rhamnopyranosyl-D-glucopyranose (3).

A significant downfield shift of the methylene carbon appearing at C 69.1 and assigned to C-6 of glucose, indicated a (1 to 6) type of interglycosidic linkage to the rhamnose moiety. Quercetin-rhamnopyranosyl-D-glucopyranose isolated from *C. quadrangula* for first time, it was reported in many plants as *Taverniera aegyptiaca*^{25,26}.

CONCLUSION

The isolation and identification 10 α -hydroxyoplopan-4-one, dihydroxyeudesm-4(15)-ene, and quercetin-rhamnopyranosyl-D-glucopyranose (Rutin), from the roots of *C. quadrangula*. The work was carried out by means of various physical (solvent extraction, column chromatography, radial chromatography, preparative thin layer chromatography and melting points) and spectral techniques.

AUTHOR'S CONTRIBUTION

The manuscript was carried out, written, and approved in collaboration with all authors.

ACKNOWLEDGEMENTS

The authors extend their thanks and appreciation to the Sana'a University, Yemen to provide necessary facilities for this work.

CONFLICT OF INTEREST

No conflict of interest associated with this work.

REFERENCES

- Rajendra R, Kamala. USP filed 2004; 4: 6376657.
- Mohammad D, Al-Khateeb M, Riyadh E, Al-Hashem F, Mohammad KN. *In vivo*, acute, normo-hypoglycemic, antihyperglycemic, insulinotropic actions of orally administered ethanol extract of *Citrullus colocynthis* (L.) Schrab pulp. American J Biochem Biotech 2009; 5(3):119-26.
<https://doi.org/10.3844/ajbbsp.2009.118.125>
- Prabhakar KP, Doble M. A target based therapeutic approach towards diabetes mellitus using medicinal plants. Curr Diabetes Rev 2008; 4: 291–308.
<https://doi.org/10.2174/157339908786241124>
- Sattar EA, Harraz FMH, Ghareib ESA, et al. Antihyperglycemic and hypolipidaemic effects of the methanolic extract of *Caralluma tuberculata* in streptozotocin-induced diabetic rats. Nat Prod Res 2011; 25: 1171–1179.
<https://doi.org/10.1080/14786419.2010.490782>
- Sattar EA, Abdel-Naim AB, Khedr A, Shehata IA. Anti-hyperglycemic activity of *Caralluma tuberculata* in streptozotocin-induced diabetic rats. Food Chem Toxicol 2013; 59: 111–117.
<https://doi.org/10.1016/j.fct.2013.05.060>
- Habibuddin M, Daghriri HA, Humaira T, et al. Anti-diabetic effect of alcoholic extract of *Caralluma sinaica* L. on streptozotocin-induced diabetic rabbits. J Ethnopharmacol 2008; 117: 215–220.
<https://doi.org/10.1016/j.jep.2008.01.021>
- Latha S, Rajaram K, Suresh KP. Hepatoprotective and antidiabetic effect of methanol extract of *Caralluma fimbriata* in streptozotocin induced diabetic albino rats. Int J Pharm Pharm Sci 2014; 6: 665–668.
- Gushash AS. Plants in the Mountains of Sarat and Hejaz. Sarawat, Al Madinah press, KSA, 2006.
- Abdel-Sattar E, Ahmed AA, Hegazy ME, Farag MA, Al-Yahya MA. Acylated pregnane glycosides from *Caralluma russeliana*. Phytochem 2007; 68: 1459–1463. <https://doi.org/10.1016/j.phytochem.2007.03.009>
- Oyama M, Iliya I, Tanaka T, Linuma M. Five new steroidal glycosides from *Caralluma dalzielii*. Helvetica Chim Acta 2007; 90, 63–71.
<https://doi.org/10.1002/hlca.200790022>
- Ahmad MM, Qureshi S, Shah A, et al. Anti-Inflammatory activity of *Caralluma tuberculata* alcoholic extract. Fitoterapia 1983; 46: 357–360.
- Zakaria MN, Islam MW, Radhakrishnan R, et al. Antinociceptive and anti-inflammatory properties of *Caralluma arabica*. J Ethnopharmacol 2001; 76: 155–158. [https://doi.org/10.1016/S0378-8741\(01\)00208-2](https://doi.org/10.1016/S0378-8741(01)00208-2)
- Ramesh YN, Rao AV, Rao MC, et al. Antinociceptive and anti-inflammatory activity of a flavonoid isolated from *Caralluma attenuata*. J Ethnopharmacol 1998; 62: 63–66. [https://doi.org/10.1016/S0378-8741\(98\)00048-8](https://doi.org/10.1016/S0378-8741(98)00048-8)
- Dembitsky VM. Chemistry and biodiversity of the biologically active natural glycosides. Chem Biodiv 2004; 1: 673–781.
<https://doi.org/10.1002/cbdv.200490060>
- Kumar DS. A medicinal plants survey for treatment of obesity. J Pharm Res 2011; 4: 597–600.
- Bader A, Braca A, Tommasi ND, Morelli I. Further constituents from *Caralluma negevensis*. Phytochem 2003; 62: 1277–1281.
[https://doi.org/10.1016/S0031-9422\(02\)00678-7](https://doi.org/10.1016/S0031-9422(02)00678-7)
- Braca A, Bader A, Morelli I, et al. Tommasi. New pregnane glycosides from *Caralluma negevensis*. Tetrahedron 2002; 58: 5837–5848.
- Malladi S, Ratnakaram VN, Babu SK, Pullaiah T. Phytochemical investigation of *C. lasiantha*: isolation of stigmasterol, an active immunomodulatory agent. Int J Chem Sci 2017; 15: 339–407.
- Trease GE, Evans WC. Pharmacognosy, Saunders. Elsevier, Amsterdam, The Netherlands 2002; 36: 51.
- Gupta AK. Introduction to Pharmaceutics-1. CBS publication, New Delhi 2004.
- Yindengzhi G, Haofeng Z, Junfei Z, Haiqing Z. Isolation and characterization of Sesquiterpenoids from Cassia buds and their antimicrobial activities. Agri Food Chem 2017; 65(28): 5614–5619.
<https://doi.org/10.1021/acs.jafc.7b01294>
- Al-Mahweety J, Azzam S, Ali A. A study of phytochemical constituents in *Caralluma quadrangula*. Universal J Pharm Res 2020; 5(2):28–31.
<https://doi.org/10.22270/ujpr.v5i2.385>
- Lene H, Boll PM. The polyacetylenic falcarinol as the major allergen in *Schefflera arboricola*. Phytochem 1986; 5 (2): 529–530.
[https://doi.org/10.1016/S0031-9422\(00\)85517-X](https://doi.org/10.1016/S0031-9422(00)85517-X)
- Helena G, Palma BFMS, Torre MCDL. Sterols from *Teucrium abutiloides* and *T. betonicum*. Phytochem 1996; 43(3): 613–61.
[https://doi.org/10.1016/0031-9422\(96\)00337-8](https://doi.org/10.1016/0031-9422(96)00337-8)
- Hasan A, Hhadiga F, Sayed A, John N, Soren BC. A new flavonol glycoside and other flavonoids from the aerial parts of *Taverniera aegyptiaca*. Nat Prod Res 2019; 33(8):1135–1139.
<https://doi.org/10.1080/14786419.2018.1460834>
- Amal M, Youssef M, Ahmed I, Mahmoud A. Isolation, structural elucidation of flavonoid constituents from *Leptadenia pyrotechnica* and evaluation of their toxicity and antitumor activity. Nat Prod Res 2009; 23(3):539–552. <https://doi.org/10.1080/13880200902875065>