Available online on 15.09.2020 at <http://ujpr.org>
Universal Journal of Pharmaceutical Research
 An International Peer Reviewed Journal

Open access to Pharmaceutical research
 This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial Share Alike 4.0 License which permits unrestricted non commercial use, provided the original work is properly cited

Volume 5, Issue 4, 2020

Open Access

REVIEW ARTICLE

EXPLORING THE ANTIPARASITIC ACTIVITY OF MEDICINAL PLANTS

Gehan Fawzy Abdel Raouf^{1*} , Ahmed Wael Lethy Mohamed²

¹Pharmacognosy Department, Pharmaceutical and Drug Research Industries Division, National Research Centre, Dokki, Giza 12622, Egypt.

²College of Biotechnology, Misr University for Science and Technology (MUST), 6th of October City, Egypt.

ABSTRACT

Parasitic disease is one of the most important challenges impacting a large number of people. Worldwide, more than three billion cases of parasitic disease are reported yearly. Due to the resistance of parasite to the synthetic drugs, it is necessary to search for alternative sources. Plants contain different phytoconstituents with different biological activities. Many previous researches showed that many plants exerted antiparasitic activity due to its secondary metabolites. This review highlights the antiparasitic effects of different medicinal plants and mentions the mode of action of different phytochemicals against parasites. In conclusion, the medicinal plants play a vital role as antiparasitic agents, but further studies are needed to isolate and test the active constituents of the medicinal plants as antiparasitic agents targeting to enter the drug discovery area.

Keywords: Antiparasitic, medicinal plants, parasitic disease, phytoconstituents.

Article Info: Received 20 June 2020; Revised 24 July; Accepted 26 August, Available online 15 September 2020

Cite this article-

Raouf GFA, Mohamed AWL. Exploring the antiparasitic activity of medicinal plants. Universal Journal of Pharmaceutical Research 2020; 5(4):71-75.

DOI: <https://doi.org/10.22270/ujpr.v5i4.442>

Address for Correspondence:

Dr. Gehan F. Abdel Raouf, Pharmacognosy Department, National Research Centre, Dokki, 12622, Giza, Egypt, Tel: +201223417848, E-mail: gehankandeel9@yahoo.com

INTRODUCTION

Parasitic diseases are the most important public health problem impacting a large number of people worldwide. Parasites are considered to be a major problem in our life. A large variety of parasites have developed during the evolution of humans. Typically, a parasite will not destroy its host instantly. Most internal parasites are weakening our health, while think of lice and fleas are unpleasant for us. If the patients do not get appropriate therapy, some parasitic infections can be deadly, such as Chagas, trypanosomiasis or malaria¹. The transmission of parasites is facilitated because of the bad hygienic conditions. People have often tried to mitigate the parasitic infection. Mechanically, it may minimize or eliminate external parasites. While, internal parasites are more complicated to treat². A number of drugs have been synthesized from the medicinal chemists which can be used as antiparasitic drugs, but some parasites showed resistance to these drugs. So, searching for antiparasitic drugs from natural origin is necessary to be an alternative to synthetic drugs³. For several thousands of years, humans have used medicinal plants to treat diseases and health problems. For centuries, medicinal plants have been used as antiparasitic agents, and up till now, are still used for this purpose⁴. Many medicinal plants showed antiparasitic activity against

different parasites. These medicinal plants contain different biologically active compounds that showed antiparasitic activity. For example, saponins impact on the cell membrane permeability of the parasites causing vacuolization of teguments⁵. The aim of this review is to highlight the antiparasitic effects of different medicinal plants and to know the mode of action of different phytochemicals against parasites.

PARASITES

A parasite lives in or on a host; it depends on the resources of its host to maintain its life cycle. Most of parasites are invisible by the naked eye, but others can reach a length of 30 meters or more as some worm parasites. Parasites can cause spreading of different diseases⁶. Parasites cause large numbers of infections and lead to several million deaths every year⁷. There are different ways for parasitic infections as polluted vegetables, food, soil and water leading to different complications as allergies, anemia, malnutrition and gastrointestinal disorders. The parasitic infections cause many tropical diseases, such as helminthiasis, leishmaniasis, onchocerciasis, malaria, lymphatic filariasis, Chagas disease, trypanosomiasis and schistosomiasis². Helminth (parasitic worm) can exist as individuals or as parasites dependent on plant or animal hosts. In human beings helminthic infections

are known as one of the most common infections⁸. The lives of billions of people worldwide are affected by protozoan parasites that cause large economic impacts⁹.

Table 1: Examples of some antiparasitic drugs.

Category of the drug	Examples
Anticestoda	Praziquantel, Levamisole, Niclosamide
Antirematoda	metrifonate, oxamniquine, praziquantel, bithionolsulfoxide
Antinematoda	diethyl carbamazepine, ivermectin, piperazine, pyrantel, mebendazole, thiabendazole

Types of parasites

There are two main types of parasites endoparasites and ectoparasites. Human endoparasites live inside their hosts, in the alimentary canal or within cells or tissues. There are two types of endoparasites; Protozoa which include the unicellular organism called Plasmodium. The other type are helminthes (worm parasites) such as; tapeworm, fluke, pinworm, roundworm and trichina spiralis. On the other hand, ectoparasites live on, rather than in their hosts. They include fleas and lice¹⁰.

Prevention

To avoid and prevent parasitic infections, there are several precaution should be followed such as; washing the hands regularly, drink clean water and avoid swallowing water from ponds streams or lakes, cook

food to its recommended internal temperature, avoid cat feces and litter especially for pregnant woman and safe sex practicing¹¹.

Diagnosis

There are different ways for diagnosis of the parasitic infections such as; a blood test, a fecal exam to check the presence of parasites or their eggs in the stool sample, colonoscopy or endoscopy in which the doctor will pass a thin tube into the digestive system through the mouth or rectum of patient to examine his intestinal tract, the other ways for the diagnosis of parasitic infections by some scans to inspect the presence of any lesions or damages of organs by parasites, these scans such as; magnetic resonance imaging (MRI), computerized axial tomography (CAT), or X-ray¹¹.

Treatment

A number of drugs have been synthesized from the medicinal chemists that can be used as antiparasitic drugs. Some parasites showed resistance to these drugs. So, searching for antiparasitic drugs from natural origin is necessary to be used instead of synthetic drugs³. For the pharmaceutical industry, the production of new synthetic antiparasitic drugs is a risky affair due to a high price of the drugs and because a lot of parasitic infestations exist in developing countries where it is difficult for the people to pay. So, new drugs derived from natural products or their derivatives are necessary to be an alternative to synthetic drugs². Natural products play a vital role in medicine; large numbers of new drugs were derived from natural products or their derivatives⁴. Table 1 showed examples of some anthelmintic drugs.

Table 2: Antiparasitic medicinal plants and their secondary metabolites.

No.	Scientific name	Family	Chemical constituent
1	<i>Cinchona officinalis</i> ¹⁴	Rubiaceae	Quinine, quinidine, cinchonine, and cinchonidine
2	<i>Artemisia annua</i> ²²	Asteraceae	sesquiterpene
3	<i>Streblus asper</i> ²	Moraceae	Asperoside, strebluside
4	<i>Carica papaya</i> ²¹	Caricaceae	Papain, Benzyl isothiocyanate
5	<i>Cichorium intybus</i> ³	Asteraceae	sesquiterpene lactones
6	<i>Buteamonosperma</i> ¹⁶	Fabaceae	Palasonin and tannins
7	<i>Zingiber officinale</i> ¹⁶	zingiberaceae	Zingiberene, gingerols, shogaols and bisabolene
8	<i>Dryopteris filix-mas</i> ²	Dryopteridaceae	Vermicidal phloroglucinols
9	<i>Punicagranatum</i> ¹³	Lythraceae	Alkaloid, tannins, glycosides
10	<i>Artemisia herba-alba</i> ¹³	Asteraceae	Alkaloid, tannins and phenol
11	<i>Ailanthus altissima</i> ¹⁷	Simaroubaceae	Quassinoids, ailanthone
12	<i>Allium sativum</i> ^{18,20}	Amaryllidaceae	Allicin and ajoene
13	<i>Dillenia suffruticosa</i> ¹⁵	Dilleniaceae	MethylGlycolate, phenol, tridecanal
14	<i>Dichrostachys cinerea</i> ⁵	Fabaceae	Cardiac glycosides, flavonoids, tannins, triterpenoids and saponins

Antiparasitic medicinal plants and their secondary metabolites

Herbal medicine plays a vital role in the management of many diseases as it has become a very safe, non-toxic, and easily available source. Owing to the existence of diverse bioactive compounds with antioxidant properties, the extracts of medicinal plants can be used as a natural treatment for the infestation of

parasites. Some parasites showed resistance for many of synthetic drugs. So all the world is going to produce new drugs from natural origins and plants¹³.

Cinchona officinalis

Cinchona officinalis belongs to family Rubiaceae. From *Cinchona officinalis* and related *Cinchona* species, the first medications to cure malaria were developed. Quinoline alkaloids are the main active

constituents of Cinchona bark. Cinchonidine (Quinimax) was the bitter-tasting quinine that is used to cure the *Plasmodium* phases in the blood. The basic structure for synthesizing multiple antimalarial drugs is quinine. Quinine as well as its conjunction with clindamycin, doxocycline or tetracycline are important in the treatment of severe *P. falciparum* infections¹⁴.

***Dichrostachys cinerea* (Family Fabaceae)**

Dichloromethane extract of *Dichrostachys cinerea* stem bark exhibited a potent *in vitro* antiplasmodial effect against the chloroquine resistant strain of *Plasmodium falciparum*. In addition, it showed a potent suppression of parasite and capacity to recover disease in the animal model of *P. berghei*. Moreover, *Dichrostachys cinerea* methanol extract exhibited cytotoxicity with concentration 178.35µg/ml, indicating that *Dichrostachys cinerea* methanol extract may be defined as anti-malarial drug⁵.

Cichoriumintybus

Cichoriumintybus (Chicory) belongs to family Asteraceae. It showed antiparasitic activity as it

contains different bioactive compounds. Sesquiterpene lactones rich extracts from chicory exhibited a potent activity against different gastrointestinal helminths of livestock, as well as it showed antimalarial properties³.

***Dillenia suffruticosa* (Dilleniaceae)**

Different bioactive compounds are presented in the leaves of *D. suffruticosa*. The methanol extract of the leaves showed a potent effect against the deleterious leeches of hybrid groupers. It was found that *D. suffruticosa* methanol extract revealed strong antiparasitic activity against the marine leech *Zeylanicobdella arugamensis* with 100% mortality¹⁵.

Butea monosperma

Butea monosperma belongs to family Fabaceae. The methanol extract of *B. monosperma* showed anthelmintic activity *in vitro*. The different species of *Butea* have been reported to exhibit anthelmintic activity against *Dipylidium caninum*, *Taenia*, earth worm, *A. galli* and *Ascaris lumbricoides*. *Butea monosperma* contains many bioactive compounds as tannins and palasonin¹⁶.

Table 3: The mode of action of different phytochemicals.

Phytochemical	Action
Phenolic compounds ^{5,23}	Uncoupling the oxidative phosphorylation leads to disturbance in energy generation mechanism
Alkaloids ^{13, 24}	- At as an antioxidant - Steroidal alkaloid and oligoglycosides inhibit the transfer of sucrose from the stomach to the small intestine. - Cause paralysis due to its effect on central nervous system.
Tannins ^{5,25}	-Uncoupling the oxidative phosphorylation leads to disturbance in energy generation mechanism -Binding glycoprotein on the cuticles of the worms or the free protein of the GI tract of the host animal
Cysteine proteinases ²⁶	Cause digestion of nematode cuticle
Isoflavones ²⁷	Disturb the Ca ²⁺ homeostasis in the parasites. Inhibit the enzymes of glycogenolysis and glycolysis

***Zingiber officinale* Rosc (Ginger)**

Zingiber officinale belongs to family zingiberaceae. Zingiberene, gingerols, shogaols and bisabolene are the main active compounds of *Zingiber officinale*. A monocyclic sesquiterpene (zingiberene) is the main component of ginger oil. It gives ginger its distinct flavoring and represents about 30% of the essential oils in ginger rhizomes. The alcoholic extract of rhizomes of *Z. officinale* showed the anthelmintic activity against human *A. lumbricoides*. Also it exhibited antischistosomal activity¹⁶.

Dryopterisfilix-mas

Dryopterisfilix-mas (Dryopteridaceae) contains biologically active compounds that effective against cestodes and causing worm muscles paralysis, these compounds (vermicidal phloroglucinols) such as deaspidin, filixic acid and aspidin².

Punicagranatum

Punicagranatum (Lythraceae) is effective against gastrointestinal nematodes. The crude extracts of *P. granatum* showed a potential anthelmintic activity. By

comparing with the negative control, the extract showed a potent nematocidal effect at the highest concentration (10 mg/ml). Moreover, plant extract showed a potent inhibitory effect against hatching of the egg within 48hr of exposure¹³.

Artemisia herba-alba

Artemisia herba-alba (Asteraceae family) is used as an anthelmintic agent. The crude extract of *A. herba-alba* showed a potential anthelmintic activity at all dose levels. *Artemisia herba-alba* flower extract showed a potent inhibitory effect against egg hatching (98.67%) at concentration 1 mg/ml comparing with the negative control, this result confirmed the nematocidal activity of *Artemisia herba-alba*¹³.

Ailanthus altissima

Ailanthus altissima belongs to Simaroubaceae family. From the active extracts of *Ailanthus altissima*, 6 alpha tigloyloxy chaparrinone and ailanthone were isolated and showed a potent inhibitory effect against *Plasmodium falciparum* strains *in vitro*. Extracts of *Ailanthus altissima* (Mill.) Swingle have been tested

for activity. The chloroformic extract showed a potent effect on *P. berghei* in mice and on *P. falciparum* *in vitro*. In addition, the presence of the quassinoid anthrone in the plant plays an important role in this activity¹⁷.

Allium cepa (onion) and Allium sativum (garlic)

The effects of onion (*Allium cepa*) and garlic (*Allium sativum*) on adult parasite *Lernantropus kroyeri* (*L. kroyeri*) were studied. Results showed that onion and garlic juices exhibited the inhibitory effect on the females of *Lernantropus kroyeri* in a concentration and time dependent manner¹⁸⁻²⁰.

Carica papaya

Carica papaya belongs to family Caricaceae. Benzylisothiocyanate and Papain are the main active constituents of Papaya. Papain, papaya proteinase I, is a cysteine protease enzyme present in seed, fruit and leaves of papaya. The latex containing papain showed anthelmintic properties against intestinal nematodes of poultry²¹. Papain is comprised of a polypeptide chain with three disulfide bridges and a sulfhydryl group required for the enzyme activity which is responsible for digestion of nematodes cuticle¹⁶.

Streblus asper

Streblus asper family Moraceae exhibited a potent antifilarial activity. The main active components of *Streblus asper* are the cardiac glycosides strebluside and asperoside².

Artemisia annua (Asteraceae)

Artemisia annua contains the sesquiterpene artemisinin which plays a vital role as an antimalarial agent. It showed a potent activity against *P. falciparum*. Various semi-synthetic derived products from artemisinin have been developed, and today in clinical practice²². Table 2 summarized the most important antiparasitic medicinal plants and their secondary metabolites. While Table 3 showed the mode of action of different phytochemicals.

CONCLUSION

This review highlights the antiparasitic effects of different medicinal plants as well as the mode of action of different phytochemicals against parasites. So, the current review provides recommendations based on proof to understand the role of medicinal plants as antiparasitic agents aiming to be the first step towards the production of novel drugs for controlling the parasitic infection. Furthermore, more research is needed to run clinical trial to confirm the effect of medicinal plants as antiparasitic agents. In future, medicinal plants should be the first choice for the management of parasitic infection.

ACKNOWLEDGEMENTS

The authors extend their thanks and appreciation to the National Research Centre, Giza, Egypt to provide necessary facilities for this work.

CONFLICTS OF INTEREST

No conflict of interest associated with this work.

REFERENCES

1. Panda SK, Luyten W. Antiparasitic activity in Asteraceae with special attention to ethnobotanical use by the tribes of Odisha, India. *Parasite* 2018; 25(10):1-25. <https://doi.org/10.1051/parasite/2018008>
2. Wink M. Medicinal Plants: A source of anti-parasitic secondary metabolites. *Molecules* 2012; 17(11):12771-12791. <https://doi.org/10.3390/molecules171112771>
3. Peña-Espinoza M, Valente AH, Thamsborg SM, Simonsen HT, Boas U, Enemark HL, Muñoz RL, Williams A R. Antiparasitic activity of chicory (*Cichorium intybus*) and its natural bioactive compounds in livestock: a review. *Parasit Vect* 2018; 11(475):1-14. <https://doi.org/10.1186/s13071-018-3012-4>
4. Newman D J, Cragg GM. Natural Products as Sources of New Drugs over the Last 25 Years. *J Nat Prod* 2007; 70(3):461-477. <https://doi.org/10.1021/np068054v>
5. Kweyamba PA, Zofou D, Efang N, Assob JC N, Kitau J, Nyindo M. *In vitro* and *in vivo* studies on anti-malarial activity of *Commiphora africana* and *Dichrostachys cinerea* used by the Maasai in Arusha region, Tanzania. *Malaria J* 2019; 18(119):1-6. <https://doi.org/10.1186/s12936-019-2752-8>
6. Bundy DAP, Appleby LJ, Brooker SJ. Nematodes Limited to the Intestinal Tract (*Enterobius vermicularis*, *Trichuris trichiura*, *Capillaria philippinensis* and *Trichostrongylus* spp.). *Hunter Tropical Med Emerg Infect Diseases* 2020; 10:834-839. <https://doi.org/10.1016/B978-1-4160-4390-4.00107-7>
7. Peters W, Pasvol G. Atlas of Tropical Medicine and Parasitology. 6th ed.; Mosby-Elsevier: Philadelphia, PA, USA. 2007; 1-376.
8. Raman VK, Saini M, Sharma A, Parashar B. *Morchella esculenta*: a herbal boon to pharmacology. *Int J Dev Res* 2018; 8(3):19660-19665.
9. Murray CJL, Rosenfeld LC, Lim SS, et al. Global malaria mortality between 1980 and 2010: A systematic analysis. *Lancet* 2012; 379:413-431. [https://doi.org/10.1016/S0140-6736\(12\)60034-8](https://doi.org/10.1016/S0140-6736(12)60034-8)
10. Jacobsen KH, Ribeiro PS, Quist BK, Rydbeck BV. Prevalence of intestinal parasites in young Quichua children in the highlands of rural Ecuador. *J Health Popul Nut* 2007; 25:399-405. [PMID: 18402182](https://pubmed.ncbi.nlm.nih.gov/18402182/)
11. Bahk YY, Shin EH, Cho SH, Ju JW, Chai JY, Kim TS. Prevention and control strategies for parasitic infections in the Korea centers for disease control and prevention. *The Korean J Parasitol* 2018; 56(5):401-408. <https://doi.org/10.3347/kjp.2018.56.5.401>
12. Bahmani M, Kopaei M R, Hassanzadazar H, Saki K, Karamati S A, Delfan B. A review on most important herbal and synthetic antihelmintic drugs. *Asian Pac J Trop Med* 2014; 7:29-33. [https://doi.org/10.1016/S1995-7645\(14\)60200-5](https://doi.org/10.1016/S1995-7645(14)60200-5)
13. Ahmed AH, Ejo M, Feyera T, Regassa D, Mammed B, Huluka SA. *In vitro* anthelmintic activity of crude extracts of *Artemisia herba-alba* and *Punicagranatum* against *Haemonchus contortus*. *J Parasitol Res* 2020; 1:1-7. <https://doi.org/10.1155/2020/4950196>
14. Willcox M. Improved traditional phytomedicines in current use for the clinical treatment of malaria. *Planta Med* 2011; 77:662-671. <https://doi.org/10.1055/s-0030-1250548>
15. Shah MD, Maran BAV, Iqbal M, Ching FF, Lal MTM, Othman R B, Shapawi R. Antiparasitic activity of the medicinal plant *Dillenia suffruticosa* against the marine leech *Zeylanicobdella arugamensis* (Hirudinea) and its phytochemical composition. *Aquacul Res* 2019; 51:215-221. <https://doi.org/10.1111/are.14367>
16. Bauri RK, Tigga M N, Kullu SS. A review on use of medicinal plants to control parasites. *Indian J Nat Prod Res* 2015; 6(4):268-277.
17. Al-Snafi AE. The pharmacological importance of *Ailanthus altissima*- a review. *Int J Pharmacy Rev Res* 2015; 5(2):121-129.

18. Krstin S, Sobeh M, Braun M S, Wink M. Anti-parasitic activities of *Allium sativum* and *Allium cepa* against *Trypanosoma brucei* and *Leishmania tarentolae*. *Medicines* 2018; 5(37):1-13. <https://doi.org/10.3390/medicines5020037>
19. Upadhyay RK. Nutraceutical, pharmaceutical and therapeutic uses of *Allium cepa*: A review. *Int J Green Pharm* 2016; 10(1):46-64. <http://dx.doi.org/10.22377/ijgp.v10i1.612>
20. Yildiz HY, Van QP, Parisi G, Sao MD. Anti-parasitic activity of garlic (*Allium sativum*) and onion (*Allium cepa*) juice against crustacean parasite, *Lernantropus kroyeri*, found on European sea bass (*Dicentrarchus labrax*). *Italian J Animal Sci* 2019; 18(1):833-837. <https://doi.org/10.1080/1828051X.2019.1593058>
21. Adu OA, Akingboye KA, Akinfemi A. Potency of Papaya (*Carica papaya*) latex as an anthelmintic in poultry production. *Botany Res Int* 2009; 2(3):139-142.
22. Efferth T, Herrmann F, Tahrani A, Wink M. Cytotoxic activity towards cancer cells of secondary constituents derived from *Artemisia annua* L. in comparison to its designated active constituent artemisinin. *Phytomed* 2011; 18:959-969. <https://doi.org/10.1016/j.phymed.2011.06.008>
23. John J, Mehta A, Shukla S, Mehta P. A report on anthelmintic activity of *Cassia tora* leaves. *J Sci Tech* 2009; 31(3):269-271.
24. Roy H, Chakraborty A, Bhanja S, Nayak BS, Mishra RS, Ellaiah P. Preliminary phytochemical investigation and anthelmintic activity of *Acanthospermum hispidum* DC. *J Pharm Sci Tech* 2010; 2(5):217-221.
25. Patel J, Kumar G S, Qureshi MS, Jena PK. Anthelmintic activity of ethanolic extract of whole plant of *Eupatorium odoratum*. *Int J Phytomed* 2010; 2:127-132.
26. Caffrey CR, Goupil L, Rebello KM, Dalton JP, Smith D. Cysteine proteases as digestive enzymes in parasitic helminthes. *PLOS Negl Trop Dis* 2018; 12(8):1-20. <https://doi.org/10.1371/journal.pntd.0005840>
27. Stepek G, Lowe AE, Buttle DJ, Duce IR, Behnke JM. *In vitro* and *in vivo* anthelmintic efficacy of plant cysteine proteinases against the rodent gastrointestinal nematode, *Trichuris muris*. *Parasitol* 2006; 132:681-689. <https://doi.org/10.1017/S003118200500973X>