



Available online on 15.05.2017 at <http://ujpr.org>

Universal Journal of Pharmaceutical Research

An International Peer Reviewed Journal

Open access to Pharmaceutical research

©2017, publisher and licensee UJPR, This is an open access article which permits unrestricted non commercial use, provided the original work is properly cited

Volume 2, Issue 2, 2017

RESEARCH ARTICLE

ANTI-HYPERGLYCEMIC AND ANTI-OXIDANT ACTIVITIES OF METHANOL EXTRACT OF *GONGRONEMA LATIFOLIUM*

Edenta Chidi¹, Nwobodo Ndubuisi Nwobodo², Nweje-Anyalowu Paul C³

¹Department of biochemistry, Renaissance University, Enugu, Nigeria.

²Department of Pharmacology and Therapeutics, Division of Clinical Pharmacology and Translational Medicine, University of Science and Technology, Enugu State, Nigeria.

³Clifford University, Owerri, Abia State, Nigeria

*Corresponding Author's Email: edentachidi1@gmail.com

DOI: <http://doi.org/10.22270/ujpr.v2i2.R4>

Received 1 April 2017; Revised 12 May; Accepted 13 May, Available online 15 May 2017

ABSTRACT

Gongronema latifolium (Asclepiadiaceae) is used as bitter spice or flavouring agent in many traditional Nigerian dishes. It is well known for medicinal and nutritional purposes like anti-hyperglycemic, anti-oxidant, antitussive and widely distributed in the southeastern states of Nigeria. Objective of present study was to evaluate the anti-hyperglycemic and anti-oxidant effect of the methanol extract of *Gongronema latifolium* leaves. The methanol extract of *G. latifolium* at doses of 250 and 500 mg/kg were studied for anti-hyperglycemic effect in alloxan-induced hyperglycemic rats. The variation in blood glucose level in normal and experimental rats on 0, 7, 14 and 21 days of treatment has also been recorded. Treatment with *G. latifolium* extract showed signs of recovery as comparable with the standard drug glibenclamide (0.25 mg/kg). The effect of different treatment on body weight of rats was also determined. The antioxidant properties of the extracts as determined by the DPPH free radical scavenging assay. The result showed that *G. latifolium* extract caused a concentration dependent percentage increase of antioxidant activity. Study concludes that methanol leaf extract of *G. latifolium* possesses significant anti-hyperglycemic and anti-oxidant activities.

Keywords: Anti-hyperglycemic, anti-oxidant, carrageenan-induced paw edema, *Gongronema latifolium*.

INTRODUCTION

Plants are extremely an important source of medicinal agents. Since ancient times, plants have been used due to their impactful pharmacological properties. WHO estimates that 80% of the population living in rural areas is dependent on herbal medicine for their health needs. Crude plant extracts in the form of infusion, decoction, tincture or herbal extracts have been traditionally used by the population for the treatment of various diseases¹. Plant used in treatment of diseases is as old as civilization and traditional medicines are still a major part of habitual treatments of different maladies².

Diabetes mellitus is the commonest endocrine disorder that affects more than 171 million people worldwide. It is a condition in which a person has a high blood sugar level. The reason may be either the body doesn't produce enough insulin, or because body cells don't properly respond to the insulin that is produced³. When the blood glucose increases for example, after eating food material, insulin is released from the pancreas to normalize the glucose level. If the body cells do not absorb glucose, the glucose accumulates in the blood, leading to vascular, neuron, and other complications⁴.

Our body is exposed to a variety of oxidizing agents. Oxidation is a chemical reaction that produces free radicals, leading to chain reactions that may damage cells⁵. Body is equally inbuilt with antioxidants to cater for the free radicals generated from the oxidants, so a balance between the production of free radicals and neutralization by antioxidants is remains maintained. Antioxidants are substances known to protect the body from damage caused by reactive oxygen species induced oxidative stress⁶. Oxidative stress results because of imbalance between formation and neutralization of free radicals by antioxidants, it results to oxidative stress. Oxidative stress has been implicated in the etiology of diseases such as cardiovascular diseases and lung cancer.

Antioxidants from natural sources have a higher bioavailability and therefore higher protective efficacy comparison of synthetic antioxidants⁷.

Gongronema latifolium (Asclepiadiaceae) is used for medicinal and nutritional purposes widely distributed in the southeastern states of Nigeria⁸. Apart from being used as bitter spice or flavouring agent in many traditional Nigerian dishes the plant leaves has been found very efficacious as an antidiarrhoeal and antitussive. Aerial parts are taken to treat cough, intestinal worms, dysentery, dyspepsia and malaria⁹. It

is also taken as a tonic to treat loss of appetite. A decoction of leaves or leafy stems is commonly taken to treat diabetes and high blood pressure¹⁰.

The objective of the present study was to evaluate the anti-hyperglycemic and anti-inflammatory properties of methanol leaf extract of *Gongronema latifolium*.

MATERIALS AND METHODS

Collection and preparation of leaf samples

Fresh leaves and roots of *G. latifolium* were collected from Niger Delta and authenticated.

Extraction of plant materials

Fresh leaves of *G. latifolium* were washed and allowed to dry air at room temperature (28 to 30°C) for two weeks, to avoid the escape of volatile components by oven-drying. The dried leaves were milled into fine powder using the Christy-Norris hammer mill and passed through a 1mm sieve to obtain a fine powder. 2 kg of the sample was percolated in methanol for 48 hrs; this was then filtered through a Whatman filter paper No.1. The filtrate was evaporated to dryness on a hot plate at an initial temperature of 100°C and the dry powder obtained was suspended in 10ml distilled water, stirred and refiltered¹¹.

Assessment of hypoglycaemic activity

The approval of the Institutional Animal Ethics Committee was obtained before starting the study. An international protocol for conducting experiments on animals were followed. Healthy wistar rats of either sex having weight 150-200 g were selected for this activity¹². They were housed in standard condition of temperature (25±2°C) with 12 h light per day cycle. Before induction of diabetes weigh and normal glucose levels of rats were determined and recorded as Day 0. The acclimatized animals were fasted for 24 h with water ad libitum¹³.

Hyperglycemia was induced using a single intraperitoneal injection of alloxan monohydrate (160 mg/kg).

All animals were returned to their cages and given free access to food and water. Blood glucose levels were monitored by using a Glucometer after 72 h of injection and recorded as 1st day.

Rats with fasting blood glucose >7.8 mmol/l or 140 mg/dl were considered hyperglycemic and were selected for the study.

Diabetic rats were randomly assigned to 5 groups, each group contains six animals.

The animals were grouped as follow

- Group I: Normal control
- Group II: Diabetic control
- Group III: Diabetic rats treated with *G. latifolium* extract (250 mg/kg)
- Group IV: Diabetic rats treated with *G. latifolium* extract (500 mg/kg)
- Group V: Diabetic rats treated with glibenclamide (0.25 mg/kg).

Blood samples were obtained from the cut tail tip of the conscious rat and glucose test strip soaked with blood and then inserted to be read by the glucometer. Blood glucose levels were examined after 2, 12, 24, 72 hrs of orally administration of test drugs¹⁴.

Anti oxidant study

DPPH antioxidant assay

The free radical scavenging activity of *G. latifolium* extract was analyzed by, 1, 1-diphenyl-2-hydrazyl (DPPH) photometric assay. 2 ml of test extract at concentrations ranging from 10 to 400 µg/ml was each mixed with 1 ml of 0.5mM DPPH in methanol¹⁴. Absorbance at 520 nm was taken after 30 min incubation in the dark at room temperature using a spectrophotometer¹⁵. The experiment was done in triplicates and the percentage antioxidant activity was calculated as follows:

$$\% \text{ of DPPH free radical scavenging} = (\text{Absorbance of (control} - \text{blank)}) / (\text{Absorbance of control}) \times 100$$

Where 1ml methanol and 2ml extract were used as blank, while 1 ml 0.5 mM DPPH solution and 2 ml methanol was used as control. Ascorbic acid was used as reference standard.

RESULTS AND DISCUSSION

The anti-hyperglycemic effect of methanol extract of *G. latifolium* leaves was evaluated for 21 days. Alloxan-induced hyperglycemia is due to selective toxicity of alloxan on the pancreatic beta cells, generation of superoxide radicals and cytotoxic action mediated by generation of reactive oxygen species. The cytotoxic action of alloxan is mediated by reactive oxygen species, with a simultaneous massive increase in cytosolic calcium concentration, leading to a rapid destruction of beta cells¹⁶.

The effect of administration of extracts of *G. latifolium* on blood glucose level of diabetic rats is shown in Table 1. Treatment of extract of *G. latifolium* showed a significant reduction in the blood glucose level.

It also shows that 0.25 mg/kg glibenclamide is lowering glucose level significantly compared to normal control. Leaf extract at a dose of 500 mg/kg was more effective in reducing the blood glucose level than a dose of 250 mg/kg. The variation in blood glucose level in normal and experimental rats on 0, 7, 14 and 21 days of treatment has also been recorded. Treatment with *G. latifolium* extract showed signs of recovery as comparable with the standard drug glibenclamide. There was significant loss in body weight of diabetic rats compared to normal rats. In diabetics, glucose is not available therefore the cells use alternatively proteins for energy; consequently due to excessive breakdown of tissue protein a loss in body weight occurs. Body weight slightly increased in the normal control rats compared to initial body weight, whereas in diabetic control rats there was a significant decrease in body weight. Groups that were treated with glibenclamide and *G. latifolium* extract (250 and 500 mg/kg) showed significant reduction in body weights. The final body weights of treated groups were significantly lower than the final weights of normal control group. Hence, present study showed a good antidiabetic response of leaf extract against the experimental animals. The antioxidant activity was done *in vitro* using 1, 1-diphenyl-2-hydrazyl (DPPH) spectrophotometric assays. Figure 1 shows the antioxidant activity of *G. latifolium* and ascorbic acid standard. The result showed that *G. latifolium* extract

caused a concentration dependent percentage increase of antioxidant activity. DPPH is a stable free radical, it accepts an electron or hydrogen radical to become a stable diamagnetic molecule which is widely used to investigate radical scavenging activity. The degree of discoloration indicates the radical-scavenging potential of the antioxidant. Result shows that DPPH scavenging was increased in a concentration dependent manner compared to ascorbic acid¹⁶.

Statistical Analysis

Graph Pad Prism Version 5.0 for Windows was used for all statistical analyses. Data are presented as mean±SEM and analyzed by one-way ANOVA followed by Dunnett's multiple comparison test.

CONCLUSION

At present scenario, many researchers are showing their interest in medicinal plants for routine scientific investigation of numerous plants extract for biological effects and potential therapeutic properties in human. Methanol leaf extract of *G. latifolium* have anti-hyperglycemic and anti-oxidant activities. The methanol extract have antihyperglycemic activity in diabetic rats, most likely to be associated with glucose uptake increasing mechanism. Lower doses of the extract should be tried in future study to establish the most appropriate dose for clinical trial. *G. latifolium* leaves has shown to be a potential anti-diabetic and thus it can be a promising source for anti-diabetic agent. In the present study, the observed DPPH scavenging activity of the methanolic extract of *G. latifolium* might be useful for the development of newer and more potent natural antioxidants. So, present study concluded that detailed investigation of plants used in local health traditions and pharmacological evaluation of these plants and their taxonomical relatives can lead to the development of invaluable plant drug for many dreaded diseases including diabetes mellitus.

CONFLICT OF INTEREST

No conflict of interest is related to this paper.

REFERENCES

- Bernhoft A. A brief review on bioactive compounds in plants, In: Bioactive compounds in plants – benefits and risks for man and animals, Oslo: The Norwegian Academy of Science and Letters. 2010; 11-17.
- Firn R. Nature's Chemicals. Oxford University Press, Oxford. 2010; 74-75.

- Piero MN, Nzaro GM, Njagi JM. Diabetes mellitus- a devastating metabolic disorder. As J Biomed Pharm Sci; 04 (40). 2014; 1-7.
- Chamnan P, Simmons RK, Forouhi NG, Luben RN, Khaw KT, Wareham NJ, Griffin SJ. Incidence of type 2 diabetes using proposed HbA1c diagnostic criteria in the European prospective investigation of cancer–Norfolk Cohort implications for preventive strategies. Diabetes Care. 2011; 34(4): 950-956.
- Del CM, Sacchetti G, Di Mattia C, Compagnone D, Mastrocola D, Liberatore L, Cichelli A. Contribution of the phenolic fraction to the antioxidant activity and oxidative stability of olive oil. J Agric Food Chem. 2004; 52(13): 4072–9.
- Cook NR, Albert CM, Gaziano JM. A randomized factorial trial of vitamins C and E and beta carotene in the secondary prevention of cardiovascular events in women: results from the women's antioxidant cardiovascular study. Arch Intern Med. 2007; 167(15): 1610–8.
- Jain S, Jain N. Antioxidant: literature review, World J Pharm Pharm Sci. 2016; 5(10): 365-377.
- Edet EE, Akpanabiatu MI, Eno AE, Umoh IB, Itam EH. Effects of *Gongronema latifolium* crude leaf extract on some cardiac enzymes of alloxan-induced diabetic rats. Afr J Biochem Res. 2009; 3(11):366-369.
- Balogun ME, Besong EE, Obimma JN, Mbamalu OS, Djibissie SFA. *Gongronema latifolium*: A phytochemical. nutritional and pharmacological review. J Phys Pharm Adv. 2016; 6(1):811-824.
- Morebise O. A review on *Gongronema latifolium*, an extremely useful plant with great prospects. Eur J Med Pl. 2015; 10(1):1-9.
- Morebise O, Fafunso MA, Makinde JM. Membrane stabilizing activity: Possible mechanism of action for the anti-inflammatory property of *Gongronema latifolium* leaves. Int J Biom Heal Sci. 2005; 1(1):15-19.
- Kuhara T. Consensus recommendations on effective institutional animal care and use committees. Information on overseas technologies in laboratory animal science. 1988; 7: 14-17.
- Naowaboot J, Pannangpetch, P, Kukongviriyapan V, Kongyingoes B, Kukongviriyapan U. Antihyperglycemic, antioxidant and antiglycation activities of mulberry leaf extract in streptozotocin-induced chronic diabetic rats. Plant Foods Human Nutrition. 2009; 64: 116-121.
- Batra S, Batra N, Nagori BP. Preliminary phytochemical studies and evaluation of antidiabetic activity of roots of *Cayratia trifolia* (L.) Domin in alloxan induced diabetic albino rats. J Appl Pharm Sci. 2013; 3:97-100.
- Chioma DE. Anti-inflammatory and anti-oxidant activities of methanol extract of *baphia nitida*. Univ J Pharm Res. 2016; 1(2): 42-47.
- Jorns A, Munday R, Tiedge M, Lenzen S. Comparative toxicity of alloxan, N-alkyl-alloxans and ninhydrin to isolated pancreatic islets *in vitro*. J Endocrinol. 1997; 155: 283-293

Table 1: The effect of different treatment on fasting blood glucose levels in alloxan induced diabetic rats

Treatment	0 th Day	Blood glucose level (mg/dl)		
		7 th Day	14 th Day	21 st Day
Normal control	90.4±0.25	93.21±0.26	92.47±0.22	91.65±0.15
Diabetic control	247.58±0.43	265.28±0.31	280.37±0.41	300.48±0.42
<i>G. latifolium</i> (250 mg/kg)	245.72±0.51	200.82±0.42	150.47±0.53	122.37±0.37
<i>G. latifolium</i> (500 mg/kg)	247.38±0.09	195.28±0.51	142.37±0.08	120.25±0.22
Glibenclamide (0.25 mg/kg)	248.46±0.13	179.23±0.38	125.38±0.06	112.37±0.07

N=6, p < 0.05

Table 2: The effect of different treatment on body weight of alloxan induced diabetic rats


Treatment	Change in body weight (gm)		
	Initial	10 th Day	21 th Day
Normal control	154±0.31	174.00±0.8	180.50±0.32
Diabetic control	181±0.28	165.00±0.64	140.67±0.43
<i>G. latifolium</i> (250 mg/kg)	158±0.17	147.67±2.10	148.33±0.7
<i>G. latifolium</i> (500 mg/kg)	174±0.36	149.00±0.7	144.37±1.2
Glibenclamide (0.25 mg/kg)	176±0.28	157.00±0.53	167.83±0.9

N=6, p < 0.05

Table 3: Antioxidant activity of the methanol extract of *G. latifolium*

Sample	Conc (µg/ml)	Absorbance	Absorbance of control	Mean % DPPH scavenging activity
Ascorbic acid	10	0.069	0.812	86.43±0.17
	100	0.060		90.63±0.32
	200	0.049		89.49±0.53
	400	0.038		93.69±0.82
Methanol extract of <i>G. latifolium</i>	10	0.269		70.33±0.38
	100	0.250		76.09±0.24
	200	0.215		78.18±0.31
	400	0.190		80.27±0.78

N=3, p<0.05

Figure 1: Percentage antioxidant activity of *G. latifolium* extract using DPPH photometric assay.**Cite this article-**

Chidi E, Nwobodo NN, Paul NAC. Anti-hyperglycemic and anti-oxidant activities of methanol extract of *Gongronema latifolium*. Universal Journal of Pharmaceutical Research. 2017; 2(2): 35-38.