



Available online on 15.07.2017 at <http://ujpr.org>

Universal Journal of Pharmaceutical Research

An International Peer Reviewed Journal

Open access to Pharmaceutical research

© 2017, publisher and licensee UJPR, This is an open access article which permits unrestricted non commercial use, provided the original work is properly cited

Volume 2, Issue 3, 2017

REVIEW ARTICLE

SOLID DISPERSION TECHNOLOGY, A CONTEMPORARY OVERVIEW ON A WELL ESTABLISHED TECHNIQUE

Mahmoud A. Younis

Department of Industrial Pharmacy, Faculty of Pharmacy, Assiut University, Assiut 71526, Egypt.

*Corresponding Author's Email: mahmoud_ab32@yahoo.com

DOI: <http://doi.org/10.22270/ujpr.v2i3.RW1>

Received 17 June 2017; Revised 25 June; Accepted 1 July, Available online 15 July 2017

ABSTRACT

Solubility is a significant physicochemical parameter that affects the absorption, bioavailability and therapeutic effectiveness of any drug. Formulation development would fail if drug has a poor aqueous solubility. The low aqueous solubility of drug substances will lead to inadequate absorption and consequently, low bioavailability. Improvement of the aqueous solubility and dissolution rate of hydrophobic drugs remains one of the most difficult challenges in drug development process. Among various techniques used to improve poor aqueous solubility of drugs, solid dispersion technology has been extensively used in the literature and has become one of the well-established pharmaceutical procedures during formulation process. Although the technique seems to be "a part of the past", literature tells us that it is still used and developed to suit current needs of pharmaceutical industry. This review article highlights recent advances in solid dispersion technology and its applications in contemporary pharmaceutical research.

Keywords: Dissolution rate, hydrophilic carriers, poorly-soluble drugs, solid dispersion, solubility.

INTRODUCTION

Solubility is a significant physicochemical parameter that affects all characteristics of a drug substance¹. According to literature, about 70% of newly-discovered drug substances have aqueous solubility problems². This represents a great challenge to pharmaceutical industry as it leads to poor absorption, poor bioavailability and probably, failure of treatment¹. Over decades, scientists have developed several approaches to overcome solubility problems including the use of surfactants³, cosolvents³, inclusion complexes⁴, self-emulsifying systems⁵, prodrugs⁶ and micro-environmental buffering systems⁷. Solid dispersion technology has become one of the well-established techniques for solubilization of poorly-soluble drugs⁸. It has been extensively used in pharmaceutical literature for solubilization of massive number of hydrophobic drugs. Although the technique seems to be "a part of the past", literature tells us that it is still be used and developed to suit current needs of pharmaceutical industry. This review article highlights recent advances in solid dispersion technology and its applications in contemporary pharmaceutical research.

History and definitions of solid dispersion technology

In 1961, Sekiguchi and Obi⁹ first used solid dispersions technology to increase the dissolution and oral absorption of poorly water-soluble drugs. They assumed the formation of an eutectic mixture of a

poorly water-soluble drug with a physiologically inert, hydrophilic carrier. In 1966, Mayersohn and Gibaldi¹⁰ described solid dispersion as a "solid-state dispersions" or "a solid solution". In 1971, Chiou and Riegelman¹¹ defined solid dispersion as "the dispersion of one or more active ingredients in an inert carrier matrix at solid-state prepared by the melting, solvent or melting-solvent method". In 1985, Corrigan¹² defined the term as "a product formed by converting a fluid drug-carrier combination to the solid state". Recently, Dhirendra *et al.*¹³ defined solid dispersion as "a group of solid products consisting of at least two different components, generally a hydrophilic matrix and a hydrophobic drug. The matrix can be either crystalline or amorphous. The drug can be dispersed molecularly, in amorphous particles (clusters) or in crystalline particles".

CLASSIFICATION OF SOLID DISPERSION SYSTEMS

In 2007, Chokshi *et al.*¹⁴ classified solid dispersion systems according to the crystalline status of drug within the system. If the drug is converted to amorphous form and forms one phase system with polymer, it can be classified as a "solid solution", whereas if the drug exists as microcrystalline dispersion, i.e., forms two-phase system, it is generally referred to as a "solid dispersion". In 2015, Meng *et al.*¹⁵ extended the classification according to the

crystalline status of both drug and carrier. They classified the systems as the following:

- i. **Class C-C:** Crystalline active pharmaceutical ingredient (API) dispersed in crystalline carrier.
- ii. **Class C-A:** crystalline API dispersed in amorphous carrier.
- iii. **Class A-C:** Amorphous API dispersed in crystalline carrier.
- iv. **Class A-A:** Amorphous API dispersed in amorphous carrier.
- v. **Class M-C:** API molecularly dispersed in crystalline carrier.
- vi. **Class M-A:** API molecularly dispersed in amorphous carrier.

Examples of drugs formulated as solid dispersions

A massive number of drugs have been formulated as solid dispersions in the pharmaceutical literature (Table 1).

Advantages of solid dispersion technology

The major advantage of solid dispersions is that they improve the solubility of poorly water soluble drugs in pharmaceutical dosage forms which results in rapid dissolution rates and improved bioavailability of drugs. Moreover, the approach also offers others advantages including¹⁶:

- i. Rapid dispersion into water which is beneficial in the formulation of fast-dissolving tablets that can be used as an alternative to parenteral therapy enabling patient for self-medication even without the aid of water.
- ii. High degree of particle size reduction which is advantageous over traditional grinding in that it is suitable for waxy substances produces no dust and there is no danger for explosion.
- iii. Ability to produce a controlled pattern of drug release using the suitable retarding excipients like cellulose derivatives.
- iv. Ability to formulate liquid drugs into solid dosage forms through their mixing with molten carriers followed by rapid cooling and pulverization. This method was successful with several liquid drugs including methyl salicylate and clofibrate.
- v. Solid dispersions can result in precipitation of drugs in amorphous state with higher solubility and dissolution rate than the more stable crystalline state.

Disadvantages of solid dispersion technology

1. Most of the polymers used in solid dispersions can absorb moisture, which may results in phase separation, crystal growth or conversion from the amorphous state to the crystalline state or from a metastable crystalline form to a more stable structure during storage. This may results in decreased solubility and dissolution rate¹⁷.
2. Poor scale-up for the purposes of manufacturing.¹⁷
3. Stability problems: Younis¹⁸ reported the dissolution rate lowering of solid dispersions on aging owing to recrystallization of the amorphous drug in system and/or polymorphic transitions that are frequently associated with the aged solid dispersions.

COMMON MATERIALS USED IN PREPARATION OF SOLID DISPERSIONS

There are several materials that can be used as hydrophilic carriers in preparation of solid dispersions. Some examples are briefly discussed below.

Polyethylene glycols (PEGs)

Polyethylene glycols (PEGs) are polymers of ethylene oxide, with a molecular weight falling in the range 200-300,000. For the manufacture of solid dispersions, PEGs with a molecular weight 1500-20,000 are usually employed. PEGs have many advantages including high aqueous solubility, low cost, low toxicity and low melting points (below 65 °C) making them suitable carriers for preparation of solid dispersions by melting method.¹⁹

Polyvinylpyrrolidone (PVP)

Polymerization of vinylpyrrolidone leads to polymers of molecular weight ranging from 2500 to 300,000. They are inert, safe, highly water-soluble and soluble in a wide variety of solvents including alcohol. The main disadvantages of PVPs are hygroscopicity and high melting points (above 265 °C) making them more suitable for preparation of solid dispersions by solvent-based methods instead of heating-based methods.²⁰

Urea

It is the end product of protein metabolism and can also be synthesized by chemical reactions. It is highly water-soluble, soluble in many common organic solvents and has a moderate melting point (132-135°C) making it suitable for preparation of solid dispersions by different methods.²¹

Sugars

Although sugars have high aqueous solubility, they have many drawbacks regarding their use as carriers in solid dispersions. Most of them have high melting points making them problematic in the preparation of solid dispersions by heating methods. They are also poorly-soluble in most organic solvents creating problem in preparation of solid dispersions by solvent methods. Despite these draw backs, mannitol and sorbitol were reported to be used as carriers for many drugs²².

Poloxamers

Poloxamers are poly (oxyethylene)-poly (oxypropylene) copolymers, with trade names as Supronic, Pluronic or Tetronic. They have been introduced in 1950 and were since then very famously used in diverse pharmaceutical applications. They are composed of two hydrophilic chains of polyethylene oxide (PEO) sandwiching one hydrophobic polypropylene oxide chain (PPO). They are classified according to the proportions of hydrophilic and hydrophobic chains including poloxamer 124, 188, 237, 338 and 407 as the most common types. They are used as gelling agents, surfactants, stabilizers and hydrophilic carriers. They are soluble in different solvents and have low melting points (52-57°C) making those suitable carriers for preparation of solid dispersions by different methods.²³

METHODS OF PREPARATION OF SOLID DISPERSIONS

There are several methods used to prepare solid dispersions which have been developed from simple manual procedures to advanced techniques requiring special equipment to fulfill the needs of modern pharmaceutical industry. Some of these various techniques are briefly discussed below.

1. Co-melting method

This method involves the preparation of physical mixture of a drug and a water-soluble carrier and heating it directly till melting. The molten mixture is then solidified rapidly in an ice-bath under vigorous stirring. The final solid mass is crushed, pulverized and sieved. The modification in the method can be done by pouring the homogenous melt in the form of a thin layer onto a ferrite plate or a stainless steel plate and cooled by flowing air or water on the opposite side of the plate. In addition, a super-saturation of a solute or drug in a system can often be obtained by quenching the melt rapidly from a high temperature. Under such conditions, the solute molecule is arrested in the solvent matrix by the instantaneous solidification process. The quenching technique gives a much finer dispersion of crystallites when used for simple eutectic mixtures. Advantage of co-melting method is that it is economic and solvent less process, however this method is not suitable for the drug or carrier which is unstable at fusion temperature or evaporates at higher temperature. Some of the means to overcome these problems could be by heating the physical mixture in a sealed container or melting it under vacuum or in presence of inert gas like nitrogen to prevent oxidative degradation of drug or carrier²⁴.

2. Fusion method

It is a modification of co-melting method. The carrier is placed in a porcelain dish and heated till melting over steam bath. The accurately weighed amount of drug is dispersed into molten carrier gradually using a glass rod. After complete dispersion of drug within carrier, the dish is removed from steam bath and left aside to cool at room temperature till solidification of its contents. Then, the solid dispersion formed is pulverized and sieved. This method is useful in reducing thermal decomposition of drugs²⁵.

3. Solvent evaporation method

Both drug and carrier are dissolved in a common volatile solvent which is then, removed under vacuum. The formed solid dispersion is pulverized and sieved²⁶.

4. Kneading technique

In this method, carrier is permeated with water and transformed to paste. Drug is then added and kneaded for particular time. The kneaded mixture is then dried and passed through sieve if necessary. This method is suitable for thermolabile drugs but, it is not suitable for drugs sensitive to moisture²⁷.

5. Co-precipitation method

Required amount of drug is added to the solution of carrier. The system is kept under magnetic agitation and protected from the light. The formed precipitate is separated by vacuum filtration and dried at room temperature²⁸.

6. Co-grinding method

Physical mixture of drug and carrier is mixed for some time employing a blender at a particular speed. The mixture is then charged into the chamber of a vibration ball mill. The powder mixture is pulverized. Then, the product is collected and kept at room temperature in a screw capped glass vial until use²⁹.

7. Gel entrapment technique

Hydroxyl propyl methyl cellulose (used as a carrier) is dissolved in organic solvent to form a clear and transparent gel. Then, the drug is dissolved in gel by sonication for few minutes. Organic solvent is evaporated under vacuum. Solid dispersions are reduced in size by mortar and pestle and sieved³⁰.

8. Spray drying method

Drug is dissolved in suitable solvent and the required amount of carrier is dissolved in water. Solutions are then mixed by sonication or other suitable method to produce a clear solution, which is then spray dried using a spray dryer to produce solid dispersion in form of fine, free-flowing particles³¹.

9. Electrospinning method

Electrospinning is a process in which solid fibers are produced from a polymeric fluid stream solution or melt delivered through a millimeter- scale nozzle. This process involves the application of a strong electrostatic field over a conductive capillary attached to a reservoir containing a polymer solution or melt and a conductive collection screen. Upon increasing the electrostatic field strength up to a critical value, charge species accumulated on the surface of a pendant drop destabilize the hemispherical shape into a conical shape (commonly known as Taylor cone). Beyond the critical value, a charged polymer jet is ejected from the apex of the cone and carried to the collection screen via the electrostatic force³². This technique has potential for the preparation of nanofibres and controlling the release of biomedicines. As it is simple and cheap, this technique can be utilized for the preparation of solid dispersions in future³³.

10. Freeze-drying method

This process consists of dissolving the drug and carrier in a common solvent, which is immersed in liquid nitrogen until it is fully frozen. Then, the frozen solution is further lyophilized. Although it is concluded in literature that this is a promising and suitable technique to incorporate drug substances in stabilizing matrices, the technique is poorly exploited for the preparation of solid dispersions due to economical reasons. Advantages of freeze drying include that the drug is subjected to minimal thermal stress during the formation of the solid dispersion and the risk of phase separation is minimized³⁴.

11. Supercritical fluid (SCF) method

Supercritical fluid methods are mostly applied with carbon dioxide (CO₂), which is used as either a solvent for drug and matrix or as an anti-solvent. This technique consists of dissolving the drug and the carrier in a common solvent that is introduced into a particle formation vessel through a nozzle, simultaneously with supercritical CO₂ (the gas is heated beyond its critical temperature and pressure). When the solution is sprayed, the solvent is rapidly

extracted by the SCF, resulting in the precipitation of solid dispersion particles on the walls and bottom of the vessel. Advantages of this technique include reduction of particle size and residual solvent content as well as the high yield³⁵.

12. Direct capsule filling

The technique includes direct filling of hard gelatin capsules with the liquid melt of drug and carrier. This molten dispersion forms a solid plug inside the capsule upon cooling to room temperature. Advantages include avoidance of grinding-induced changes in the crystallinity of drug, reduction of cross contamination and operator exposure in a dust-free environment, better fill weight and content uniformity³⁶.

CHARACTERIZATION OF SOLID DISPERSION SYSTEMS

There are several techniques used for characterization of solid dispersion systems including¹⁶:

- i. Microscopic methods: Scanning Electron Microscopy (SEM) and Transmission Electron Microscopy (TEM).
- ii. Thermal methods: Differential Scanning Calorimetry (DSC) and Dissolution Calorimetry (DC).
- iii. Powder X-ray diffractometry (P-XRD).
- iv. Infrared Spectroscopy (IR).
- v. Powder flowability: angle of repose, compressibility.
- vi. Solubility studies.
- vii. *In-vitro* dissolution rate studies.
- viii. *In-vivo* studies: bioavailability, pharmacokinetics.
- ix. Stability Studies (effect of humidity, recrystallization of amorphous drug).

CONCLUSION

Solid dispersion technology has been considered one of the well-established techniques in formulation of poorly-soluble drugs. Like every technique, it has several advantages and disadvantages that were discussed briefly in this article. There are several materials that can be used as carriers in formulation of solid dispersion ranging from natural sugars to synthetic polymers. There are several methods used in preparation of solid dispersions which vary from simple hand-made methods to advanced techniques requiring special equipment. Eventually, it can be concluded that solid dispersion technology still can be used and further developed to suit the needs of modern pharmaceutical industry.

CONFLICT OF INTEREST

No conflict of interest associated with this work.

REFERENCES

1. Leuner C, Dressman J. Improving drug solubility for oral delivery using solid dispersions. *Eur J Pharm Biopharm* 2000; 50(1): 47-60.
2. Yohei K, Koichi W, Manabu N, Shizuo Y, Satomi O. Formulation design for poorly water-soluble drugs based on biopharmaceutics classification system: Basic approaches and practical applications. *Int J Pharm* 2011; 420(1): 1-10.
3. Kawakami K, Oda N, Miyoshi K, Funaki T, Ida Y. Solubilization behavior of a poorly soluble drug under combined use of surfactants and cosolvents. *Eur J Pharm Sci* 2006; 28(1-2): 7-14.
4. Ruan LP, Yu BY, Fu GM, Zhu DN. Improving the solubility of ampelopsin by solid dispersions and inclusion complexes. *J Pharm Biomed Anal* 2005; 38(3): 457-464.
5. Balata GF, Essa EA, Shamardl HA, Zaidan SH, Abourehab MAS. Self-emulsifying drug delivery systems as a tool to improve solubility and bioavailability of resveratrol. *Drug Design Deve Therap* 2016; 10: 117-128.
6. Hu Longqin. Prodrugs: Effective solutions for solubility, permeability and targeting challenges. *I Drugs* 2004; 7(8): 736-742.
7. Tatavarti AS, Hoag SW. Micro environmental pH modulation based release enhancement of a weakly basic drug from hydrophilic matrices. *J Pharm Sci* 2006; 95(7): 1459-1468.
8. Yanbin Huang, Wei-Guo Dai. Fundamental aspects of solid dispersion technology for poorly soluble drugs. *Acta Pharmaceutica Sinica B* 2014; 4(1): 18-25.
9. Sekiguchi K, Obi N. Studies on absorption of eutectic mixture. I. A comparison of the behavior of eutectic mixture of sulfathiazole and that of ordinary sulfathiazole in man. *Chem Pharmaceut Bull* 1961; 9: 866-872.
10. Mayersohn M, Gibaldi M. New method of solid state dispersion for increasing dissolution rates. *J Pharm Sci* 1966; 55: 1323-1342.
11. Chiou WL, Reigelman S. Pharmaceutical applications of solid dispersion systems. *J Pharm Sci* 1971; 60(9): 1281-1302.
12. Corrigan OI. Mechanisms of dissolution of fast release solid dispersions. *Drug Dev Ind Pharm* 1985; 11: 697-724.
13. Dhirendra K, Lewis S, Udupa N, Atin K. Solid Dispersions: A Review. *Pak J Pharm Sci* 2009; 22 (2): 234-246.
14. Chokshi RJ, Zia H, Sandhu HK, Shah NH, Malick WA. Improving the dissolution rate of poorly water soluble drug by solid dispersion and solid solution-pros and cons. *Drug Delivery* 2007; 14(1): 33-45.
15. Fan M, Urvi G, Chauhan H. Classification of solid dispersions: correlation to (i) stability and solubility (ii) preparation and characterization techniques. *Drug Dev Ind Pharm* 2015; 41(9): 1401-1415.
16. Dalvi PB, Gerange AB, Ingale PR. Solid dispersion: strategy to enhance solubility. *J Drug Deliv Therap* 2015; 5(2): 20-28.
17. Singh S, Singh Baghel R, Yadav L. A review on solid dispersion. *Int J Pharm Life Sci* 2011; 2(9): 1078-1095.
18. Mahmoud A. Younis. Development and evaluation of new dosage forms of domperidone. Master thesis, Faculty of Pharmacy. Assiut University, Egypt 2016.
19. Ford JL, Stewart AF, Dubois JL. The properties of solid dispersions of indomethacin or phenylbutazone in polyethylene glycol. *Int J Pharm* 1986; 28(1): 11-22.
20. Teodorescu M, Bercea M. Poly(vinylpyrrolidone)-A versatile polymer for biomedical and beyond medical applications. *Polymer Plastics Tech Eng* 2015; 54(9): 923-943.
21. Okonogi S, Oguchi T, Yonemochi E, Puttipatkhachorn S, Yamamoto K. Improved dissolution of ofloxacin via solid dispersion. *Int J Pharm* 1997; 156: 175-180.
22. R Jachowicz. Dissolution rates of partially water-soluble drugs from solid dispersion systems. I. Prednisolone. *Int J Pharm* 1987; 35: 1-5.
23. Devi DR, Sandhya P, Hari BNV. Poloxamer: A Novel functional molecule for drug delivery and gene therap. *J Pharm Sci Res* 2013; 5(8): 159-165.
24. Kalaiselvan R, Mohanta GP, Manna PK, Manavalan R. Studies on mechanism of enhanced dissolution of albendazole solid dispersions with crystalline carriers. *Indian J Pharm Sci* 2006; 68(5): 599-607.
25. Akiladevi D, Shanmugapandiyam P, Jebasingh D, Basak S. Preparation and evaluation of paracetamol by solid dispersion technique. *Int J Pharm Pharm Sci* 2011; 3(1): 188-191.

26. Abd Alaziz DM, Sammour OA, Elshamy AA, Neseem DI. Formulation and evaluation of binary and ternary solid dispersions of domperidone by solvent evaporation method. *Afr J Pharm Pharmacol* 2014; 8(3): 66-80.
27. Ghareeb MM, Abdulrasool AA, Hussein AA, Noordin MI. Kneading technique for preparation of binary solid dispersion of meloxicam with poloxamer 188. *AAPS Pharm Sci Tech* 2009; 10(4): 1206-1215.
28. Shah N, Iyer RM, Mair HJ, Choi DS, Tian H, Diodon R, Fahrnich K, A Pabst-Ravot, K Tang, E Scheubel *et al.* Improved human bioavailability of vemurafenib, a practically insoluble drug, using an amorphous polymer-stabilized solid dispersion prepared by a solvent-controlled coprecipitation process. *J Pharm Sci* 2013; 102(3): 967-981.
29. Nokhodchi A, Talari R, Valizadeh H, Jalali MB. An investigation on the solid dispersions of chlordiazepoxide, *Int J Biomed Sci* 2007; 3(3): 211-216.
30. Bhise SB, Rajkumar M. Effect of HPMC on solubility and dissolution of carbamazepine III in simulated gastrointestinal fluids, *Asian J Pharm* 2008; 2(1): 38-42.
31. Bakatselo V, Oppenheim RC, Dressman JB. Solubilization and wetting effects of bile salts on the dissolution of steroids. *Pharm Res* 1991; 8(12): 1461-1469.
32. Hohman MM, Shin M, Rutledge G, Michael PB. Electrospinning and electrically forced jets. II. Applications, *Physics Fluids* 2001; 13(8): 2221-2236.
33. Neamark A, Rujiravanit R, Supaphol P. Electrospinning of hexanoyl chitosan, *Carbohydrate Polymers* 2006; 66: 298-305.
34. Van DJ, Drooge WL, Hinrichs MR, Visser HW Frijlink. Characterization of the molecular distribution of drugs in glassy solid dispersions at the nano-meter scale, using differential scanning calorimetry and gravimetric water vapour sorption techniques. *Int J Pharm* 2006; 310(1): 220-229.
35. Majerik V, Charbit G, Badens E, Horváth G, Szokonya L, Bosc N, N Teillaud N. Bioavailability enhancement of an active substance by supercritical antisolvent precipitation. *J Supercrit Fluids* 2007; 40(1), 101-110.
36. Serajuddin ATM, Sheen PC, Mufson D, Bernstein DF, Augustine MA. Effect of vehicle amphiphilicity on the dissolution and bioavailability of a poorly water soluble drug from solid dispersions. *J Pharm Sci* 1988; 77(5), 414-417.
37. Ahmed E. Aboutaleb, Sayed I. Abdel-Rahman, Mahrous O. Ahmed, Mahmoud A. Younis. Improvement of domperidone solubility and dissolution rate by dispersion in various hydrophilic carriers. *J App Pharm Sci* 2016; 6(7): 133-139.
38. Ahmed E. Aboutaleb, Sayed I. Abdel-Rahman, Mahrous O. Ahmed and Mahmoud A. Younis. Design and evaluation of domperidone sublingual tablets. *Int J Pharm Pharm Sci* 2016; 8(6): 195-201.
39. Vyas V, Sancheti P, Karekar P, Shah M, Pore Y. Physicochemical characterization of solid dispersion systems of tadalafil with poloxamer 407. *Acta Pharm* 2009; 59: 453-461.
40. Zaini E, Umar S, Firdaus N. Improvement of dissolution rate of valsartan by solid dispersion system using D (-) mannitol. *Asian J Pharm Clin Res* 2017; 10(3): 288-290.

Table 1: Examples of drugs formulated as solid dispersions.

Drug name	Use	References
Ampelopsin	Anti-inflammatory, antimicrobial, relieving cough, antioxidant, antihypertensive, hepatoprotective and anticarcinogenic.	4
Albendazole	Anthelmintic	24
Carbamazepine	Anti-epileptic	30
Chlordiazepoxide	Sedative and hypnotic	29
Domperidone	Anti-emetic	37, 38
Indomethacin	Anti-inflammatory	19
Meloxicam	Anti-inflammatory, analgesic	27
Paracetamol	Analgesic, antipyretic	25
Sulfathiazole	Antibacterial	9
Tadalafil	Treatment of erectile dysfunction	39
Valsartan	Antihypertensive	40



How to cite this article-

Mahmoud A. Younis. Solid dispersion technology, a contemporary overview on a well established technique. *Universal Journal of Pharmaceutical Research*. 2017; 2(3): 15-19.