Available online on 15.07.2021 at <http://ujpr.org>
Universal Journal of Pharmaceutical Research
 An International Peer Reviewed Journal

Open access to Pharmaceutical research
 This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial Share Alike 4.0 License which permits unrestricted non commercial use, provided the original work is properly cited

Volume 6, Issue 3, 2021

Open Access

RESEARCH ARTICLE

MENOPAUSAL SYMPTOMS ON THE QUALITY OF LIFE AMONG YEMENI WOMEN IN SANA'A CITY

Dua'a Jamal Mutahar¹ , Ali Mohamed Assabri² , Hassan Abdulwahab Al-Shamahy³ ,
 Ammar Jaber⁴ , Samar Al qadi¹

¹Arab Board for Medical Specialization Scientific Council for Family & Community Medicine.

²Prof of Community Medicine, Faculty of Medicine and Health Sciences, Sana'a University, Republic of Yemen.

³Department of Medical Microbiology and clinical Immunology, Faculty of Medicine and Health Sciences, Sana'a University.

⁴Dubai Medical Collage, Dubai Medical University, Dubai, UAE.

ABSTRACT

Background: Menopause is a physiological phenomenon that can strongly affect the quality of life (QOL) of women.

Objectives: To analyze the menopausal associated symptoms among contributors and their effect on quality of life. To also investigate sociodemographic factors affecting QOL.

Methods: A cross-sectional study was conducted among 188 perimenopause and menopausal Yemeni women selected randomly from outpatient clinics of the main public hospitals in Sana'a, Yemen. Data was acquired using the Menopause-Specific Quality of Life (MENQOL) questionnaire.

Results: The mean age at menopause was found to be 47 ± 3.4 . The majority of participants were postmenopausal (71.27 %), and (71.80 %) were married. The most prevalent symptoms were aching muscles and joints (55%), dissatisfaction with personal life (37.7%), night sweat (35%) and avoidance of intimacy (30%). The total scores of menopausal quality of life for each MENQOL domain revealed that the maximum mean score was recorded for physical domain (42.81 ± 17.87), followed by psychosocial (14.11 ± 5.72), then sexual (6.93 ± 2.46), and finally vasomotor domain (4.78 ± 3.10).

Conclusion: The physical and psychosocial domains had upper mean MENQOL scores than vasomotor and sexual domains. There is a need to educate women about menopause and its symptoms and the need to seek medical advice from specialists and find a center for that.

Key words: Menopausal symptoms, Menopause Specific Quality of Life Questionnaire (MENQOL), Quality of life (QOL), Yemeni women.

Article Info: Received 1 May 2021; Revised 4 June; Accepted 5 July, Available online 15 July 2021

Cite this article-

Mutahar DJ, Assabri AM, Al-Shamahy HA, Jaber A, Al qadi S. Menopausal symptoms on the quality of life among Yemeni women in Sana'a city. Universal Journal of Pharmaceutical Research 2021; 6(3):1-8.

DOI: <https://doi.org/10.22270/ujpr.v6i3.600>

Address for Correspondence:

Prof. Hassan A. Al-Shamahy, Faculty of Dentistry, Sana'a University, P.O. Box 775 Sana'a, Yemen, Tel-+967-770299847; E-mail: shmahe@yemen.net.ye

INTRODUCTION

Menopause is an innate phenomenon that have an effect on all women and occurs in a woman's life 12 months or more after her menstrual period stops due to decrease in ovarian function¹. The manifestations of menopausal symptoms vary with each individual. Every woman's knowledge of menopause is exclusive as she may experience the whole of the symptoms or not any of them². The duration, severity and impact of these symptoms vary from person to person and population to population³. These symptoms can be classified based on the time of presentation into acute, subacute, and chronic, which are then grouped into: vasomotor, somatic, psychological or sexual

complaints⁴. Similar to general health, menopause is influenced by a range of cultural, socioeconomic and lifestyle factors which affect women's lives to different levels⁵. Quality of life (QOL) of premenopausal and menopausal women are significantly impacted by social, cultural and economic settings in which they live. They have to adapt to different issues from hot flushes and night sweats to discomfort of vaginal dryness^{2,6}. Adequate elucidation of women and raising the perception of certain changes that arise during menopause aids women with greater preparedness to deal with these changes and ultimately boost their QOL⁷. In the absence of intervention, over 75% of these females will suffer the difficult situation of menopause which can lead to immense stress and

disability⁷. In Yemen, studies on women's and children's health have been limited, and recently few studies have been conducted and discussed topics related to maternal and child health, including the consequence of menopausal symptoms on the quality of life of Yemeni women⁸⁻¹⁶.

Furthermore, as a result of the dearth of understanding of changes initiated by menopause, women do not search for medical counsel for relief of difficult menopausal symptoms. This study was performed to evaluate the menopausal related symptoms among participants from Sana'a, Yemen, assess the impact of menopausal symptoms on their quality of life, and analyzing socio-demographic factors that influence QOL. Achieving these objectives is imperative, as menopause has not been comprehensively researched in Yemen. There is only one current study on menopause among Yemeni Women¹⁶. This study will help health care providers to develop plans for these women by providing them better awareness and elucidation of the impact of menopause on QOL and identifying sociodemographic factors related to decline in QOL. This will also help women to help them in early recognize the symptoms of menopause and seek appropriate medical care such as hormone replacement therapy (HRT) if needed, alternative treatment strategies and lifestyle interventions for enhancing QOL particularly through radio and television media in addition to information dissemination by general practitioners.

SUBJECTS AND METHODS

A cross-sectional study was carried out among premenopausal and menopausal Yemeni women aged among 35 to 54 years. The sample size was calculated using Epi Info version 7 software accordingly: 1- The number of women aged 35-55 in Sana'a city was estimated to be around 500,000, with a menopausal acute effect on QOL equal to 24%, and with an acceptable margin of error of 6.1%, the required sample size is 188 women aged 35-55 years with a confidence level of 95%. The participants were chosen based on random selection from female waiting areas in different outpatient clinics of the four major public hospitals in Sana'a using a stratified sampling (Proportionate to size).

The inclusion criteria: women that have not menstruated for a period of 3-12 months or had irregular menses within the last one year, women aged between 35 and 54 years, and those that attend the outpatient clinics of the main public hospitals in Sana'a (AL-Thawra, AL Kuwait, AL Gumhory, AL-Sabeen hospitals).

Data collection: The respondents were given guaranteed discretion of information, and then voluntary verbal permission was acquired from them. The consistency of the questionnaire was tested using a pilot study that was conducted on 20 participants. Prior to administering the questionnaire, it was translated to Arabic language and disseminated among 20 participants that attend outpatient clinics of major

public hospitals in Sana'a. The questionnaire was finalized based on notes and remarks of participants who responded to the pretest questionnaires, and then translated back to English. The questionnaire comprises two sections. The first part covers sociodemographic characteristics, and the second section involves assessment of QOL using the menopause specific quality of life (MENQOL) questionnaire³.

Data analysis

Analysis was carried out using epi info version 7. Socio-demographic characteristics and the frequency of menopausal symptoms were presented as means, standard deviations, and percentages. The total score of each item was the sum of scores, the higher score, the worse quality of life (poor QOL) and vice versa. The mean score was considered as cut-off for QOL evaluation. Student-T, ANOVA tests and Mann-Whitney/Wilcoxon, Kruskal-Wallis tests were utilized to comparatively analyze the QOL scores between each domain and selected demographic variables. The statistical significance was fixed at $P < 0.05$.

RESULTS AND DISCUSSION

This study is an attempt to resolve that the impact of menopausal symptoms experienced by women on their normal daily activities. This is important, as understanding the problems associated with menopause, at the onset, can provide solutions and treatment measures, eventually improving the QOL of a woman. With respect to the demographic characteristics selected for this study, the mean age of menopause was found to be 47.9 ± 3.4 (Table 1) which is consistent with some earlier studies performed in Yemen¹⁶, and Middle East countries^{17,18}. However, the mean age is lower compared to those reported in western countries which varied between 50.1 to 52.8 years¹⁹, but higher in countries such as India which recorded 45.02 ± 4.35 ²⁰. As regards the level of education, majority of the participants (77.66%) are illiterate, with only a small proportion having high education (3.7%) (Table 1).

Studies have reported variations in menopausal symptoms of women across the world. For instance, Asian women experience less vasomotor and psychological symptoms compared to women in Western countries²¹. The assessment of menopausal quality of life (MENQOL) for each domain analyzed in this study indicated that physical domain has the highest mean score (42.81 ± 17.87), followed by psychosocial (14.11 ± 5.72) then sexual (6.93 ± 2.46) and finally vasomotor (4.78 ± 3.10) (Table 2).

Figure 1 showed a majority of participants (82%) complained from mild physical symptoms reflecting good QOL and able to cope with their menopause symptoms, (70%) complained from moderate sexual symptoms reflecting average QOL, (63%) complained from moderate vasomotor symptoms reflecting average QOL and (59%) complained from moderate psychosocial symptoms reflecting average QOL.

Table 1: Distribution of the participants according to the Socio- Demographic characteristics (n=188) (Sana'a, Yemen 2018).

Characteristics	Frequency (%)	Characteristics	Frequency (%)
Age group in years		Oral contraceptives	
<37	36(19)	Users	96(51.06)
37-49	92(49)	Not users	92(48.94)
>49	60(32)		
Menopausal status		HRT	
Postmenopause	134(71.28)	Users	9(4.79)
Perimenopause	29(15.43)	Not users	179(95.21)
Surgical menopause	25(13.29)		
Marital status		Smoking	
Married	135(71.81)	Yes	61(32.45)
Widow	39(20.74)	No	127(67.55)
Divorced	11(5.85)		
Single	3(1.60)		
Number of children		Chewing khat	
1-3 children	22(11.70)	Yes	115(61.17)
4-7 children	87(46.28)	No	73(38.83)
>7 children	69(36.70)		
No children	10(5.32)		
Education status		Type of disease	
Illiterate	146(77.66)	HTN	34(37.36)
Read & write	25(13.30)	DM	36(39.56)
Basic school	7(3.72)	Heart disease	18(19.78)
Secondary school	3(1.60)	Cancer	3(3.30)
University & higher studies	7(3.72)		
Employment status		Chronic diseases	
Employed	9(4.79)	Yes	91(48.40)
Unemployed	179(95.21)	No	97(51.60)
Family income		BMI	
High income	2(1.06)	Underweight <22	34(18)
Moderate income	54(28.72)	Normal 22-32	124(66)
Low income	132(70.22)	Overweight >32	30(16)
Residence place		Physical activity	
Urban	126(67.02)	Yes	110(58.51)
Rural	62(32.98)	No	78(41.49)

BMI Categories: Underweight=<18.5; Normal weight=18.5–24.9; Overweight=25–29.9; Obesity=BMI of 30 or greater.

These findings are consistent with studies conducted in Saudi Arabia²⁰, Asian countries²², Malaysia²³ and Ethiopia²⁵.

The physical menopausal symptoms are mild in severity which indicates a good QOL (Table 2). This result is steady with a study conducted in Riyadh, SA²⁰. However, majority of the menopausal women in this study frequently experience muscle or joint aches (55.30%) which is in agreement with other studies carried out in Asian countries²², but higher than value reported in a similar study performed in Yemen (9.8%)¹⁶, and lower than those recorded in Saudi Arabia (83.9%-96.1%)¹⁸, Malaysia (80.1%)²³, Ibadan, Nigeria (74%)²⁴ and Ethiopia (65.9%)²⁵.

In addition, less than half of the participants (32%) complained about feeling lack of energy, which was lower than that reported from Makkah, SA (61.3%)²⁶. Also observed, (24%) reported flatulence or gas pain and (25%) mentioned feeling bloated, while (21.80%) reported frequent urination which is similar to a finding in Ibadan, Nigeria (18.3%)²⁴. This study also revealed that very few participants reported they experience weight gain (1%) and increased facial hair (0.50%) which are lower than the value recorded by a study conducted in West Bengal, India (5%)²⁷.

This disparity in the frequency of symptoms is considered to be random i.e. non-specific to the menopause and may be due to several factors, which is more plausible. Another prevailing menopausal symptom revealed in this study was vasomotor symptoms. The occurrence of vasomotor symptoms in current study showed moderate severity, which indicates average QOL. (33.50%) of the participants complained of hot flashes which is in accordance with studies performed in Makkah, SA. (29%)²⁶, although lower than values reported in Yemen(100%)¹⁶, Western, Turkey (96%)⁶, Alexandria Egypt (90%)¹⁸, in Abha, SA (81%)¹⁷, Ethiopia (65.9%)²⁵ and India (60%)²⁷. Furthermore, 35.10% reported night sweating as a menopausal symptom which is similar to the finding of a study performed in Northern India (36%)²⁸. However, this result is lower than that reported in Abha, SA (80.7%)¹⁷, but higher than that of a study conducted in Ibadan, Nigeria (18%)²⁴. Vasomotor symptoms are generally associated with hormonal changes during menopause periods; hence this disparity is due to hereditary or socio-cultural diversity²⁷.

Figure 1: Effect of menopausal symptoms on the quality of life (physical, vasomotor, psychosocial and sexual).

Table 2: Frequency of menopausal symptoms among participants (n=188) (Sana'a, Yemen 2018).

Symptoms present	Severe	Moderate	Mild	Very mild	Normal
Physical symptoms					
Frequency (%)					
Flatulence or gas pains	37(19.70)	46(24.50)	26(13.80)	51(14.90)	28(27.10)
Aching muscles or joints	51(27.10)	104(55.30)	14(7.40)	10(5.30)	9(4.80)
Feeling tired	45(23.90)	65(34.60)	29(15.40)	25(13.30)	24(12.80)
Feeling a lack of energy	37(19.70)	61(32.40)	18(9.60)	41(21.80)	31(16.50)
Aches in back of neck	45(23.90)	99(52.70)	18(9.60)	14(7.40)	12(6.40)
Decrease in physical strength	33(17.60)	67(35.60)	25(13.30)	51(27.10)	12(6.40)
Decrease in stamina	16(8.50)	39(20.70)	38(20.20)	68(36.20)	16(8.50)
Drying skin	7(3.7)	20(10.6)	47(25)	30(16)	84(44.70)
Weight gain	9(4.80)	2(1.10)	33(17.60)	18(9.60)	126(67)
Increased facial hair	1(0.50)	1(0.50)	181(96.30)	1(0.50)	5(2.70)
Changes in appearance	33(17.60)	2(1.10)	9(4.80)	11(5.90)	133(70.70)
Feeling bloated	42(22.30)	48(25.50)	32(17)	21(11.20)	45(23.90)
Low back pain	57(30.30)	100(53.20)	11(5.90)	6(3.20)	14(7.40)
Frequent urination	7(3.70)	41(21.80)	41(21.80)	17(9)	82(43.60)
Involuntary urination	4(2.10)	6(3.20)	18(9.60)	16(8.50)	144(76.60)
Vasomotor symptoms					
Hot Flashes	28(14.90)	63(33.50)	26(13.80)	24(12.80)	47(25)
Night Sweats	8(4.30)	66(35.10)	55(29.30)	16(8.50)	43(22.90)
Sweating	1(0.50)	17(9)	40(21.30)	32(17)	98(52.10)
Psychological symptoms					
Dissatisfaction of Personal Life	36(19.15)	71(37.77)	44(23.40)	26(13.83)	11(5.85)
Feeling Anxious or Nervous	37(19.68)	56(29.79)	14(7.40)	49(26.06)	16(8.51)
Poor Memory	18(9.57)	48(25.53)	23(12.23)	34(18.09)	65(34.57)
Accomplishing less than I used to	33(17.55)	57(30.32)	24(12.77)	42(22.34)	32(17.02)
Feeling down/depressed	21(11.17)	59(31.38)	39(20.74)	37(19.68)	32(17.02)
Being impatient with other people	27(14.36)	57(30.32)	35(18.62)	47(25)	22(11.70)
Feelings of wanting to be alone	13(6.91)	40(21.28)	38(20.21)	48(25.53)	49(26.06)
Sexual symptoms					
Change in sexual desire	1(0.68)	24(16.22)	44(29.73)	56(37.84)	23(15.54)
Vaginal Dryness	9(6.08)	38(25.68)	29(19.59)	26(17.57)	46(31.08)
Avoiding intimacy	17(11.49)	45(30.41)	45(30.41)	23(15.54)	18(12.16)
Relation with husband	16(8.51)	23(12.23)	40(21.28)	46(24.47)	63(33.51)

The moderate severity of the prevalence of psychosocial symptoms in this study indicates average QOL (Table 2). The most common psychosocial symptom is dissatisfaction with personal life (37.77%), lower than in Ethiopia (46.0%)²⁵ and Makkah, SA (44.8%)²⁶. (31.38%) of participants in this study, experienced depression which is lower than that Ethiopia (46.0%)²⁵ and West Bengal, India (88%)²⁷ but higher than that in Korea (5.9%)²⁹ and Yemen (5.5%)¹⁶. This wide difference in results is probably due to methodological disparity and economic situation

of Yemen, thus the depression may not be a menopausal entity, but a result of the current financial struggles³⁰. Likewise, complaints about anxiety or nervousness was (29.79%), which is higher compared to findings of previously in Yemen (14.5%)¹⁶, but lower than that in Makkah, SA (52%)²⁶, and Abha, SA (89.0%)¹⁷. The occurrence of poor memory was (25.53%) in the current study, which was relatively lower compared to that of Hyderabad Pakistan (62.10%)³¹.

The moderate severity of the prevalence of sexual symptoms indicates average QOL (Table 2). (25.68%) of the participants had vaginal dryness, which is similar to (26%) reported in West Bengal, India²⁷ and (26%) in Makkah, SA²⁶ but lower than (82%) in Yemen previously¹⁶. Moreover, (30.41%) of the participants avoided intimacy. This finding is compatible that in Turkey (39.7%)⁶. As for the women who did not feel anything at all (Not at all) for physical symptoms, vasomotor symptoms, psychological symptoms and sexual symptoms the rates were ranged from 4 to 52% (Table 2), which is in the ranges of similar studies in West Bengal, India²⁷ and in Makkah, SA²⁶ and in Turkey⁶.

Regarding the relation between MENQOL in different domains and menopausal status of participants (Table 3). The perimenopause group showed a comparatively higher and significantly different mean score for physical symptoms ($P=0.007$). This finding is consistent with studies conducted in

Riyadh, SA¹. Similarly, perimenopause group recorded the highest mean score for psychosocial symptoms, although without significant difference compared to the other menopausal statuses. Likewise, the postmenopausal group recorded the highest mean score for the sexual domain but without statistical significance, which is consistent with studies of Malaysia²³, but contradicts that of Nigeria where perimenopause group complained the most about sexual symptoms²⁴. The surgical menopause group recorded the highest mean score of vasomotor symptoms without statistical significance. This finding is consistent with a study conducted in Britain that revealed that surgical menopause group complained from the most about vasomotor symptoms³². This could be attributed to variation in levels of hormones, such as follicular-stimulating hormone and estrogen that occur in the course of menopausal transition, which gradually leads to decline in physical, psychosocial, vasomotor QOL³².

Table 3: The MENQOL mean scores in different domains (QOL) by menopausal status(n=188) (Sana'a, Yemen 2018).

	Domains of MENQOL			
	Physical	Vasomotor	Psychosocial	Sexual
Menopausal status				
Perimenopause	31.27±9.02	4.83±3.11	15.89±4.01	6.81±2.55
Postmenopausal	24.86±10.60	4.76±3.19	13.90±5.73	7.02±2.43
Surgical menopause	23.64±11.45	4.84±2.68	13.16±7.01	6.57±2.63
P value	$P=0.007^*$	$P=0.9$	$P=0.1$	$P=0.7$

*Kruskal-Wallis test

The menopausal specific quality of life (MENQOL) mean scores were obtained for the different domains in relation to sociodemographic factors of participants. Higher MENQOL scores indicate poorer QOL, and vice versa. This study found that mean scores for physical, vasomotor and psychosocial domains were significantly higher among participants who used HRT, with values of ($P=0.002$), ($P=0.001$) and ($P=0.01$) respectively (Table 3), as compared to those who had not used HRT. Their mean scores for sexual domain were also higher, but without statistical significance. This can be explained by the fact that a few participants used natural HRT as herbal remedies and vitamins for relieving menopausal symptoms, thus the absence of quality controls and unpredictability in the purity, potency and absorption of herbal remedies can result in under or over-dosing, eventually causing the treatment to be ineffective. Similarly; a cohort study among past hormone therapy users in Britain who had defaulted in their treatment were more prone to hot flashes and night sweats³². Furthermore, higher mean scores were recorded for vasomotor, psychosocial, and sexual symptoms among participants who not used oral contraceptives, while physical symptoms were more prevalent among users without significance association. Participants who had chronic diseases and smoking habit exhibited significantly higher vasomotor symptoms at ($P=0.03$) and ($P=0.004$), respectively, and displayed more physical, psychosocial, sexual symptoms without significance. This finding is

comparable with a study in Australia that revealed a relation between vasomotor symptoms and smoking⁴. Similar findings were reported in Mexico³³. However, a study performed in Riyadh, SA reported that smoking had no effect on the MENQOL score¹. This varying effect may be explained by the perplexing effects of chronic illness and the aging process which may impact on menopausal symptoms. Also, the nicotine contained in cigarettes, which like caffeine is a vasoconstrictor, so smokers are more likely to experience more hot flashes³³. In addition, participants who chew khat had significantly higher scores for physical and psychosocial symptoms at ($P=0.02$) and ($P=0.03$), respectively. They also exhibited higher scores for vasomotor symptom, but without significance. On the other hand, participants who do not chew khat had higher scores for sexual symptom without significance. This is plausible as the side effects of chewing khat include insomnia, anxiety, psychosis, decreased appetite and energy loss¹⁶. Regarding BMI, this study found no association with all the menopausal symptoms. However, physical, vasomotor symptoms were more prevalent among the overweight participants. A similar finding has been reported in Mexico³³. Conversely, more complaints were received from participants with normal weight with respect to psychosocial symptoms, while the underweight participants complained the most of sexual symptoms.

Table 4: The MENQOL mean scores in different domains of participants across sociodemographic factors (n=188) (Sana'a, Yemen 2018).

Socio-demographic factors	Domains of MENQOL			
	Physical	Vasomotor	Psychosocial	Sexual
Education				
Illiterate	26.31±11.02	4.84±3.13	14.21±5.98	6.80±2.62
Read & write	24.08±9.72	4.64±3.13	13.92±5.22	6.87±1.75
Basic school	21.85±5.87	6.14±2.41	12.43±3.99	7±1.78
Secondary school	14.66±4.93	2±2	10.66±5.13	7.33±1.15
University & higher studies	27±11.22	4±3.16	15.85±2.67	9±2.53
P value	P=0.2	P=0.3	P=0.6	P=0.3
Employment status				
Employed	28.77±11.42	4.55±4.79	16±4.24	8.13±2.29
Unemployed	25.53±10.69	4.79±3.12	14.01±5.77	6.85±2.46
P value	P=0.3	P=0.8	P=0.3	P=0.1
Residence place				
Rural	24.56±10.38	5.19±3.12	14.25±6.35	7.15±2.45
Urban	26.24±10.88	4.58±3.08	14.04±5.41	6.82±2.47
P value	P=0.3	P=0.2	P=0.8	P=0.4
Family income				
High	25±22.62	4.5±3.54	15±14.14	9.50±2.12
Moderate	25.85±12.53	4.53±3.15	14±5.83	6.66±2.51
Low	25.64±9.83	4.90±3.09	14.14±5.61	7±2.43
P value	P=0.8	P=0.7	P=0.9	P=0.2
Chronic disease				
Yes	26.58±10.92	5.28± 3.03	14.54±5.79	7.07±2.34
No	24.85±10.52	4.32±3.12	13.70±5.65	6.80±2.56
P value	P=0.3	P=0.03*	P=0.3	P=0.5
Oral contraceptive used				
Users	26±11.51	4.73±3.01	14.09±5.73	6.75±2.22
Not users	25.36±9.88	4.85±3.21	14.13±5.75	7.14±2.74
P value	P=0.8	P=0.7	P=0.9	P=0.3
HRT				
Users	39.66±8.03	7.88±2.20	18.66±4.38	7.85±1.95
Not users	24.98±10.36	4.63±3.06	13.88±5.69	6.87±2.48
P value	P=0.002**	P=0.001*	P=0.01*	P=0.3
Habit				
Smoking				
Yes	27.45±11.38	5.70±3.39	15.47±5.89	6.94±2.63
No	24.84±10.32	4.34±2.86	13.45±5.54	6.92±2.38
P value	P=0.7	P=0.004*	P=0.02*	P=0.9
Chewing Khat				
Yes	27.08±11.08	5.11±3.22	14.82±5.69	6.89±2.57
No	23.49±9.79	4.27±2.86	13±5.63	6.98±2.27
P value	P=0.02**	P=0.07	P=0.03*	P=0.8
Physical activity				
Yes	21.37±8.71	4.41±2.98	12.26±5.45	7.20±2.05
No	31.78±10.36	5.31±3.22	16.72±5.06	6.60±2.85
P value	P=0.001**	P=0.5	P=0.001*	P=0.1
BMI				
Underweight <22	24.60±12.79	4.48±3.59	12.96±6.52	5.28±1.95
Normal 22-32	25.54 ±10.52	4.74± 2.98	14.29±5.83	5.17±1.80
Overweight >32	27.12 ±25.92	5.18 ±3.23	14.25±4.51	5.25±1.71
P value	P=0.6	P=0.6	P=0.5	P=0.9

*T-test, **Mann-Whitney test

T-test: The independent samples t-test was used for two separate sets of independent and identically distributed samples were obtained, one from each of the two populations being compared. Mann-Whitney test is to assume that: All the observations from both groups are independent of each other. The responses are at least ordinal (i.e., one can at least say, of any two observations, which is the greater). Under the null hypothesis H0, the distributions of both populations are equal. The alternative hypothesis H1 is that the distributions are not equal.

This finding is consistent with a study carried out in Turkey, which revealed no association between BMI and menopausal symptoms³⁴. Contrarily, a study in Turkey showed that women with no education had non-significantly higher scores of physical,

psychological and vasomotor symptoms³⁵. This finding possibly indicates that awareness among educated Yemeni women towards menopausal symptoms and their impact on QOL was lower as compared to those who were not educated. In fact, a few employed

women among the participants experienced more symptoms, thus employment may initiate more stress in some women facing menopause compared to those who are jobless.

With regards to residence, the vasomotor, psychosocial, sexual symptoms were more dominant among participants who lived in rural areas while physical symptoms were more prevalent among participants who lived in urban areas. On the contrary, a study conducted in Ranbir, India revealed significantly higher scores of psychosocial, physical symptoms with respect to the residence place³⁶. This disparity in frequency of symptoms may be due to differences of culture, lifestyle, and diet. In relation to marital status, married participants complained more commonly of physical and vasomotor symptoms, while psychosocial symptoms were more prevalent among divorced participants. In fact, majority of the participants were married and complained about suffering from sexual symptoms, which may be due to hormonal changes in the menopause phase, leading to a decline in sexual desire and vaginal dryness. In contrast, the complaints from divorced participants about psychosocial symptoms may be due to the absence of social support and stability with a spouse³⁵. In addition, a few participants with high income level had higher mean scores of psychosocial, sexual symptoms, while participants with low income level had higher scores of vasomotor symptoms. Physical symptoms were more prevalent among participants with moderate level income. Unexpectedly, only a few of the participants with high income had better QOL. In contrast, study conducted in India reported an association of higher income with lower scores and better QOL³⁷.

CONCLUSION

The QOL of women is affected by symptoms associated with menopause. This study confirmed the negative impact of menopausal symptoms on QOL. The Yemeni women selected for this study complained more commonly of physical and psychosocial symptoms as compared to vasomotor and sexual symptoms. The most prevalent symptoms were aching muscles and joints, dissatisfaction with personal life, night sweat and avoidance of intimacy. However, there was a variation in severity of symptoms, as reflected by the average QOL. In spite of the acknowledged benefits of HRT, especially for relieving vasomotor and sexual symptoms, none of the participants used HRT mainly because majority are inadequately informed about its accessibility and effectiveness. Few participants used herbal products. However, there is no consistent evidence to support their efficacy and safety.

RECOMMENDATION

Health facilities and health service providers should be encouraged to actively participate in educating menopausal women about improving their quality of life and engaging in HRT, especially women undergoing surgical menopause, alternative treatment

strategies and lifestyle interventions. More community-based research is needed to address postmenopausal women's health needs and quality of life using a larger sample population and different geographic locations, rather than a facility-based study, when possible.

AUTHOR'S CONTRIBUTION

This research work is part of the MD study. The first author (DJM) did fieldwork and wrote the thesis. The corresponding author (HAA) and other authors supervised the work and reviewed and edited the paper.

ACKNOWLEDGEMENTS

We are grateful to all the staff and the health directors make it easy to allow us to collect data from participants and use the SECA Weight Scale Tool. We thank all study participants; otherwise this study would not have been fulfilled.

CONFLICT OF INTEREST

There is no conflict of interest with this research.

REFERENCES

1. Al Dughaiter A, Ateeq M, Al Mutairy H. Menopausal symptoms and quality of life among Saudi women visiting primary care clinics in Riyadh, Saudi Arabia. *Int J Womens Health* 2015; 7:645-653. <https://doi.org/10.2147/IJWH.S84709>
2. Ande AB, Omu OP, Ande OO, Olagbuji NB. Features and perceptions of menopausal women in Benin City, Nigeria. *Ann Afr Med* 2011; 10:300-4. <https://doi.org/10.4103/1596-3519.87048>
3. Bosworth HB, Bastian LA, Kuchibhatla MN, Steffens DC, McBride CM, Skinner CS, et al. Depressive symptoms, menopausal status and climacteric symptoms in women at midlife. *Psychosom Med* 2001; 63:603-8. <https://doi.org/10.1097/00006842-200107000-00013>
4. Hunter M, Rendall M. Bio-psycho-socio-cultural perspectives on menopause. *Baillieres Best Pract Res Clin Obstet Gynaecol* 2007; 21(2):261-274. <https://doi.org/10.1016/j.bpobgyn.2006.11.001>
5. Ahmed K, Jahan P, Nadia I, Ahmed F and AL-Emran A. Assessment of Menopausal Symptoms among Early and Late Menopausal Midlife Bangladeshi Women and their Impact on the Quality of Life. *Menopausal Med* 2016; 22(1):39-46. <https://doi.org/10.6118/jmm.2016.22.1.39>
6. Ayranli U, Orsal O, Arslan G and Emeksiz D. Menopause status and attitudes in a Turkish midlife female population: an epidemiological study. *BMC Women's Health* 2010; 10:1. <https://doi.org/10.1186/1472-6874-10-1>
7. Rotem M, Kushnir T, Levine R, Ehrenfeld M. A psycho-educational program for improving women's attitudes and coping with menopause symptoms. *J Obstet Gynecol Neonatal Nurs* 2005; 34(2):233-40. <https://doi.org/10.1177/0884217504274417>
8. Al-Rukeimi AA, Al-Hatami SM, Al-Danany DA, Al-Shamahy HA, Al-Rukeimi RA. Prevalence and risk factors associated with vulvovaginal candidiasis during pregnancy in Sana'a, Yemen. *Universal J Pharm Res* 2020; 5(3). <https://doi.org/10.22270/ujpr.v5i3.407>
9. Al Rukeimi AD, Al Shamahy, HA, et al. Association of Cytomegalovirus and Rubella Virus Infections in Pregnant Women with bad Obstetric History. *WJ Gynecol Women's Health* 2019; 2(3):1-5. <https://doi.org/10.33552/WJGWH.2019.02.000538>

10. Al-Rukeimi A AD, Jamial NA, Al Shamahy HA, *et al.* Assessment of bacterial contamination at the time of embryo transfer, and its impression on the *in-vitro* Fertilization/Pregnancy Outcome, in Sana'a City, Yemen. *W J Gynecol Women's Health* 2019; 2(2). <https://doi.org/10.33552/WJGWH.2019.02.000533>
11. Almoaish AMA, Al-Shamahy HA, Al-Hajj MA, *et al.* Prevalence of cytomegalovirus IgG antibodies, potential risk factors and awareness of congenital cytomegalovirus among female doctors. *Universal J Pharm Res* 2018; 3(5):1-5. <https://doi.org/10.22270/ujpr.v3i5.199>
12. Aljedry ZA, Shaib SH, Al-Shamahy HA, Al-Jaufy AY. Tetanus immunization among pregnant women: coverage rate and rate of protection at time of delivery. *Universal J Pharm Res* 2019; 4 (1):1-6. <https://doi.org/10.22270/ujpr.v4i1.233>
13. Edrees WH, Al-Asbahi AA, Al-Shehari WA, Qasem EA. Vulvovaginal candidiasis prevalence among pregnant women in different hospitals in Ibb, Yemen. *Universal J Pharm Res* 2020; 5(4):1-5. <https://doi.org/10.22270/ujpr.v5i4.431>
14. Al Rukeimi AAD, Al Shaibani E M, Al Shamahy HA, Al Hadad AM, Al Tabreezi HM, *et al.* Uterine rupture and associated factors during labor amongst women delivered in Saudi Hajjah Hospital in Hajjah City North West Yemen. *W J Gynecol Women's Health* 2018; 1(2):1-8. <https://doi.org/10.33552/WJGWH.2018.01.000508>
15. Alqaisi NA, AL-Mekhlafi AM, Al-Shamahy HA, *et al.* Toxoplasmosis in pregnant women in Yemen: the immune status and potential risk factors. *Universal J Pharm Res* 2021; 6(2): 1-6. <https://doi.org/10.22270/ujpr.v6i2.570>
16. Abdul Halim F, Mehras A, Assabri A, Alkobaty M, Alawadi S and Albourhi A. Menopause among Yemeni Women. *Scientific Research Publishing* 2018; 7:65-77. <https://doi.org/10.4236/aar.2018.74006>
17. Al-Musa H, Ahmed R, Alsamghan A, *et al.* The prevalence of symptoms experienced during menopause, influence of socio-demographic variables on symptoms and quality of life among women at Abha, Saudi Arabia. *J Biomed Res* 2017; 28(6):1-6. <https://doi.org/10.5114/jpm.2015.48637>
18. El-Juhany L, Abdel Aziz F.A, Dabbous N and Hassan M. Women's perception and experience of menopause: a community-based study in Alexandria, Egypt. *East. Mediterr. Health J* 2006; 12(2):1-10.
19. Palacios S, Henderson VW, Siseles N, Tan D, Villaseca P. Age of menopause and impact of climacteric symptoms by geographical region. *Climacteric J* 2010;13(5):419-428. <https://doi.org/10.3109/13697137.2010.507886>
20. Kapur P, Sinha B, Pereira B. Measuring climacteric symptoms and age at natural menopause in an Indian population using the Greene climacteric scale. *Menopause* 2009; 16(2):378-384. <https://doi.org/10.1097/gme.0b013e31818a2be9>
21. Fuh JL, Wang SJ, Lu SR, Juang KD, Chiu LM. The Kinmen women-health investigation (KIWI): a menopausal study of a population aged 40-54. *Maturitas* 2001; 39(2):117-24. [https://doi.org/10.1016/s0378-5122\(01\)00193-1](https://doi.org/10.1016/s0378-5122(01)00193-1)
22. Gartoulla P, Bell R, Fradkin P, Davis S. Prevalence of menopausal symptoms in Asian midlife women: a systematic Prevalence of menopausal symptoms in Asian midlife women: a systematic review. *Climacteric J* 2015; 18(2):157-176. <https://doi.org/10.3109/13697137.2014.937689>
23. Rahman SA, Zainudin SR, Mun VL. Assessment of menopausal symptoms using modified Menopause Rating Scale (MRS) among middle age women in Kuching, Sarawak, Malaysia. *Asia Pac Fam Med* 2010; 9:5-10. <https://doi.org/10.1186/1447-056X-9-5>
24. Ibraheem OM, Oyewole OE, Olaseha IO. Experiences and perceptions of menopause among women in Ibadan South East Local Government area, Nigeria. *Afr J Biomed Res* 2015; 18:81-94.
25. Yisma E, Eshetu N, Ly S and Dessalegn B. Prevalence and severity of menopause symptoms among perimenopausal and postmenopausal women aged 30-49 years in Gulele sub-city of Addis Ababa, Ethiopia. *BMC Women Health J* 2017; 17: 124. <https://doi.org/10.1186/s12905-017-0484-x>
26. Mohammed H, Lamadah S, Al Zamil L. Quality of life among menopausal women. *Glob J Bio Agri Health Sci* 2014; 4(11):1-6. https://doi.org/10.4103/jmh.JMH_78_16
27. Karmakar N, Majumdar S, Dasgupta A, Das S. Quality of life among menopausal women: A community-based study in a rural area of West Bengal. *J Mid-life Health* 2017; 8:21-7. https://doi.org/10.4103/jmh.JMH_78_16
28. Vijayalakshmi S, Chandrababu R, Eilean Victoria L. Menopausal transition among Northern Indian women. *Nitte Univ J Health Sci* 2013; 3:73-79. <https://doi.org/10.1055/s-0040-1703658>
29. Sun P, Junga J, Shinb A, Kangab D. Menarche age, menopause age and other reproductive factors in association with post-menopausal onset depression: Results from Health Examinees Study (HEXA). *J Affect Disord* 2015;187:127-135. <https://doi.org/10.1016/j.jad.2015.08.047>
30. Kulkarni J. Perimenopausal depression-an-under-recognised entity. *Aust Prescr.* 2018; 41(6):183-185. <https://doi.org/10.18773/austprescr.2018.060>
31. Nusrat N, Nishat Z, Gulfareen H, Aftab M and Asia N. Knowledge, attitude and experience of menopause. *J Ayub Med Coll Abbottabad* 2008; 20(1):56-9. [PMID: 19024187.](https://pubmed.ncbi.nlm.nih.gov/19024187/)
32. Hunter MS, Gentry-Maharaj A, Ryan A, *et al.* Prevalence, frequency and problem rating of hot flushes persist in older postmenopausal women: Impact of age, body mass index, hysterectomy, hormone therapy use, lifestyle and mood in a cross-sectional cohort study of 10,418 British women aged 54-65. *BJOG* 2012; 119:40-50. <https://doi.org/10.1111/j.1471-0528.2011.03166.x>
33. Ortega-Ceballos PA, Moran C, Blanco-Munoz J, *et al.* Reproductive and lifestyle factors associated with early menopause in Mexican women 2006; 48 (4):300-307. <https://doi.org/10.1590/s0036-36342006000400004>
34. Neslisah M, Kartal M, Guldal D. The effect of physical activity and body mass index on menopausal symptoms in Turkish women: a cross-sectional study in primary care. *BMC Women's Health* 2014; 14:38-42. <https://doi.org/10.1186/1472-6874-14-38>
35. Yanikkerem E, OrucKoltan S, Goker Tamay A, Dikayak S. Relationship between women's attitude towards menopause and quality of life. *Climacteric J* 2012; 15(6):552-562. <https://doi.org/10.3109/13697137.2011.637651>
36. Sharma S, Mahajan N. Menopausal symptoms and its effect on quality of life in urban versus rural women: A cross-sectional study. *J Midlife Health* 2015; 6(1):16-20. <https://doi.org/10.4103/0976-7800.153606>
37. Poomalar G K, Bupathy Arounassalame. The quality of life during and after menopause among rural women. *J Clin Diagnost Res* 2013; 1:135-9. <https://doi.org/10.7860/JCDR/2012/4910.2688>