



Available online on 15.9.2017 at <http://ujpr.org>
Universal Journal of Pharmaceutical Research

An International Peer Reviewed Journal

Open access to Pharmaceutical research

This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial Share Alike 4.0 License which permits unrestricted non commercial use, provided the original work is properly cited

Volume 2, Issue 4, 2017



Open Access

REVIEW ARTICLE

AN UPDATED REVIEW ON TRANSFERSOMES: A NOVEL VESICULAR SYSTEM FOR TRANSDERMAL DRUG DELIVERY

Neelam Chauhan¹ , Kapil Kumar¹ , Navin Chandra Pant² 

¹Global Institute of Pharmaceutical Education and Research, Kashipur, Uttarakhand, India

²Bhagirathi Institute of Pharmacy, Lohaghat, Uttarakhand, India

ABSTRACT

Transdermal route is an interesting option in this respect because a transdermal route is convenient and safe, avoid first pass metabolism, predictable and extended duration of activity, minimizing undesirable side effects, utility of short half-life drugs, improving physiological and pharmacological responses, avoiding the fluctuation in drug levels and inter and intra-patient variations. However it has got its own limitations its inability to transport large molecules, inability to overcome the barrier properties of stratum corneum and many more. Transfersomes hold great prospective in delivery of huge range of drug substances which includes large molecules like peptides, hormones and antibiotics, drugs with poor penetration due to unfavorable physicochemical characters. Formulating the drug in a transfersome is one such approach to solve these problems. Transfersome is an ultradeformable vesicle, elastic in nature which can squeeze itself through a pore which is many times smaller than its size owing to its elasticity. Present article deals with the properties of transfersomes, method of preparation and different evaluation parameters.

Keywords: First pass metabolism, transdermal route, transfersome.

Article Info: Received 29 August 2017; Revised 5 September; Accepted 9 September, Available online 15 September 2017



Cite this article-

Chauhan N, Kumar K, Pant NC. An updated review on transfersomes: a novel vesicular system for transdermal drug delivery. Universal Journal of Pharmaceutical Research 2017; 2(4): 42-45.

DOI: <http://doi.org/10.22270/ujpr.v2i4.RW2>

Address for Correspondence:

Neelam Chauhan, Global Institute of Pharmaceutical Education and Research, Kashipur, Uttarakhand, India, Tel: +91-9045529093, E-mail: neelamchauhan9015@gmail.com

INTRODUCTION

Since the last few years, the vesicular systems have been promoted as a mean of sustained or controlled release of drugs. The word “transfersome” and the underlying concept were introduced in 1991 by Gregor Cevc. The name ‘Transfero’ is derived from the latin word meaning to carry across and the Greek word ‘soma’ for a body¹. A transfersome is a highly adaptable and stress-responsive, complex aggregate. Its preferred form is an ultra deformable vesicle possessing an aqueous core surrounded by the complex lipid bilayer. Vesicles are water-filled colloidal particles. The walls of these capsules consist of amphiphilic molecules (lipids and surfactants) in a bilayer conformation². These vesicles serve as a depot for the sustained release of active compounds in the case of topical formulations, as well as rate-limiting membrane barrier for the modulation of systemic absorption in the case of transdermal formulations³. Transfersomes consist of a phospholipids component along with a surfactant mixture. The ratio of individual surfactants and total amount of surfactants control the flexibility of the vesicle.

The uniqueness of this type of drug carrier system lies in the fact that it can accommodate hydrophilic, lipophilic as well as amphiphilic to drugs⁴. Transfersomes are applied in a non-occluded method to the skin and have been shown permeate through the stratum corneum lipid lamellar regions as a result of the hydration or osmotic force in the skin. Transfersomes can deform and pass through narrow constriction (from 5 to 10 times less than their own diameter) without measurable loss. Transfersomes can pass through even tiny pores (100nm) nearly as efficiently as water, which is 1500 times smaller.




Figure 1: Structural representation of one transfersome unit

They can act as a carrier for low as well as high molecular weight drugs e.g. analgesic, anesthetic, corticosteroids, sex hormone, anticancer, insulin, gap junction protein, and albumin. They protect the encapsulated drug from metabolic degradation. They act as depot, releasing their contents slowly and gradually. They can be used for both systemic as well as topical delivery of drug. Easy to scale up, as procedure is simple, do not involve lengthy procedure and unnecessary use or pharmaceutically unacceptable additives⁵.

Advantages

1. High entrapment efficiency, for lipophilic drug it is near to 90%.
2. Can encapsulate both hydrophilic and lipophilic moieties.
3. Suitable as a carrier for low as well as high molecular weight drugs e.g., analgesic, corticosteroids, hormones, anticancer drugs, insulin, proteins, etc⁶.
4. Can deform and pass through narrow constriction (from 5 to 10 times less than their own diameter) without measurable loss.
5. Suitable for both systemic as well as topical delivery of drug⁷.
6. Protect the encapsulated drug from metabolic degradation⁸.
7. Biodegradability and lack of toxicity⁹.

Limitations

1. Chemically unstable, highly susceptible to oxidative degradation.
2. Formulations are expensive¹⁰.

Mechanism of penetration of transfersomes

After penetration through the outermost skin layers, transfersomes reach the deeper skin layer. From there, they are normally washed out into the blood circulation. If it is applied under suitable conditions, resulting in access to all body tissues¹¹.




Figure 2: Penetration pathway of transfersomes

The mechanism for penetration includes generation of “osmotic gradient” due to evaporation of water while applying the transfersomes on the skin surface. The transport of these elastic vesicles is thus independent of concentration. This osmotic gradient is developed due to the skin penetration barrier, prevents water loss through the skin and maintains a water activity difference in the viable part of the epidermis. As the vesicles are elastic, they can squeeze through the pores in stratum corneum (though these pores are less than one-tenth of the diameter of vesicles). Transfersomes by enforcing its own route induce hydration that widen

the hydrophobic pores of skin, through the widen pores there is gradual release of drug occurs that binds to targeted organ. Transfersomes act as penetration enhancers that disrupt the intercellular lipids from stratum which ultimately widens the pores of skin and facilitate the molecular interaction and penetration of system across skin¹².

Additives and methods for preparation of transfersome

Transfersomes composed of phospholipids like phosphatidyl choline which self assembles into lipid bilayer in aqueous environment and closes to form vesicle. A bilayer softening component (such as a biocompatible surfactant or an amphiphile drug) is added to increase lipid bi layer flexibility and permeability. This second component is called as edge activator¹³.

1. Vortexing-sonication method

In this method, mixed lipids (i.e. phosphatidylcholine, EA and the therapeutic agent) are blended in a phosphate buffer and vortexed to attain a milky suspension. The suspension is sonicated, followed by extrusion through poly-carbonate membranes¹⁴.

2. Suspension homogenization process

In this process, transfersomes are prepared by mixing an ethanolic soybean phosphatidylcholine solution with an appropriate amount of edge-active molecule, e.g. sodium cholate. This prepared suspension is subsequently mixed with Triethanolamine-HCl buffer to yield a total lipid concentration. The resulting suspension is sonicated, frozen, and thawed for 2 to 3 times¹⁵.

3. Modified handshaking process

In this process, the transfersomes are prepared by modified hand shaking, ‘lipid film hydration technique’. Drug, lecithin (PC) and edge activator were dissolved in ethanol: chloroform (1:1) mixture. Organic solvent was removed by evaporation while hand shaking above lipid transition temperature (43°C). A thin lipid film was formed inside the flask wall with rotation. The thin film was kept overnight for complete evaporation of solvent. The film was then hydrated with phosphate buffer (pH 7.4) with gentle shaking for 15 minute at corresponding temperature¹⁵.

4. Aqueous lipid suspension process

In this process, Drug-to-lipid ratio in the vehicles is fixed between 1/4 and 1/9. Depending upon the particular formulation type, the composition is preferred. This would ensure the high flexibility of the vesicle membrane in comparison to the standard phosphatidylcholine vesicles in the fluid phase. Specifically, vesicles with the size ranging from 100-200 nm are prepared by using soyphosphatidylcholine with the standard deviation of size distribution (around 30%). This formulation could be prepared by suspending the lipids in an aqueous phase wherein the drug is dissolved¹⁶.

5. Centrifugation process

In this process, phospholipids, surfactants and the drug are dissolved in alcohol. Then the solvent is removed by rotary evaporation under reduced pressure at 40°C. Final traces of solvent are removed under vacuum. Then the deposited lipid film is hydrated with the

appropriate buffer by centrifuging at 60 rpm for 1 hour at room temperature. At room temperature, the resulting vesicles are swollen for 2 hours. The multi-lamellar lipid vesicles obtained which are further sonicated at room temperature¹⁶.

CHARACTERIZATION AND EVALUATION OF TRANSFERSOMES

1. Determination of vesicle diameter

The vesicle size is one of the key issues during the manufacturing process of transfersomes. It gives important information about the control of the preparation technique and can be utilized for process optimisation. Very small vesicles (smaller than 40 nm) are prone to fusion processes due to the high curvature of their bilayer membrane. It can be determined using photon correlation spectroscopy or dynamic light scattering (DLS) method¹⁷.

2. Determination of Vesicle shape and type

Transfersomes vesicles can be visualized by TEM, Phase contrast microscopy, etc¹⁸.

3. Determination of vesicle size distribution and zeta potential

Vesicle size, size distribution and zeta potential were determined by Dynamic Light Scattering Method (DLS) using a computerized inspection system by Malvern Zetasizer¹⁸.

4. Determination of number of vesicle per cubic mm

This is an important parameter for optimizing the composition and other process variables. Non-sonicated transfersome formulations are diluted five times with 0.9% sodium chloride solution. Haemocytometer and optical microscope can then be used for further study¹⁸.

5. Determination of entrapment efficiency

The entrapment efficiency is expressed as the percentage entrapment of the drug added. Entrapment efficiency can be determined by separating the untrapped drug. After centrifugation (to separate the untrapped drug), the vesicle can be ruptured¹⁹.

6. Determination of drug content

The drug content can be determined using one of the instrumental analytical methods such as modified high performance liquid chromatography method (HPLC) method using a UV detector, column oven, auto sample, pump, and computerized analysis program depending upon the analytical method of the pharmacopoeial drug¹⁹.

7. Turbidity measurement

Turbidity of drug in aqueous solution can be measured using nephelometer²¹.

8. Surface charge and charge density

Zetasizer is used to determine surface charge and charge density of transfersomes. Surface charge and Charge density of transfersomes can be determined using Zetasizer²¹.

9. Confocal scanning laser microscopy study²²

In this technique lipophilic fluorescence markers are incorporated into the transfersomes and the light emitted by these markers used for following purpose:

- a) Investigation of the mechanism of penetration of transfersomes across the skin.

- b) Determination of histological organization of the skin, shapes and architecture of the skin penetration pathways.

- c) Comparison and differentiation of the mechanism of penetration of transfersomes with liposomes, niosomes and micelles.

10. *In-vitro* drug release

In vitro drug release study is performed for determining the permeation rate. For determining *in vitro* drug release, beaker method is used in which transfersomes suspension is incubated at 32°C using cellophane membrane and the samples are taken at different times and then detected by various analytical techniques (UV, HPLC, HPTLC) and the free drug is separated by minicolumn centrifugation, then the amount of drug release is calculated²³.

APPLICATION OF TRANSFERSOMES

1. Delivery of insulin

Insulin is generally administered by subcutaneous route that is inconvenient. Encapsulation of insulin into transfersomes (transfersulin) overcomes the problems of inconvenience, larger size (making it unsuitable for transdermal delivery using conventional method) along with showing 50% response as compared to subcutaneous injection²⁴.

2. Delivery of corticosteroids

Transfersomes improves the site specificity and overall drug safety of corticosteroid delivery into skin by optimizing the epicutaneously administered drug dose. Transfersomes based corticosteroids are biologically active at dose several times lower than the currently used formulation for the treatment of skin diseases²⁴.

3. Delivery of proteins and peptides

Transfersomes have been widely used as a carrier for the transport of proteins and peptides. Proteins and peptides are large biogenic molecules which are very difficult to transport into the body, when given orally they are completely degraded in the GI tract and transdermal delivery suffers because of their large size²⁵.

4. Delivery of anticancer drugs

Anti cancer drugs like methotrexate were tried for transdermal delivery using transfersome technology. The results were favorable. This provided a new approach for treatment especially of skin cancer²⁵.

5. Delivery of anesthetics

Transfersome based formulations of local anesthetics-lidocaine and tetracaine showed permeation equivalent to subcutaneous injections, with less than 10 min. Maximum resulting pain insensitivity is nearly as strong (80%) as that of a comparable subcutaneous bolus injection, but the effect of transfersosomal anesthetics last longer.

6. Delivery of herbal drugs

Transfersomes can penetrate stratum corneum and supply the nutrients locally to maintain its functions resulting maintenance of skin²⁵.

CONCLUSION

Transdermal drug delivery system has several advantages but there is a major limitation, transportation

of the larger size molecule. That is why vesicular system like transfersomes are developed to overcome these limitations. This carrier system does not depend upon the concentration gradient and mainly works on the principle of hydrotaxis and elasto-mechanics. Transfersomes are highly deployed in the delivery of hormones, proteins, anticancer drugs, anesthetics and insulin transdermally. Transfersomes hold great prospective in delivery of huge range of drug substances which includes large molecules like peptides, hormones and antibiotics, drugs with poor penetration due to unfavorable physicochemical characters. All above discussed properties of this technology conclude its good future in transdermal drug delivery.

REFERENCES

- Honeywell-Nguyen PL, Bouwstra JA. Vesicles as a tool for transdermal and dermal delivery. *Drug Discovery Today: Technologies* 2005; 2(1): 67-74. <https://doi.org/10.1016/j.ddtec.2005.05.003>
- Hofl HEJ, Bouwstra JA, Spies F, Gooris G, Nagelkerke JF. Interaction of liposomes and niosomes with human skin. *J Pharm Sci* 1994; 83, 1192-1196. [https://doi.org/10.1016/0169-409X\(95\)00076-J](https://doi.org/10.1016/0169-409X(95)00076-J)
- Fang YP, Tsai YH, Wu PC, Huang YB. Comparison of 5-aminolevulinic acid-encapsulated liposome versus ethosome for skin delivery for photodynamic therapy. *Int J Pharm* 2008; 356 (1-2): 44-52. <https://doi.org/10.1016/j.ijpharm.2008.01.020>
- Prajapati ST, Patel CG, Patel CN. Transfersomes: A vesicular carrier system for transdermal drug delivery, *Asian J Biochem Pharm Res* 2011; 2 (1), 507-524. <https://doi.org/10.3797/scipharm.1208-02>
- Kombath RV, Minumula SK, Sockalingam A, Subadhra S, Parre S, Reddy TR, David B. Critical issues related to transfersomes- novel Vesicular system. *Acta Sci Pol Technol Aliment* 2012; 11 (1), 67-82.
- Walve JR, Bakliwal SR, Rane BR, Pawar SP. Transfersomes: A surrogated carrier for transdermal drug delivery system. *Int J App Biol Pharm Tech* 2011; 2 (1): 201-214.
- Modi CD, Bharadia PD. Transfersomes: New dominants for transdermal drug delivery. *American J Pharmtech Res* 2012; 2 (3): 71-91.
- Sheo DM, Shweta A, Ram CD, Ghanshyam M, Girish K, Sunil KP. Transfersomes- a novel vesicular carrier for enhanced transdermal delivery of stavudine: development, characterization and performance evaluation. *J Scientific Spec Res* 2010; 1 (1): 30-36. <https://doi.org/10.3797/scipharm.1208-02>
- Patel R, Singh SK, Singh S, Sheth NR, Gendle R. Development and characterization of curcumin loaded transfersome for transdermal delivery. *J Pharm Sci Res* 2009; 1 (4): 71-80.
- Shivanand P, Manish G, Viral D, Jarina F. Transfersomes: A novel approach for transdermal drug delivery. *Scholars Research Library Der Pharmacia Lettre* 2009; 1(2): 143-50.
- Gregor C, Dieter G, Juliane S, Andreas S, Gabriele B. Ultra-flexible vesicles, Transfersomes, have an extremely low pore penetration resistance and transport therapeutic amounts of insulin across the intact mammalian skin. *Biophysica Acta* 1998; 1368: 201-215. [https://doi.org/10.1016/S0005-2736\(97\)00177-6](https://doi.org/10.1016/S0005-2736(97)00177-6)
- Maghraby EI, Williams GM, Barry BW. Skin delivery of oestradiol from lipid vesicles: importance of liposome structure. *Int J Pharmaceutics* 2000; 204: 1-2, 159-69. [https://doi.org/10.1016/S0378-5173\(00\)00493-2](https://doi.org/10.1016/S0378-5173(00)00493-2)
- Trotta M, Peira E, Carlotti ME, Gallarate M. Deformable liposomes for dermal administration of methotrexate. *Int J Pharmaceutics* 2004; 270: 119. <https://doi.org/10.1016/j.ijpharm.2003.10.006>
- Woo HJ, Carraro C, Chandler D. Assembly of extended interfaces and micelles: Charge frustrated models of amphiphilic mixtures. *Faraday Discuss* 1996; 104: 183-191. <https://doi.org/10.1039/FD9960400183>
- Zhdanov RI, Podobed OV, Vlassov VV. Cationic lipid-DNA complexes-lipoplexes-for gene transfer and therapy. *Bioelectrochemistry*. 2002; 58: 53. [https://doi.org/10.1016/S1567-5394\(02\)00132-9](https://doi.org/10.1016/S1567-5394(02)00132-9)
- Strittmatter P, Enoch HG. Formation and properties of 1000-A-diameter, single-bilayer phospholipid vesicles. *Methods Enzymol* 1978; 52: 188-193.
- Sullivan SM, Huang L. Enhanced delivery to target cells by heat-sensitive immunoliposomes. *Proc Natl Acad Sci*. 1986; 83: 6117. <https://doi.org/10.1073/pnas.83.16.6117>
- Lum K, Chandler D, Weeks JD. Hydrophobicity at small and large length scales. *J Phys Chem B* 1999; 103: 4570-4577. <https://doi.org/10.1021/jp984327m>
- Cevc G, Blume G, Schatzlein A. Transfersomes-mediated transperidermal delivery improves the regio-specificity and biological activity of corticosteroids *in vivo*. *J Control Release* 1997; 45:211-226. [https://doi.org/10.1016/S0168-3659\(96\)01566-0](https://doi.org/10.1016/S0168-3659(96)01566-0)
- Cevc G, Gebauer D. Hydration-driven transport of deformable lipid vesicles through fine pores and the skin barrier. *Biophysical J* 1984; (4): 1010-1024. [https://doi.org/10.1016/S0006-3495\(03\)74917-0](https://doi.org/10.1016/S0006-3495(03)74917-0)
- Irfan M, Verma S, Ram A. Preparation and Characterization of Ibuprofen loaded transfersomes as a novel carrier for transdermal drug delivery. *Asian J Pharm Clin Res* 2012; 5(3): 162-165. <https://doi.org/10.3797/scipharm.1208-02>
- Kumar K, Rai AK. Proniosomal formulation of Curcumin having Anti-inflammatory and Anti-arthritis activity in different experimental animal models. *Die Pharmazie*. 2012; 10:852-857. <https://doi.org/10.1691/ph.2012.1164>
- Nanda A, Nanda S, Dhall M, Rao R. Transfersomes-a novel ultradeformable vesicular carrier for transdermal drug delivery. *Transdermal Delivery* 2005; 5 (9). <https://doi.org/10.4103/2231-4040.85524>
- Doaa A, Dalia R, Sally H and Mohamed N. Formulation and evaluation of fluconazole topical gel. *Int J pharm pharmaceutical sciences* 2012; 4: 176-183.
- Merin P. Eldhose, Flowerlet M, Neethusha J. Transfersomes - a review. *Int J Pharm Pharm Sci* 2016; 6(4): 436-452.